

Review

## AN OVERVIEW OF THE METHOD, MANAGEMENT, PROBLEM AND THEIR SOLUTION IN THE PEARL OYSTER (*Pinctada margaritifera*) CULTURE

Ludi Parwadani Aji

Technical Implementation Unit For Marine Life Conservation, Research Centre For Oceanography,  
Indonesian Institute of Science (LIPI), Biak – West Papua

Received : March, 7, 2011 ; Accepted : May, 31, 2011

### ABSTRACT

*Pearl culture operations can be divided into three categories which are collection/hatchery production, on-growing and pearl culture. For the hatchery, the pearl oyster industry relies on spat collection at natural production atolls where spat is abundant during the warm season and also from broodstock in the laboratory condition. After that, hatchery grown juveniles are put into the sea on the material which they settle upon. The spat are left to grow for 2 years till an average size of 90 mm. Pearl culture involves the implantation of a spherical nucleus together with a piece of mantle tissue (Saibo) from a sacrificial oyster, into the gonads. Although pearl culture is extensive with little control over weather, the use of good management methods can drastically increase productivity and result in higher profitability. Therefore, management of culture system such as site selection, settlement, feeding, stocking density and pearl culture technique is essential. For example, site selection is the most critical factor affecting pearl oyster productivity and spat collection, as the oysters spend most of their growing time exposed to water elements. Site selection must take into account important water quality parameters like temperature, salinity and turbidity. Moreover, it was identified several problems in pearl oyster culture including predation, disease and biofouling. They can result in massive loss in productivity. However, pearl industries have solution to deal with those problems. For instance, it can be done by cleaning mesh bag, biofouling organisms and pearl oyster regularly. For the future, the genetic approach like to create faster growing oysters, resistance to diseases and production of higher quality pearls has given promising results. Therefore, the productivity of pearl oyster can be improved.*

**Keywords:** pearl oyster ; management ; culture ; *Pinctada margaritifera*

**Correspondence:** Phone +62-98-181112; E-mail: [ludi\\_bio@yahoo.co.id](mailto:ludi_bio@yahoo.co.id)

### INTRODUCTION

*Pinctada margaritifera* is distinguished by the black color of the outer surface of the shell and non-nacreous border. The silver nacre inside the shell is dark towards the distal rim, hence the name black lip (Saville-Kent, 1983). The black lip pearl oyster ranges from Gulf of California, Mexico to the Eastern Mediterranean sea (George, 1978), from French Polynesia to Cook Islands, and across the northern coast of Australia from champion bay in Western Australia to Moreton bay in Queensland. Aquaculture of the species began in 1961 after the French government started an experimental study on the species in collaboration with

Nippo Pearls Co. of Japan in Tahiti. The pearl industry grew steadily and by 1998, black-lipped cultured pearls were cultured largely in Tahiti, Cook Islands, Fiji, Marshall Islands and Japan, with Tahiti dominating the market with 95% of total production (USD 143 million) and contributing to 25% of the total pearl production in the world (Hisada and Fukuhara, 1999). This essay will examine methods of culture of the pearl oyster and methods of management across different species which will result in maximum productivity and maximum profits. Moreover, the problem of Pearl oyster


culture is also reviewed by giving some solution to deal with that obstacle.

## METHOD OF CULTURE

Pearl culture operations can be divided into three categories which are (1) collection/hatchery production, (2) ongrowing and (3) pearl culture. Each phrase of production permits a degree of specialization by farmer and allows people of different income and different technical expertise to get involved in cultivation.

### *Collection/ hatchery production*

In French Polynesia, the pearl oyster industry relies on spat collection at natural production atolls where spat is abundant, during the warm season (November to May). Farmers place their collectors, made of black shade mesh and suspended from a longline every 30 cm at a depth of 3 m (Cabral *et al.*, 1985). Spat oysters are left on collectors (**Fig. 1**) for 1 year till 4-5cm in length though some farmers may choose to have them for a year and a half or till they reach 14cm, where they are big enough to be grafted, and skip the ongrowing process altogether. (Le Moullac *et al.*, 2003).



**Fig. 1.** Spat pearl oyster attached to spat collector (Southgate and Lucas, 2008)

In Australia, however, some of the spat are produced in hatcheries, where broodstock are first conditioned on  $50-100 \times 10^3$  cells/ml of a 1:1 mixture of *Isochrysis galbana* and *Chaetoceros simplex* for 6 weeks. To induce spawning, oysters are held overnight at  $20^\circ\text{C}$ , and then transferred to a raceway with the seawater temperature at  $30-32^\circ\text{C}$ . The first spawning male triggers spawning by other oysters and this occur within a few minutes to 1 hour (Southgate and Beer, 1997). Spawning oysters are removed individually into individual containers. Larvae are reared in static water exchange systems where water is changed every 2-3 days. *Isochrysis galbana* and *Pavlova lutheri* were given once a day concentrations from 5 - 15 cells/ $\mu\text{l}$  until metamorphosis (Doroudi *et al.*, 1999). At day 20, when most of

them reach  $150\mu\text{m}$ , they are transferred to seawater settlement tanks.

### *On growing*

Hatchery grown juveniles are put into the sea on the material which they settled upon. The material is hung from longlines in areas of calm water (**Fig. 2**). Longline are horizontal ropes which are supported by floats and anchored at the final part of ropes to maintain position. A protective mesh covers the material, and mesh size is increased as spat grow. The spat are left to grow for 2 years till an average size of 90 mm. A hole is then drilled through the posterior ear and the oysters are hung down in pairs on a down line.



**Fig. 2.** Longline used for Pearl oyster culture. Note the panel nets containing pearl oyster juveniles suspended from the longline (Southgate and Lucas, 2008)

### *Pearl culture*

Pearl culture involves the implantation of a spherical nucleus together with a piece of mantle tissue (Saibo) from a sacrificial oyster, into the gonads. The mantle tissue grows around the nucleus and secretes nacreous deposits to form a pearl. At a minimum size of 100mm, the preoperative pearl oyster undergoes a weakening process for 40 days for the muscular and gonadal epithelium to disintegrate. They are either crowded together, limiting the oxygen and food supply or moved to deeper water and then moved up the water column to create stress, which causes the oyster to spawn.

Implantation takes place during the cooler months, when the conditioned oysters are wedged open, an incision is made in the gonads and the nucleus and Saibo is added. The wound is cleaned with antiseptic. The oysters are then returned to the sea and placed in calm conditions. Cells in the pearl sac secrete nacre onto the outer surface of the nucleus and cultured pearls are harvested when nacre are 1 mm thick after 18-24 months (Gervis and Sims, 1992).

### **MANAGEMENT METHODS**

Although pearl culture is extensive with little control over weather, the use of management methods can drastically increase productivity and result in higher profitability. Therefore, management of culture system such as site

selection, settlement, feeding, stocking density and pearl culture technique is essential.

### *Site selection*

Site selection is the most critical factor affecting pearl oyster productivity and spat collection, as the oysters spend most of their growing time exposed to water elements. Site selection must take into account important water quality parameters like temperature, salinity and turbidity. Temperature affects transparency level, absorption efficiency of suspension feeding, respired energy and excreted energy. Temperature was found to significantly increase respiratory energy of *P. margaritifera* at high temperatures reducing potential for growth (Yukihira *et. al.*, 2000). Compared to *P. maxima* (32 °C), optimum growth for *P. margaritifera* occurred at 23-28 °C and is therefore suitable for cooler climates. High temperatures were also found to increase oxygen consumption and ammonia excretion in another species *P. mazatlanica* (Saucedo *et. al.*, 2004). The effect of ammonia excretion can be mitigated through co cultivation with red alga *Kappaphycus alvarezii*, which can remove nitrogenous waste and CO<sub>2</sub> released by oysters and provide O<sub>2</sub> from photosynthesis (Wu *et al.*, 2003). Temperature also has a pronounced effect on larval development in nursery culture. At 18-26°C, *P. imbricata* increased rate and number of embryo development into D-stage veliger with increasing temperature, under 14 °C, larvae did not develop into veliger.

Temperature also affects byssal attachment, with the highest rate at 18°C (O'Connor and Lawler, 2004). Most importantly, temperature controls quality of nacre. Higher temperatures accelerate nacre deposition, however, lower temperatures decrease the deposition rates but increasing the quality of the nacre (Shirai, 1970). Colder temperatures have a positive influence on lustre (O'Sullivan *et al.*, 1998). Therefore, seeding of pearls is always done at the coldest times of the year.

Salinity also plays an important role in larvae development. Survival of *P. imbricata* larvae was highest at 32 and 35 ppt and at 29-35ppt, percentage of embryo developing to D-stage larvae increases significantly with increasing salinity (O'Connor and Lawler, 2004). Turbidity also has an influence on growth and survival of oysters. Turbidity is caused by suspended particulate matter (SPM), and is inversely proportional to organic particulate matter (Yukihira *et al.*, 1999), of which oysters feed from. *P. margaritifera* do not survive and grow well in areas of high SPM (Yukihara *et al.*, 2006). Furthermore, *P. margaritifera* has developed a trophic strategy involving clearance of large amount of water to gain sufficient nutrients in nutrient poor water (Pouvreau *et al.*, 2000), and this process is energy consuming and reduces the scope of growth.

### Settlement

A major part of a successful pearl venture depends on collecting maximum numbers of spat, both in the hatchery and from the sea. Settlement of larval spat is affected by various factors. Firstly, use of natural biofilm induces settlement in *P. maxima* larvae (Zhao *et al.*, 2003). Types of substrate also influence larvae settlement. Deep colour material like red and blue attracted significantly more larvae from *P. martensii* than lighter coloured green and yellow material. In addition, rougher material also promoted more settlement, so does coating of substrate with tissue extract from the same species (Su *et al.*, 2007).

Depth and chemical cues also affect spat recruitment, with an intermediate depth of 60-90cm collecting the most *P. Mazatlanica* (Saucedo *et al.*, 2005). High settlement of larvae occurred in the presence of GABA at concentrations of  $10^{-3}$  and  $10^{-4}$  M. However,

mortality rate was also high (32%) at a concentration of  $10^{-3}$  M (Doroudi and Southgate, 2002), suggesting unsuitability of use. Other less invasive chemicals include excess potassium as well as 50mM calcium chloride (Zhao *et al.*, 2003b).

In hatcheries, larvae tend to settle on side of settlement tanks and removal is by scraping (Rose and Baker, 1994) which is time consuming and may damage shells. Exposing the oysters to hypersaline conditions of 45 ppt for one hour or lowering the pH of water to 4 can cause voluntarily detachment of spat from substrate without any mortality (Taylor *et al.*, 1997a), saving time in the process. This also has implications for transport of spat, as spat will detach if transported under conditions that cause them to dehydrate.

### Feeding

Feeding of spat cannot be controlled for farming in the sea. However, survival of larvae and juvenile oysters in hatchery tanks requires management of feed. Optimum diet for *P. margaritifera* larvae was found to be 1:1 mixture of *Pavlova salina* and *Chaetoceros simplex* and dried *Tetraselmis suecica* and commercial yeast L-10, Microfeast (Southgate *et al.*, 1998). Recent results showed that addition of another diatom *C. muelleri* resulted in better growth for D and umbone stage larvae (Martinez-Fernandez and Southgate, 2007). Apparently, microalgae diets vary with species, as *Isochrysis aff. galbana* (T-ISO) and *Pavlova lutheri* provide for optimum growth of *P. sterna* larvae (Martinez-Fenandez *et al.*, 2004).

### Stocking density

Stocking density has been found to affect larvae, juvenile and spat growth of oysters. Taylor *et al.*, (1997b) found the maximum rate of survival and growth in terms of wet weight, shell length and height of *P. maxima* juveniles at 1.3 juvenile per 100cm<sup>3</sup>. Higher stocking density tends to affect water flow, which directly affects food availability. Under prolonged low food availability, oysters partition energy into increasing shell weight and thickness over body tissue growth (Wilson, 1987). This effect is also determined by stage of culture. Juveniles of *Pteria sterna* are better able to withstand higher stocking density. An

optimum protocol has been worked out to start nursery culture at more than 100 individuals/tray and perform density reduction until there are 100 individuals/tray after 3 months. Then reduce individuals to 50/tray from the 4<sup>th</sup> month onwards (Monteforte *et al.*, 2005). Moreover, self-thinning will also occur in oyster populations at high density (Frechette *et al.*, 1996). For juveniles on suspended box culture, a 30% stocking density is the optimum method for culturing *P. margaritifera* (Southgate and Beer, 2000).

### *Pearl culture techniques*

During seeding operation, high numbers of oysters will reject the nucleus and the rest will produce poor quality pearls. Time of implantation is of importance. As the oyster grows, there is a progressive reduction in growth rate. This implies that the sooner the nucleus is implanted, the greater is the rate of nacreous deposition, and the shorter is the time to obtain a marketable pearl (Pouvreau *et al.*, 2000). Implantation of pearl when temperatures are low also resulted in high quality mabe pearls of the oyster *P. sterna* (Ruiz-Rubio *et al.*, 2006).

Use of a relaxant will allow relaxation of abductor muscle and opening of shell valve so that the muscles will not be damaged from overstretching during operation. It also decreases bead rejection. Propylene phenoxetol is effective at concentrations of 2.0 to 3.0 ml/l result in rapid induction and 100% recovery and survival after 7 days (Norton *et al.*, 1996). Use of 1:50 aqueous dilution of 10% povidone iodine antiseptic also reduces mortality and promotes recovery. Pearl shape was improved when the surgical incision was closed with a flexible cyanoacrylate adhesive (Norton *et al.*, 2000).

Anaesthetic can also enable donor oysters to recover their mantle tissue after excision. Therefore, donor oysters need not be sacrificed and genetic selection can take place. *P. fukata* can grow the excised tissue within 105 hours (Acosta-Salmón and Southgate, 2006) and *P. margaritifera* when exposed to 1200 mg/l of benzocaine resulted in 30 minute relaxation time and rapid recovery with no mortality after operation (Acosta-Salmón *et al.*, 2005).

## **PROBLEM AND THEIR SOLUTION IN PEARL OYSTER CULTURE**

### *Predation*

Predation of oyster spat and juveniles can result in massive loss in productivity. Predators include excavating sponges of the family *Clionaidae* (*Porifera: Demospongiae*) (Fromont *et al.*, 2005), gastropods *Cymatium* (Family: Ranellidae), and Crustacea. There are some ways to deal with the problem. Firstly, spat can be removed from collectors at a shorter period 3-4 months instead of 6 months, when mean size is 15mm DVM, and reared in panel nets (Friedman and Bell, 2000). Alternatively, the spat can be covered with mesh, and a mesh size of 3 mm improved spat survival of *P. maxima* while not fouling so easily (Taylor *et al.*, 1998). However, in environments like Solomon Islands, mesh bags should not be used, as they trap larvae of predators such as *Cymatium* spp, gastropods and portunid crabs and fouled easily, impeding water flow to the collector (Friedman and Bell, 1996). For juvenile culture, ear hanging and 24 pocket juvenile panel nets were the preferred method, showing highest growth and lowest fouling and mortality from predators (Southgate and Beer, 2000).

### *Biofouling*

Removal of biofouling organisms is done by manual labour (**Fig. 3**) and contributes to most of the expenses of a pearl oyster farm. *P. maxima* left uncleaned for 4 weeks or more had stunted growth and shell deformities were observed after 8 weeks, caused by biofouling organisms like barnacles and polychaete worms (Taylor *et al.*, 1997). However, cleaning also affect survival rates, and frequent cleaning every 4 weeks resulted in high mortality for *P. margaritifera*. Handling oysters will stress them and therefore optimum cleaning rate for this species is once every 8 weeks (Pit and Southgate, 2003). Age of oyster may also affect biofouling rate as the periostracum, a thin, flexible, fibrous and sclerotized protein layer covering the calcified shell functioning as a physical defence against fouling organism, wears off over time (Scardino *et al.*, 2003). A novel way of dealing with biofouling involves painting them with anti biofouling chemicals

but as evident in the research done by de Nys and Ison (2000), the coating only last for 3 months in the northern territory and therefore it

is more economical to perform manual cleaning of oysters.



**Fig. 3.** Manual removal of fouling from pearl oyster culture nets using knives (Southgate and Lucas, 2008).

### Diseases

Oyster diseases have the potential to affect productivity. Pearl oyster *P.maxima* suffered high mortality in Australia in 1996 due to the bloom of algae *Trichodesmium erythraeum*. The oyster exhibit dilated digestive gland lumens, sloughing of epithelial cells and large numbers of residual bodies and with increased numbers of brown cells and granulocytes under the epithelial layer, indicative of an inflammatory response. This is due to the algae producing a paralytic shellfish toxin, saxitoxin (Negri *et al.*, 2004) and this cannot be prevented. However, another commonly occurring oyster disease, the reddening of abductor muscle caused by infection in *P. fucata* can be prevented by lowering the temperature of water to less than 19 °C (Nagai *et al.*, 2007).

### Genetics

The future of pearl oyster aquaculture lies in genetic selection, which can create faster growing oysters, resistance to diseases and

production of higher quality pearls. This first step in doing so is to ensure quality gametes from selected broodstock are kept at their optimum conditions. Crytopreservation of sperm can ensure the preservation of desirable genes and good motility of sperms. For *P. fucata*, 10% METmethanol plus 90% diluents comprising 80% seawater and 20% FBSfetal bovine serum, cooled to 50 °C at a cooling rate between 15 and 20 °C/min is a good method of crytopreservation (Kawamoto *et al.*, 2007). *P. margaritifera* has a different method of crytopreservation- 0.45 M trehalose with 0, 0.64, 1.02 or 1.53 M Me2SO and frozen using slow to medium-rapid cooling rates (2.1–5.2 °C/min) (Acosta-Salmon *et al.*, 2007). When needed, sperms can be activated using 2.0 mM NH<sub>3</sub> solution (Ohta *et al.*, 2007).

Selection in *P. fucata* has already shown effect. By selectively crossing white male and female oysters, frequency of inferior yellow coloured pearls was reduced (Wada and Komaru, 1996). One of the worries of selection through inbreeding is loss in genetic diversity, which may lead to inferior growth and survival (Wada and Komaru, 1994). Fortunately, in

French Polynesia, no evidence has been found to indicate a change in the number of alleles and heterozygosity for current farming methods (Arnaud-Haond *et al.*, 2003). Caution needs to be exercised to prevent genetic loss from happening in future.

## CONCLUDING REMARKS

Pearl oyster cultures have several constraints that can affect their productivity such as predation, biofouling and disease. On the other hand, pearl farming has gone a long way in establishing control of production through various management methods. The future scope for expansion in pearl farming is in genetics, where more valuable quality pearls can be produced in a shorter time in higher quantity and pearl oysters can improve resistance to diseases. However, genetics is a double edged sword and future management decisions must ensure the preservation of genetic diversity as a buffer against unforeseen circumstances. Therefore, for the future research which improves pearl oyster production especially through genetic approach is important.

## REFERENCES

- Acosta-Salmo'n, H., E. Mart'inez-Ferna'ndez, and P.C. Southgate, 2005. Use of relaxants to obtain saibo tissue from the blacklip pearl oyster (*Pinctada margaritifera*) and the Akoya pearl oyster (*Pinctada fucata*), *Aquaculture*, 246: 167– 172
- Acosta-Salm' on, H., and P.C. Southgate, 2006. Wound healing after excision of mantle tissue from the Akoya pearl oyster, *Pinctada fucata*, *Comparative Biochemistry and Physiology, Part A*, 143: 264–268
- Acosta-Salmo' n, H., D.R. Jerry, and P.C. Southgate, 2007. Effects of cryoprotectant agents and freezing protocol on motility of black-lip pearl oyster (*Pinctada margaritifera* L.) spermatozoa. *Cryobiology*, 54: 13–18
- Arnaud-Haonda, S., V. Vonaua, F. Bonhommeb, P. Boudry, J. Proua, T. Seaman, M. Veyreta, and E. Goyard. 2003. Spat collection of the pearl oyster (*Pinctada margaritifera* cumingii) in French Polynesia: An evaluation of the potential impact on genetic variability of wild and farmed populations after 20 years of commercial exploitation, *Aquaculture*, 219: 181–192
- Cabral, P., K. Mizuno, and A. Tauru, 1985. Preliminary data on the spat collection of mother of pearl (*Pinctada margaritifera*, Bivalve, Mollusc) in French Polynesia. In: Delesalle, B., Galzin, R., Salvat, B. (Eds.), Proc. Vth Int. Coral Reef Cong. French Polynesian Coral Reefs, Antenne Museum-EPHE, Moorea, French Polynesia, 1: 177– 182.
- De Nys, R. and O. Ison, 2004. Evaluation of antifouling products developed for the Australian pearl industry, FRDC project no. 2000/254, James Cook University, Queensland, Australia, 114 pp.
- Doroudi, M.S. and P.C. Southgate, 2002. The effect of chemical cues on settlement behaviour of blacklip pearl oyster (*Pinctada margaritifera*) larvae, *Aquaculture*, 209: 117–124
- Doroudi, M.S., P.C. Southgate, and R.J. Mayer, 1999. The combined effect of temperature and salinity on embryos and larvae of the black-lip pearl oyster, *Pinctada margaritifera* (L.), *Aquaculture. Res.*, 30: 271– 277
- Friedman, K.J. and J.D. Bell, 1996. Effects of different substrata and protective mesh bags on collection of spat of the pearl oysters, *Pinctada margaritifera* (Linnaeus, 1758) and *Pinctada maculata* (Gould, 1850), *J. shellfish Res.*, 15(3): 535-541
- Friedman, K.J. and J.D. Bell, 2000. Shorter immersion times increase yields of the blacklip pearl oyster, *Pinctada margaritifera* Linne., from spat collectors in Solomon Islands, *Aquaculture*, 187: 299–313

- Fromont, J., R., Craig, L. Rawlinson, and J. Alder. 2005. Excavating sponges that are destructive to farmed pearl oysters in Western and Northern Australia, *Aquaculture. Res.*, 36: 150-162
- George, C.D. 1978. The pearl. A report to the government of Papua New Guinea, the FAO, and the Asian development bank. Samarai, Milne Bay province, Papua New Guinea, 169 pp
- Gervis, M.H. and N.A. Sims. 1992. The biology and culture of pearl oysters, Overseas development administration of the United Kingdom, London, England, 49pp.
- Hisada, Y. and T. Fukuhara. 1999. Pearl marketing trend with emphasis on black pearl market, FAO Field Document No.13, 1999
- Kawamoto, T., T. Narita, K. Isowa, H. Aoki, M. Hayashi, A. Komaru, and H. Ohta. 2007. Effects of cryopreservation methods on post-thaw motility of spermatozoa from the Japanese pearl oyster, *Pinctada fucata martensii*, *Cryobiology*, 54: 19-26
- Martínez-Fernández, E. and P.C. Southgate. 2007. Use of tropical microalgae as food for larvae of the black-lip pearl oyster *Pinctada margaritifera*. *Aquaculture*, 263: 220-226
- Martinez-Fernandez, E., H. Acosta-Salmon, and C. Rangel-Davalos. 2004. Ingestion and digestion of 10 species of microalgae by winged pearl oyster *Pteria sterna* (Gould, 1851) larvae. *Aquaculture*, 230: 417-423
- Monteforte, M., H. Bervera, J.J. Ramirez, P. Saucedo, and C. Lopez, 2005. Effect of stocking density on growth and survival of the rainbow pearl oyster *Pteria sterna* (Gould 1852) during nursery and late culture in Bahia de La Paz, Baja California Sur, Mexico. *Aquaculture Int.*, 13: 391-407
- Moullac, G.L., E. Goyard, D. Saulnier, P. Haffner, E. Thouard, G. Nedelec, J., Goguenheim, C. Rouxel, and G. Cuzon. 2003. Recent improvements in broodstock management and larviculture in marine species in Polynesia and New Caledonia: genetic and health approaches. *Aquaculture*, 227: 89-106
- Nagai, K., J. Go. S. Segawa, and T. Honjo. 2007. A measure to prevent relapse of reddening adductor disease in pearl oysters (*Pinctada fucata martensii*) by low-water temperature culture management in wintering fisheries. *Aquaculture*, 262: 192-201
- Negria, A.P, O. Bunter, B. Jones, and L. Llewellyn. 2004. Effects of the bloom-forming alga *Trichodesmium erythraeum* on the pearl oyster *Pinctada maxima*, *Aquaculture*, 232: 91-102
- Norton, J.H., M. Dashorst, T.M. Lansky, and R.J. Mayer. 1996. An evaluation of some relaxants for use with pearl oysters. *Aquaculture*, 144: 39-52
- O'Connor, W.A. and N.F. Lawler. 2004. Salinity and temperature tolerance of embryos and juveniles of the pearl oyster, *Pinctada imbricata* Roding, *Aquaculture*, 229: 493-506
- Ohta, H., T. Kawamoto, K. Isowa, H. Aoki, M. Hayashi, T. Narita, and A. Komaru. 2007. Motility of spermatozoa obtained from testes of Japanese pearl oyster *Pinctada fucata martensii*. *Fish. sci.*, 73: 107-111
- O'Sullivan, D., R. Scoones, D. Cropp, and O. Bunter. 1998. The old and the new of the Australian pearl production, *World Aquaculture Society*, Aquaculture 1998, Book of Abstracts, Las Vegas, NV, U.S.A. Feb. 15-19, 1998
- Pit, J.H. and P.C. Southgate. 2003. Fouling and predation; how do they affect growth and survival of the blacklip pearl oyster, *Pinctada margaritifera*, during



- nursery culture. *Aquaculture Int.*, 11: 545–555
- Pouvreau, S., A. Bodo, and D. Buestel. 2000a. In situ suspension feeding behaviour of the pearl oyster, *Pinctada margaritifera*: combined effects of body size and weather-related seston composition. *Aquaculture*, 181: 91–113
- Pouvreau, S., J. Tiapari, A. Gangnery, F. Lagarde, M. Garnier, H. Teissier, G. Haumani, D. Buestel, and A. Bodoy. 2000b. Growth of the black-lip pearl oyster, *Pinctada margaritifera*, in suspended culture under hydrobiological conditions of Takapoto lagoon French Polynesia. *Aquaculture*, 184: 133–154
- Rose, R.A. and S.B. Baker. 1994. Larval and spat culture of the Western Australian silver- or gold-lip pearl oyster, *Pinctada maxima* Jameson (Mollusca: Pteriidae). *Aquaculture*, 126: 35-50
- Ruiz-Rubio H., H. Acosta-Salmón, A. Olivera, P.C. Southgate, and C. Rangel-Dávalos. 2006. The influence of culture method and culture period on quality of half-pearls ('mabé') from the winged pearl oyster *Pteria sterna*, Gould, 1851. *Aquaculture*, 254: 269–274
- Saucedo, P.E., L. Ocampo, M. Monteforte, and H. Bervera. 2004. Effect of temperature on oxygen consumption and ammonia excretion in the Calafia mother-of-pearl oyster, *Pinctada mazatlanica* (Hanley, 1856). *Aquaculture*, 229: 377–387
- Saucedo, P.E., H. Bervera-Leon, M. Monteforte, P.C. Southgate, and P. Monsalvo-Spencer. 2005. Factors influencing recruitment of hatchery reared pearl oyster (*Pinctada Mazatlanica*; Hanley 1956) spat. *J. Shellfish Res.*, 24(1): 215-219
- Saville-Kent, W. 1983. Pearl and pearl-shell fisheries, pp. 204-224. In: *The great barrier reef of Australia: its products and potentialities*. W.H. Allen and Co., London, UK
- Scardino, A., R. de Nys, O. Ison, O.V. W. Connor, and P. Steinberg. 2003. Microtopography and antifouling properties of the shell surface of the bivalve molluscs *Mytilus galloprovincialis* and *Pinctada imbricata*. *Biofouling*. 19: 221– 230
- Shirai, S. 1970. The Story of Pearls. Japan Publications Inc., Japan. 132 pp.
- Southgate, P.C. and A.C. Beer. 1997. Hatchery and early nursery culture of the blacklip pearl oyster (*Pinctada margaritifera* L.). *J. Shellfish Res.*, 16: 561– 567
- Southgate, P.C. and A.C. Beer. 2000. Growth of blacklip pearl oyster *Pinctada margaritifera* juveniles using different nursery culture techniques. *Aquaculture*, 187: 97 104
- Southgate, P.C., A.C. Beer, P.F. Duncan, and R. Tamburri. 1998. Assessment of the nutritional value of three species of tropical microalgae, dried *Tetraselmis* and yeast based diet for larvae of the blacklip pearl oyster, *Pinctada margaritifera* L. *Aquaculture*, 162: 247–257
- Southgate, P. C. and J. S. Lucas, (eds). 2008. The pearl oyster. Elsevier, Amsterdam.
- Su, Z., L. Huang, Y. Yan, and H. Li, 2007. The effect of different substrates on pearl oyster *Pinctada martensii* (Dunker) larvae settlement, *Aquaculture (In press)*
- Taylor, J.J., R.A. Rose, and P.C. Southgate, 1997a. Inducing detachment of silver-lip pearl oyster (*Pinctada maxima*, Jameson) spat from collectors. *Aquaculture*, 159: 11-17
- Taylor, J.J., R.A. Rose, P.C. Southgate, and C.E. Taylor, 1997b. Effects of stocking density on growth and survival of early juvenile silver-lip pearl oysters, *Pinctada maxima* (Jameson), held in suspended nursery culture. *Aquaculture*, 153: 41-49

- Taylor, J.J., P.C. Southgate, and R.A. Rose. 1998. Effects of mesh covers on the growth and survival of silver-lip pearl oyster *Pinctada maxima*, Jameson/spat. *Aquaculture*, 162: 241–246
- Wada, K.T. and A. Komaru. 1994. Effect of selection for shell coloration on growth rate and mortality in the Japanese pearl oyster, *Pinctada fucata martensii*. *Aquaculture*, 125: 59-65
- Wada, K.T. and A. Komaru. 1996. Color and weight of pearls produced by grafting the mantle tissue from a selected population for white shell color of the Japanese pearl oyster *Pinctada jixata martensii* (Dunker). *Aquaculture*, 142: 25-32
- Wilson, J.H. 1987. Environmental parameters controlling growth of *Ostrea edulis* L. and *Pecten maximus* L. in suspended culture. *Aquaculture*, 64: 119- 131
- Wua, M., S.K.K. Mak, X. Zhang, and P.P. Qian. 2003. The effect of co-cultivation on the pearl yield of *Pinctada martensi* (Dunker). *Aquaculture*, 221: 347–356
- Yukihira, H., D.W. Klumpp, and J.S. Lucas. 1999. Feeding adaptations of the pearl oysters *Pinctada maxima* and *P. margaritifera* to variations in natural particulates. *Mar. Ecol. Prog. Ser.*, 182: 161– 173
- Yukihira, H., J.S. Lucas, and D.W. Klumpp. 2000. Comparative effects of temperature on suspension feeding and energy budgets of the pearl oysters *Pinctada margaritifera* and *P. Maxima*. *Mar. Ecol. Prog. Ser.* 195: 179-188
- Yukihira, H., J.S. Lucas, and D.W. Klumpp. 2006. The pearl oysters, *Pinctada maxima* and *P. margaritifera*, respond in different ways to culture in dissimilar environments. *Aquaculture*, 252: 208– 224
- Zhao, B., S. Zhang, and P.Y. Qian. 2003. Larval settlement of the silver- or goldlip pearl oyster *Pinctada maxima* (Jameson) in response to natural biofilms and chemical cues. *Aquaculture*, 220: 883–901