

Original Paper

## VERTICAL CHANGES OF RECENT OSTRACODE ASSEMBLAGES AND ENVIRONMENT IN THE INNER PART OF JAKARTA BAY, INDONESIA

Lili Fauzielly<sup>1,2</sup>, Toshiaki Irizuki<sup>3</sup> and Yoshikazu Sampei<sup>3</sup>

<sup>1</sup>Department of Geoscience, Interdisciplinary Graduate School of Science and Engineering, Shimane University, 1060 Nishikawatsu-cho, Matsue 690-8504, Japan (e-mail: fzelly@yahoo.co.id)

<sup>2</sup>Faculty of Geology, Padjadjaran University, Jl Raya Bandung Sumedang Km 21 Jatinangor Sumedang, West Java 45363, Indonesia

<sup>3</sup>Department of Geoscience, Interdisciplinary Faculty of Science and Engineering, Shimane University, Japan 1060 Nishikawatsu-cho, Matsue 690-8504, Japan (e-mail: irizuki@riko.shimane-u.ac.jp; sampei@riko.shimane-u.ac.jp)

Received : Accepted :

### ABSTRACT

*A short sediment core from the inner part of Jakarta Bay, Indonesia, was quantitatively analysed for ostracods (minute Crustacea), total organic carbon (TOC) and total nitrogen (TN) contents, and the vertical distributions were recorded. A total of 53 ostracod species were obtained from 80 continuous core samples. The dominant species were Keijella carriei and Loxoconcha wrighti, which are common in areas with high TOC and TN contents. Based on an analysis of ostracod assemblages and carbon/nitrogen ratio, the study site began to be influenced by organic contamination from around 1950. Although the population of Jakarta City has increased rapidly since then, TOC and TN contents which were low, have gradually increased (0.7%–0.9% and 0.10%–0.12%, respectively), probably due to addition of nutrients from river sedimentation. The increased sedimentation rate after 1950 resulted in an increasing TOC ratio. The observed correlation between TOC and dominant species shows that Phlyctenophora orientalis may be a good indicator for monitoring increases in the narrow TOC content range of 0.7%–1.1%.*

Key word: Ostracoda, TOC, CNS analysis, Jakarta Bay

Correspondence :

### INTRODUCTION

Although the northern coastline of West Java is extensive (approximately 365,000 km), knowledge of the Recent Ostracoda from this region is sparse. Recent Ostracoda from Indonesian regions have been the subject of several studies (Brady, 1880; Kingma, 1948; Keij, 1953, 1964; Carbonel and Moyes, 1987; Dewi, 1997); however, few studies have been conducted on the northern coast of West Java; Jakarta Bay: Brady (1867–1872) and Fauzielly *et al.*, (in press); Thousand Islands: Keij (1974, 1975) and Whatley and Watson (1988).

Ostracoda is a meiobenthic class belonging to the subphylum Crustacea,

containing organisms with an average size of 1 mm living in a wide range of aquatic habitats from Ordovician to Recent (Horne *et al.*, 2002). Ostracods possess two calcified valve-like shells, which can be preserved for a long time and which are abundant in sediment cores. Several recent studies using sediment cores have revealed that ostracod species could potentially be used to reconstruct the history of bottom environments and as a proxy for organic contamination.

Many studies of ostracods in short sediment cores have been conducted in East Asia, especially in Japan (Yasuhara *et al.*, 2003,

2007; Yasuhara and Yamazaki, 2005; Irizuki et al., 2011). However, no similar studies have been conducted in southeastern Asia. In inner bays adjacent to a metropolis, such as Jakarta Bay, where humans have greatly accelerated the rate of deposition of terrestrial particles in the ocean and increased water fertility, no studies have yet been conducted that record the distribution of ostracods in relation to environmental change. Core sediments from muddy inner bays are best suited for this line of inquiry, because they provide high resolution.


The aim of this study was to reconstruct the paleoenvironments in the inner part of Jakarta Bay over the past several hundred years, based on ostracod assemblages and total organic carbon (TOC) and total nitrogen (TN) contents. This is the first study to demonstrate temporal shifts in recent ostracod assemblages in Indonesian bays in relation to organic pollution, using short sediment cores.

## MATERIALS AND METHODS

### *Study area*

Jakarta Bay is a semi-enclosed bay located on the western side of the northern part of Java Island, Indonesia. It is a shallow bay, with an average depth of approximately 15 m, an area of 514 km<sup>2</sup> and a shoreline of approximately 72 km long (**Fig. 1**). The bay is very fertile as a result of the abundant supply of nutrients from rivers that cross the city.

Human activity in and around the bay began thousands of years ago. Initially, the bay was used only for fishing, although it is probable that some shipping took place along the coast. The current harbours of Tanjung Priok and Sunda Kelapa were first used more than 300 years ago. In recent decades, many of the people living around the bay have been involved in fish and shrimp farming. The sand beaches have been dredged for landfill and thousands of hectares surrounding the bay have been transformed into fish ponds, luxury residences and industrial zones (UNESCO, 2000).



**Fig. 1.** Location of Jakarta Bay and sample localities used in this study

The population in Greater Jakarta (Jabotabek) increased rapidly from 138,600 in 1905 to 8,259,300 in 1990, an increase of almost 60-fold (pemprov DKI, 2011).


Another problem facing Jakarta is flooding. The uncontrolled growth of housing along the riverside, heavy sedimentation and the dysfunctional drainage system in Jakarta, especially in the riverside areas, make it susceptible to flooding. Jakarta has experienced extensive flooding in 1918, 1976, 1996, 2002 and 2007 (Tanuwidjaja, 2010).

The high population growth rate together with the expansion of Jakarta City during the second half of the 20th century have led to serious pollution and over-exploitation of coastal resources, threatening the sustainability of the marine environment. Jakarta Bay has

undergone some of the most drastic changes over the last few decades.

#### *Sediment collection*

The 80-cm long sediment core used in this study was collected in 1994 by the Indonesian Marine Institute using a gravity corer. The water depth at the study site is 11 m. Sediments are composed of homogeneous massive dark greyish brown (2.5Y 4/2) clay from a core depth of 1–40 cm, containing small fragments ( $\phi < 1$  mm) and olive grey (5Y 4/2) clay from a core depth of 40–80 cm, containing large fragments of molluscs. The colour and size of the shell fragments gradually changed from a depth of 45 cm (**Fig. 2**).



**Fig. 2.** Temporal changes of composition of ostracode assemblages (abundance, number of species, diversity, equitability) and total organic carbon (TOC) content (wt%), total Nitrogen (TN) content (wt%) and the ratio of TOC/TN contents. Abundance means the number of specimens per 1 gr dry sediment.

*All symbols in column shows shell fragments.*

#### *Ostracod analysis*

The core was continuously sliced in 1-cm thick samples, creating a total of 80 samples. For ostracod analysis, samples were weighed and washed through a 63- $\mu$ m sieve, and subsequently, oven dried and dry sieved to separate the >125  $\mu$ m fraction. The dried sediments were weighed and mud contents were calculated based on water content and the residual weight of the washed samples.

Samples from the >125  $\mu$ m fraction containing abundant ostracod specimens were divided using a sample splitter into separate samples, each containing approximately 200 specimens. The number of specimens refers to the sum of the left and right valves. One carapace was counted as two valves. Most specimens were composed of separate valves, and carapaces were rare.

### Grain size analysis

Grain size analysis was conducted using a laser diffraction particle size analyser (SALD-3000S) after decomposition of organic matter and pyrite with 30% H<sub>2</sub>O<sub>2</sub> for several days.

### TOC analysis

TOC and TN contents were measured using the combustion method at 1000 °C in a FISONs analyzer EA 1108, after treatment to remove the carbonate fraction. This treatment was performed by adding 1 M HCl to the weighed sediment in Ag cups.

### Results

#### Grain size

The median grain size of samples in the study area ranged from approximately 7.4 to 8.8  $\phi$  (Table 1) but most samples were less than 8  $\phi$ , suggesting that cores are mainly composed of clay. The mud content of all samples was greater than 60%, except TJ56-56 (Table 1), which contained relatively coarser particles (Md  $\phi$  = 7.6  $\phi$ ). The profile of grain size was relatively constant, but some fluctuations were apparent, particularly in the uppermost portion of the core (core depth: 1–10 cm; Fig. 2).

### TOC and TN contents

The profile of TOC content showed three trends. First, TOC content was relatively constant (approximately 0.75 wt%) from the lowest portion of the core (core depth of 80 cm) to a depth of approximately 45 cm. Thereafter, TOC content gradually increased and reached approximately 0.85 wt% at a core depth of 12 cm. TOC content then increased rapidly, reaching more than 1.0 wt% in the upper part of the core. The TN profile displayed trends similar to TOC content. The C/N ratio ranged from 6.9 to 11.0. The C/N ratio was stable from the lowest portion of the core to a depth of 12 cm (approximately 7.5). It increased rapidly from that horizon and reached approximately 9.0 (Table 1, Fig. 2).

### Ostracoda

A list of the ostracod species is shown in Table 2. Fifty-three ostracod species were identified, and 48 of them comprised less than 5% of the total specimens in the samples. Fig 3 shows scanning electron microscopy photographs of the dominant species from the core.

Two species, *Keijella carriei* and *Loxoconcha wrighti*, were abundant throughout the core; 36% and 14%, respectively. These species were first reported from the Java Sea, west of Bawean Island, Indonesia (Dewi, 1997) and were abundantly distributed in the inner to middle parts of Jakarta Bay, Indonesia (Fauzielly et al., in press). In Jakarta Bay, *K. carriei* is abundant at a water depth of less than 20 m, in a sediment with a fine grain

**Table 1.** List of sample data

sample	grainsize ( $\phi$ )	mud content (%)	TN (wt%)	TOC(wt%)	C/N ratio	Diversity (H)	Evenness (H)
TJ56-1	8.4	75.7	0.13	0.99	7.70	1.6	0.5
TJ56-2	8.4	91.7	0.11	1.04	9.39	2.1	0.7
TJ56-3	8.6	94	0.11	0.97	8.84	1.7	0.4
TJ56-4	7.7	96.7	0.10	0.91	8.78	1.9	0.4
TJ56-5	8.6	99.8	0.12	0.94	7.75	1.8	0.4
TJ56-6	8.8	96.2	0.09	1.03	10.96	2.0	0.5
TJ56-7	7.4	96.1	0.13	0.99	7.84	2.0	0.5
TJ56-8	8.7	90	0.12	1.01	8.64	2.0	0.4
TJ56-9	8.7	97.3	0.11	1.08	9.39	2.1	0.4

TJ56-10	7.6	96.5	0.12	1.03	8.50	2.0	0.5
TJ56-11	8.3	97	0.12	0.95	7.92	2.2	0.5
TJ56-12	7.9	94.3	0.12	0.86	7.17	1.9	0.4
TJ56-13	8.2	98.3	0.12	0.84	7.00	2.2	0.5
TJ56-14	8.8	94.3	0.12	0.89	7.42	2.1	0.5
TJ56-15	7.9	98.3	0.13	0.9	6.92	2.0	0.6
TJ56-16	8.3	98.5	0.12	0.86	7.17	2.1	0.4
TJ56-17	8.7	97	0.12	0.87	7.25	2.1	0.4
TJ56-18	8.7	94.4	0.12	0.87	7.25	2.2	0.6
TJ56-19	8.6	97.3	0.12	0.92	7.67	1.7	0.4
TJ56-20	8.8	92	0.11	0.88	8.00	2.1	0.5
TJ56-21	8.6	96.5	0.12	0.86	7.17	2.2	0.6
TJ56-22	8	95.5	0.12	0.87	7.25	2.3	0.6
TJ56-23	8.3	94.8	0.12	0.85	7.08	2.3	0.6
TJ56-24	8.5	96.6	0.12	0.84	7.00	2.3	0.6
TJ56-25	8.1	96.6	0.12	0.91	7.58	2.3	0.6
TJ56-26	8	95.9	0.11	0.85	7.73	2.1	0.6
TJ56-27	7.9	93.9	0.12	0.87	7.25	2.1	0.4
TJ56-28	7.9	95	0.12	0.86	7.17	2.1	0.5
TJ56-29	8	95.2	0.11	0.83	7.55	2.1	0.5
TJ56-30	8.7	95.8	0.12	0.87	7.25	2.4	0.7
TJ56-31	8.2	94.3	0.11	0.84	7.64	2.1	0.5
TJ56-32	8.3	97.1	0.11	0.85	7.73	2.0	0.5
TJ56-33	8.3	96.8	0.11	0.84	7.64	2.0	0.4
TJ56-34	8.8	98.3	0.11	0.83	7.55	1.8	0.5
TJ56-35	8	97.1	0.11	0.85	7.73	1.9	0.5
TJ56-36	8.6	95.3	0.12	0.86	7.17	1.9	0.5
TJ56-37	7.9	97.1	0.1	0.8	8.00	2.2	0.5
TJ56-38	8.1	95.2	0.11	0.85	7.73	2.1	0.6

TJ56-39	7.9	95.7	0.11	0.82	7.45	2.2	0.5
TJ56-40	8.6	95.1	0.11	0.79	7.18	2.2	0.4
TJ56-41	8.5	84.6	0.09	0.74	8.22	2.0	0.4
TJ56-42	8.7	92.7	0.11	0.82	7.45	2.2	0.5
TJ56-43	8.7	86.6	0.11	0.78	7.09	2.3	0.5
TJ56-44	8.1	86.5	0.1	0.76	7.60	2.5	0.6
TJ56-45	7.9	93.1	0.09	0.73	8.11	2.0	0.5
TJ56-46	8.5	94.4	0.1	0.74	7.40	2.0	0.4
TJ56-47	8.1	94.3	0.1	0.74	7.40	2.3	0.5
TJ56-48	8	84.7	0.1	0.74	7.40	2.3	0.6
TJ56-49	8.5	79	0.09	0.7	7.78	2.1	0.5
TJ56-50	8.5	89.7	0.1	0.74	7.40	1.9	0.5
TJ56-51	8.7	93.6	0.11	0.79	7.18	2.2	0.6
TJ56-52	8.6	88.9	0.1	0.76	7.60	2.3	0.6
TJ56-53	8	93.6	0.1	0.7	7.00	2.3	0.5
TJ56-54	7.9	94.9	0.1	0.77	7.70	2.3	0.5
TJ56-55	7.8	81.1	0.1	0.76	7.60	2.1	0.5
TJ56-56	7.6	47	0.09	0.71	7.89	2.1	0.5
TJ56-57	8.2	90.2	0.09	0.76	8.44	2.0	0.5
TJ56-58	7.8	92.1	0.09	0.74	8.22	2.2	0.5
TJ56-59	8.2	90.9	0.09	0.78	8.67	2.2	0.5
TJ56-60	8.3	97.1	0.1	0.76	7.60	2.0	0.6
TJ56-61	8.7	79	0.09	0.78	8.67	2.2	0.6
TJ56-62	8.5	94.9	0.1	0.74	7.40	2.4	0.5
TJ56-63	8.5	93.3	0.09	0.69	7.67	2.2	0.5
TJ56-64	8.3	61.4	0.09	0.67	7.44	1.9	0.4
TJ56-65	8.4	82.9	0.1	0.73	7.30	2.1	0.5
TJ56-66	8.2	87.4	0.09	0.8	8.89	2.4	0.5
TJ56-67	8.5	94.7	0.1	0.74	7.40	2.3	0.6

TJ56-68	8.2	94.3	0.09	0.7	7.78	1.9	0.4
TJ56-69	8.3	87.7	0.09	0.69	7.67	2.0	0.4
TJ56-70	8.3	87.5	0.09	0.69	7.67	2.3	0.4
TJ56-71	8.5	83.4	0.1	0.72	7.20	2.2	0.6
TJ56-72	8.5	85.9	0.1	0.75	7.50	1.8	0.4
TJ56-73	8.3	87	0.09	0.66	7.33	2.0	0.5
TJ56-74	8.2	82.3	0.09	0.72	8.00	2.2	0.5
TJ56-75	7.8	88.3	0.11	0.82	7.45	2.2	0.4
TJ56-76	8.1	80.6	0.09	0.69	7.67	2.5	0.6
TJ56-77	8	84.8	0.09	0.72	8.00	2.0	0.4
TJ56-78	8.3	68.4	0.09	0.68	7.56	1.9	0.4
TJ56-79	8.5	65.8	0.08	0.71	8.88	1.9	0.4
TJ56-80	8.1	67.7	0.09	0.72	8.00	1.3	0.4

size (silt–clay), with a TOC content of 0.5–1.5 wt% (Fauzielly et al., in press), whereas to the west of the Bawean Island, this species has been found in gravelly mud at a water depth of 25 m (Dewi, 1997). *L. wrighti* is abundant at water depths of 10–30 m in Jakarta Bay (Fauzielly et al., in press) whereas in the Java Sea, it is found in gravelly mud bottoms at a water depth of 63 m (Dewi, 1997). *K. carriei* and *L. wrighti* are common in areas with high TOC and TN contents, even when they are anoxic (Fauzielly et al., in press).

The species *Pistocythereis* cf. *bradyformis*, *Neomonoceratina delicata*, *Phlyctenophora orientalis*, *Neomonoceratina iniqua*, Loxoconchidae gen. sp. indet., *Hemicytheridea reticulata* and *Propontocypris* sp. 1, were common throughout the core (3%–10%). Loxoconchidae gen. sp. indet. is the same

species as Dewi's (1997) new genus and new species (*Baweanconcha indonesiana*), but it was never formally described. Most of the species reported are commonly found living in tropical shallow seas in southeastern Asia at water depths of 20–50 m (Whatley and Zhao, 1987, 1988; Zhao and Whatley, 1989; Mostafawi, 1992; Dewi, 1997, Fauzielly et al., in press). *P. orientalis* was rarely found in the lower half of the core, whereas *H. reticulata* and Loxoconchidae gen. et sp. indet. were common in the lower half of the core (Figure 4).


To clarify the structure of the ostracod assemblages, the Shannon–Wiener function was used as an index of species diversity [H(S)]. Equitability (E) was calculated using the function of Buzas and Gibson (1969). The diversity index ranged from 1.3 to 2.53





Species	TJ56-31	TJ56-32	TJ56-33	TJ56-34	TJ56-35	TJ56-36	TJ56-37	TJ56-38	TJ56-39	TJ56-40	TJ56-41	TJ56-42	TJ56-43	TJ56-44	TJ56-45	TJ56-46	TJ56-47	TJ56-48	TJ56-49	TJ56-50	TJ56-51	TJ56-52	TJ56-53	TJ56-54	TJ56-55	TJ56-56	TJ56-57	TJ56-58	TJ56-59	TJ56-60	
<i>Alocopocythere kendengensis</i>																							1								
<i>Atjehella kingmai</i>										1																					
<i>Argillocea sp</i>	1	1	2						1	2	1	2	4	17	7	1		1					3	6	1	1			1	3	2
<i>Baweanconcha indonesiana</i>	11	4	9	2	2	1	5	5	10	4	10	20	7	7	7	18	18	25	16	17	15	12	17	10	7	14	8	14	10	4	
<i>Copypus posterosulcus</i>	6	2	4	4	1	3	1	4	4	6	1	3	6	7	3	2	9	9	4	5	12	5	4	3	3	5	2	2	4		
<i>Cytherella hemipuncta</i>	1	1			4	2			2	1	1	3	1	5				2		4	5	4	1	2	2	2		1	3		
<i>Cytherella incohota</i>				1			1									2			1										1		
<i>Cytherella javaseanse</i>										1	1													2						1	
<i>Cytherella semitalis</i>																															
<i>Cytherella koegleri</i>																															
<i>Cytherelloidea singulata</i>																															
<i>Cytherelloida cf excavata</i>																															
<i>Cytherelloida leroi</i>																															
<i>Hemicytheridea reticulata</i>	12	5	5	1	1	5	3	4	9	3	4	8	10	15	5	6	12	6	11	11	12	11	9	13	6	13	8	16	7		
<i>Hemicytheridea ornata</i>	4								1	1	1	3		3			3	1	4	3		2	7	5	4	4	6	2	6		
<i>Hemikrite petersoni</i>																		1			1										
<i>Hemikrite orientalis</i>																															
<i>Keijella carriei</i>	66	61	88	66	26	56	51	77	69	74	61	84	62	42	72	78	55	43	75	80	61	46	63	63	51	57	59	62	62	32	
<i>Keijella kloempritisensis</i>																															
<i>Keijella reticulata</i>																								2							
<i>Keijia labyrinthica</i>																															
<i>Loxoconcha wrighti</i>	40	15	26	17	3	25	20	25	28	27	22	27	36	22	34	36	31	27	47	43	30	23	37	18	28	28	27	33	32	10	
<i>Malayacythereis trachodes</i>																															
<i>Neocytheretta adunca</i>						1						2		1		4	2						1								
<i>Neocytheretta sp</i>	1								1		1																				
<i>Neocytheretta cf angulasa</i>							1																								
<i>Neocytheretta spongiosa</i>																1							1								
<i>Neocytheretta cf spongiosa</i>							2				3																			2	
<i>Neocytheretta murilineata</i>		2	1						2				1			2			4	2	3				1	1			5		
<i>Neomonoceratina delicata</i>	27	12	21	11		9	19	8		16	11	19	12	17	11	32	20	25	17		25	27	13		7	7	27		2		
<i>Neomonoceratina iniqua</i>		4	5	8	6	9	4	4	23	4	3	7	4	4	10		4	7	5	23		4	15	21	4	4	3	12	31	4	
<i>Neomonoceratina rhomboidea</i>	3	3	2	4	3		4	3	2	3	4	5	3	9	3	2	6	1	3		5	4	4	5	1	1	2	4	4	1	
<i>Neomonoceratina columbiformis</i>																															
<i>Parakrithella sp</i>									1																						
<i>Parakrite placida</i>																															
<i>Paratycerois sp</i>	2					1	3	8	3	2			2	1		1	1	2	3	2		1				1		2	1		
<i>Paracypris nuda</i>												2																			
<i>Phlyctenophora orientalis</i>	4	17	10	3	4	6	15	17	17	21	24	13	16	1	10	5	4	1	5	2	1	3	2	10	5	3	3	5	5	7	
<i>Phystocythereis sp</i>										15														7					20		
<i>Propontocypris sp 1</i>	41	5	3	6	1	1	3	12	4	4	4	3	4	6	7	9	5	2	3	3	6	11	1	5	3	1		3	4		
<i>propontocypris sp2</i>			5				2	12	6	5	1	3	5	6	2		1	12				3	6					1	1	5	
<i>propontocypris cf subangularis</i>																															
<i>Phystocythereis cf bradyformis</i>	11	12	19	13	2	18	17	21	12	1		10	13	17	16	13	22	23	18	19	20	22	17	3	27	13	30	33	1	7	
<i>Stigmatocythere cf indica</i>																															
<i>Stigmatocythere roesmani</i>																															
<i>Stigmatocythere indica</i>																													1		1
<i>Stigmatocythere sp</i>																															
<i>Tanella gracillis</i>				1	2	2	1						1			2	4				1	1	3		2						
<i>Venericythere sp</i>																						2			5						
<i>Venericythere gonia</i>							1			2	1																		4	1	
<i>Venericythere papuensis</i>	2		1						1		4		7	1		1	5	5	2		2	1	1	1	1	1	4	9	5	1	
Total number specimen	232	144	202	136	55	139	153	201	197	195	156	216	206	172	181	215	202	196	221	213	202	177	210	176	151	158	186	207	195	78	
Total number species	16	14	16	12	12	14	18	14	19	21	18	17	17	19	13	17	20	16	17	13	16	17	20	18	16	16	14	19	19	12	
sample weight	3.18	5.83	3.18	4.40	4.32	3.18	4.10	5.99	5.62	4.28	4.73	5.07	5.57	5.84	5.38	3.91	3.22	4.25	3.20	3.32	5.16	3.86	5.81	5.05	3.67	3.21	3.50	7.06	3.68	4.59	
split	4	1	1	2	2	1	1	2	2	1	4	2	2	2	1	2	2	4	4	2	4	2	2	2	4	2	1	1	1	2	1
split sample weight	0.80	5.83	3.18	2.20	2.16	3.18	4.10	2.99	2.81	4.28	1.18	2.53	2.78	2.92	5.38	1.96	1.61	1.06	0.80	1.66	1.29	1.93	2.90	1.26	1.84	3.21	3.50	7.06	1.84	4.59	
no of individu	292	25	64	62	25	44	37	67	70	46	131	85	74	59	34	110	125	184	276	128	157	92	72	139	82	49	53	29	106	17	

Species	TJ56-61	TJ56-62	TJ56-63	TJ56-64	TJ56-65	TJ56-66	TJ56-67	TJ56-68	TJ56-69	TJ56-70	TJ56-71	TJ56-72	TJ56-73	TJ56-74	TJ56-75	TJ56-76	TJ56-77	TJ56-78	TJ56-79	TJ56-80
<i>Alocopocythere kendengensis</i>										1				3	1		1			
<i>Atjehella kingmai</i>										1										
<i>Argilocea sp</i>		1	3	1			2		1	1				1	2					
<i>Baweanconcha indonesiana</i>	15	15	23	16	12	15	12	9	13	9	20	6	7	3	5	4	4	3	3	
<i>Copytus posterosulcus</i>	2	6	4	1	5	5	8	1			4				3	3	1	5	2	
<i>Cytherella hemipuncta</i>	2			1	1		5	1	1	1		1	1		3	5			1	
<i>Cytherella incohota</i>																				
<i>Cytherella javaseanse</i>		4	1	4	1	7		2						2						
<i>Cytherella semitalis</i>																				
<i>Cytherella koegleri</i>																				1
<i>Cytherelloidea singulata</i>																				
<i>Cytherelloida cf excavata</i>																	1			
<i>Cytherelloida leroi</i>																3				
<i>Hemicytheridea reticulata</i>	12	17	17	5	7	8	14	20	22	24	7	11	1	2	5	10	4	9	6	3
<i>Hemicytheridea ornata</i>	9	8	1	2	3	4	4	1	4	2	2	2	5	5	3	2	2		1	
<i>Hemikrite petersoni</i>																				
<i>Hemikrite orientalis</i>								1						2		4		1		
<i>Keijella carriei</i>	41	54	52	64	80	68	60	78	82	78	66	73	71	58	65	30	98	66	65	48
<i>Keijella kloempritisensis</i>												1					1			
<i>Keijella reticulata</i>					3								5							
<i>Keijia labyrinthica</i>																				
<i>Loxococoncha wrighti</i>	45	43	43	44	28	40	30	28	39	35	38	10	17	16	32	22	22	20	16	10
<i>Malaycythereis trachodes</i>																1				
<i>Neocytheretta adunca</i>										2	1									
<i>Neocytheretta sp</i>																				
<i>Neocytheretta cf angulasa</i>																				
<i>Neocytheretta spongiosa</i>				2					2						2					
<i>Neocytheretta cf spongiosa</i>																		2		
<i>Neocytheretta murilineata</i>		4	2			1	5	1	1	5			2	2	2	1				
<i>Neomonoceratina delicata</i>	14	17	28	23	13	30	10		13	12	14	9	12		19	15	3	1	7	
<i>Neomonoceratina iniqua</i>	8	5		6	3	6	9	15	12	6	8	4	7		4		14	11	5	13
<i>Neomonoceratina rhomboidea</i>	3	3	2	4		3	4	4		8	8		6	16	6	2	1		1	
<i>Neomonoceratina columbiformis</i>										1										
<i>Parakrihella sp</i>		1				1														
<i>Parakrite placida</i>							1													
<i>Paratyerois sp</i>				1		5	1							1	3	1		1	2	1
<i>Paracypris nuda</i>																				
<i>Phlyctenophora orientalis</i>	3	5	3	3	6	10	3	1	5	5	5	8	5	8	2	1	7	6	6	5
<i>Phystocythereis sp</i>			13	7	10	13		11	1			6	16	6			9		3	6
<i>Propontocypris sp 1</i>	2	4	5	1	1		4	4	4	2	12				4	2	4	3		
<i>propontocypris sp2</i>	4	1				8				1										
<i>propontocypris cf subangularis</i>																				
<i>Phystocythereis cf bradyformis</i>	18	1			2	3	17	1	4	17	11	3		3	29	2	5			
<i>Stigmatocythere cf indica</i>		1	1						1	4	3		3	5						
<i>Stigmatocythere roesmani</i>										1						2			2	
<i>Stigmatocythere indica</i>										1		1	3		2	2	4	3	12	
<i>Stigmatocythere sp</i>																		3		2
<i>Tanella gracillis</i>			2		3			1	1	3	1			1	1			1		
<i>Venericythere sp</i>		8		2	4					4						5	9			
<i>Venericythere gonia</i>		4	3			4		1	4	2		1	2	8		4	14	10	2	1
<i>Venericythere papuensis</i>	1	1	6	3	8	10	2	2	2	2	3	2	3	1	1	4				2
Total number specimen	179	203	209	190	190	241	191	182	212	228	203	138	166	143	194	125	204	145	136	90
Total number species	15	21	18	19	18	19	18	19	19	26	16	15	17	19	21	22	19	16	17	10
sample weight	4.32	4.82	4.58	4.74	5.69	4.51	4.75	4.51	6.65	6.30	5.50	4.78	5.30	5.83	5.48	2.00	2.94	2.00	1.40	5.01
split	4	2	1	2	2	2	2	2	4	1	2	4	1	4	2	4	1	4	4	2
split sample weight	1.08	2.41	4.58	2.37	2.85	2.26	2.38	2.26	1.66	6.30	2.75	1.20	5.30	1.46	2.74	0.50	2.94	0.50	0.35	2.51
no of individu	166	84	46	79	67	107	80	81	128	36	74	116	31	98	71	250	69	290	389	36



**Fig. 3.** Scanning electron micrographs of the dominant species. (ALV: adult left valve, ARV: adult right valve, JRV: juvenile right valve) 1, *Keijella carriei* Dewi, ALV, sample TJ 56-8; 2, *Loxoconcha wrighti* Dewi, ALV, sample TJ56-21; 3, *Pistocythereis* cf. *bradyformis* (Ishizaki), ARV, sample TJ56-21; 4 *Neomonoceratina delicata* Ishizaki and Kato, ARV, sample TJ 56-40; 5, *Phlyctenophora orientalis* (Brady),ALV,sample TJ56-25; 6, *Neomonoceratina iniqua* Brady, ALV, sample TJ56-28; 7, Loxoconchidae gen. et sp. Indet, ALV, sample TJ56-6; 8 *Hemicytheridea reticulata* Kingma, ALV, sample TJ56-6; 9, *Propontocypris* sp 2, ARV, sample TJ56- 8; 10, *Copytus posterosulcus* Wang, ARV, sample TJ 56-10; 11, *Phystocythereis* sp, ARV, sample TJ56-74; 12 *Propontocypris* sp 1, ALV, sample TJ56-39. 13, *Neomonoceratina rhomboidea* Hanai, ALV, sample TJ56-37; 14. *Argilocea* sp, ARV, sample TJ56-9; 15. *Cytherella hemipuncta*, Swanson, ARV, sample TJ56-24; 16. *Hemicytheridea ornata* Mostafawi, ALV, sample TJ56-28; 17. *Venericythere papuensis* (Brady),ALV, sample TJ56-14; 18. *Paratycerois* sp, JRV, sample TJ56-4.  
Scale bars = 0.1 mm

(low to moderate) (Table 2), and was relatively constant throughout the core but decreased slightly in the uppermost portion. Equitability ranged from 0.36 to 0.68 (moderate to high) (Table 2) and fluctuated throughout the core. Ostracod density (N: the number of specimens per 1 gram dry sediment sample) was approximately 50. However, the average density (N = 93) in the lower half of the core was higher than that in the upper half (N = 58). This trend paralleled the trend for number of species.

### Discussion

The chronology of the core was based on the  $^{210}\text{Pb}$  information from the Centre for Oceanological Research and Development of the Indonesian Institute of Sciences (Rositasari, pers. comm.), from a study site located approximately 6 km southeast of the present core. The sedimentation rate ranged from 0.4 to 2.0 cm/year, and the maximum sedimentation

rate was in the 50–60 cm layer. Thus, the age of this core ranges from prior to 1920 AD (depth > 60cm) to 1994 AD.

The stable clay sedimentation throughout the core and its location in a semi-enclosed bay indicate that sediments were accumulated at the study site in a low-energy and stable environment. The history of the depositional environment in this core can be divided into two major intervals (0–45 cm = approximately 1994–1950 AD); 45–80 cm = approximately 1950–before 1918 AD) based on TOC and TN contents. The upper interval is characterized by higher TOC (0.79–0.95 wt%) and TN (0.10–0.13 wt%) contents and dark greyish brown clay containing sporadic shell fragments. The lower interval is characterized by low TOC (0.66–0.82%) and TN (0.09–0.11%) contents and massive olive grey clay containing large fragments of molluscs ( Fig. 4)




Fig. 4. Temporal change of dominant species

The increase in TOC and TN contents can be explained by the increase in population of Jakarta City. From 1950 to 1990, the population rapidly increased by almost fourfold, from 1,733,600 to 7,515,000. Waste from domestic areas and industries that produce nutrients probably caused the increase in TOC and TN contents. Although the population increased rapidly, TOC content gradually increased, suggesting a high rate of sedimentation and dilution by natural forces.

The gradual increase in TOC content since around 1950 did not greatly impact the structure of ostracod assemblages; the diversity index showed stable values throughout the core, but species number and density decreased slightly in the upper half of the core. Ostracod assemblages were composed of high percentages of *K. carriei*, *L. wrightii*, *N. delicata* and *N. iniqua* throughout the core at relatively stable frequencies, which were not affected by the small increase in TOC content.

The percentages of *Loxoconchidae* gen. sp. indet. and *H. reticulata* decreased at a core depth of 45 cm (around 1950), where TOC content began to increase. Fauzielly et al. (in press) reported recent ostracods from core top samples collected in 1994 in Jakarta Bay and analysed the relationships between the density of dominant species in Jakarta Bay and TOC and TN contents. According to this study, *H. reticulata* is dominant in the middle part of Jakarta Bay and decreases abruptly at more than 0.8 and 0.14 wt% in TOC and TN, respectively, and prefers lower TOC and TN contents. Thus, it is considered that the increase in TOC content since around 1950 was a direct cause of the decrease in the relative abundance of *H. reticulata*. On the other hand, the relative abundance of *P. orientalis* increased rapidly between core depths of 45 cm and 40 cm, and from 40 cm upward, it was relatively more common (4%–16%) than in the lower portion of the core (1.7%–10%). *P. orientalis* was common in the inner part of Jakarta Bay in 1994 (Fauzielly et al., in press). TOC content indicates pollution and is a source of food for ostracods. A small increase in TOC content allows *P. orientalis* to thrive, and thus, *P. orientalis* may be used as an indicator of an increase in TOC content within a narrow range.

## CONCLUSIONS

1. A total of 53 ostracod species were recognized in a core from the inner part of Jakarta Bay. The dominant species were *Keijella carriei* and *Loxoconcha wrighti*.
2. Abundance of *K. carriei* and *L. wrighti* throughout the core indicates that the benthic portion of Jakarta Bay was an organic-rich environment.
3. The rapid population increase in Jakarta City over the last 80 years caused an increase in the TOC content of muddy sediment. However, the increase in the TOC content is limited to between 0.7% and approximately 1.1 %, probably due to the high mud sedimentation rate.
4. In approximately the 1950s, *Loxoconchidae* gen. sp. indet. and *L. wrighti* decreased and *Phlyctenophora orientalis* increased due to organic contamination related to the increased population of Jakarta City. Thus, these markers may become useful tools for

monitoring of coastal areas displaying the effects of organic pollution.

## ACKNOWLEDGMENTS

We gratefully acknowledge the Marine Institute of Geology, Indonesia for providing core samples and The Directorate General of Higher Education, Ministry of Education, Indonesia for financial assistance to carry out this study.

## REFERENCES

- Brady, G. S., 1867-1872, In L.DeFolin and L.Perier (eds) Les Fonds de la Mer,V,1 pt,1,XIV Quelques points de la cote septentrionale de Java, p.59-70, pl,7,8
- Brady, G. S., 1880, Report on the Ostracoda dredged by HMS Challenger during the year 1873-1876, Zoology, p. 1–184.
- Buzas, M. A. and Gibson, T. G., 1969, Species diversity: benthic foraminifera in western North Atlantic. Science, 163, 72–75.
- Carbonel, P. and Moyes, J., 1987, Late Quaternary paleoenvironments of the Mahakam Delta (Kalimantan,Indonesia). Palaeogeography, Palaeoclimatology, Palaeoecology, 61, 265-284.
- Dewi, K. T., 1997, Ostracoda from the Java Sea, west of Bawean Island Indonesia, Marine Geological Institute, Special Publication no. 4, 1997, Ministry of Mines and Energy, Directorate General Of Geology and Mineral Resources, Marine Geological Institute, Bandung, 86p.
- Hanai, T., Ikeya, N. and Yajima, M., 1980, Checklist of ostracoda from Southeast Asia. University of Tokyo Press Bulletin 17, 236p.
- Horne, D.J., Cohen, A., Martens, K., 2002,Taxonomy, morphology and biology of Quaternary and living Ostracoda. In Holmes, J. A. and Chivas,

- A. R. (eds). *The Ostracoda: Applications in Quaternary Research*. American Geophysical Union, Washington, D. C, p 5–36.
- Irizuki, T., Takimoto, A., Sako, M., Nomura, R., Kakuno, K., Wanishi, A. and Kawano, S., 2011, The influences of various anthropogenic sources of deterioration on meiobenthos (Ostracoda) over the last 100 years in Suo-Nada in the Seto Inland Sea, southwest Japan. *Marine Pollution Bulletin*, 62, 2030-2041.
- Kingma, J. T., 1948, Contributions to the knowledge of the young Cenozoic Ostracoda from the Malayan region, PhD thesis, University of Utrecht. 119p.
- Keij, A. J., 1953, Preliminary note on the recent Ostracoda of the Snellius Expedition. *Proceeding of the Koninklijke Nederlandse Akademie van Wetenschappen*, Series 56, 155–168.
- Keij, A. J., 1964, Neogen to Recent species of Chutherelloidea (Ostracoda) from Northwestern Borneo. *Micropaleontology*, 10, 415–430.
- Keij, A. J., 1974, Review of the Indopacific species of Tribelina (Ostracoda). *Proceeding of the Koninklijke Nederlandse Akademie van Wetenschappen*, Series B77, 345–358
- Keij, A. J., 1975, Note on three Holocene Indo-Malaysian ostracod species. *Proceeding of the Koninklijke Nederlandse Akademie van Wetenschappen*, Series B78, 231–241.
- Mostafawi, N., 1992, Rezente Ostacoden aus Dem Mittleren Sunda-Schelf Zwischen der Malaiischen Halbinsel und Borneo. *Senckenbergiana Lethaea*, 72, 129–168.
- Pemerintah Provinsi DKI Jakarta, 2011, Status lingkungan hidup daerah propinsi DKI Jakarta tahun 2010, Jakarta, 385p.
- UNESCO, 2000, Reducing megacity impacts on the coastal environment – Alternative livelihoods and waste management in Jakarta and the Seribu Islands. *Coastal Region and Small Island Papers 6*, UNESCO, Paris, 59 pp.
- Whatley, R. C. and Watson, K., 1988, A preliminary Account of the Distribution of Ostracoda in Recent Reef and Reef Associated Environments in the Pulau Seribu or Thousand Island Group, Java Sea, In Hanai, T, Ikeya N. and Ishizaki, K.,(eds). *Evolutionary Biology on Ostracoda: Proceeding of the Ninth International Symposium on Ostracoda*, Shizuoka, 399–411
- Whatley, R. and Zhao, Q., 1987, Recent Ostracoda of the Malacca Straits Part I. *Revista Espanola de Micropaleontologia*, 19, 327–366.
- Whatley, R. and Zhao, Q., 1988, Recent Ostracoda of the Malacca Straits Part II. *Revista Espanola de Micropaleontologia*, 20, 5–37.
- Yasuhara M. and Yamazaki, H., 2005, The impact of 150 years of anthropogenic pollution on the shallow marine ostracode fauna, Osaka bay, Japan. *Marine Micropaleontology*, 55, 63–74.
- Yasuhara, M., Yamazaki, H., Irizuki, T. and Yoshikawa, S., 2003, Temporal changes of ostracode assemblages and anthropogenic pollution during the last 100 years, in sediment cores from Hiroshima Bay, Japan. *The Holocene*, 13, 527–536.
- Yasuhara, M., Yamazaki, H., Tsujimoto, A., Hirose, K., 2007. The effect of long-term spatiotemporal variations in urbanization-induced eutrophication on a benthic ecosystem, Osaka Bay, Japan. *Limnology and Oceanography* 52, 1633–1644.
- Zhao, Q. and Whatley, R., 1989, Recent podocopid Ostracoda of the Sedili River and Jason Bay, southeastern Malay Peninsula, *Micropaleontology*, 35, 168-187.