

# The Diversity of Molluscs in Seagrass Ecosystem of Wondama Bay, West Papua, Indonesia

Ludi Parwadani Aji<sup>1,2,\*</sup>, Andriani Widyastuti<sup>3</sup>, Hadiyanto Hadiyanto<sup>1</sup>

<sup>1</sup>Research Center for Oceanography, National Research and Innovation Agency  
Jl. Pasir Putih, Pademangan, Ancol, North Jakarta, DKI Jakarta 14430, Indonesia

<sup>2</sup>Marine Evolution and Ecology Group, Naturalis Biodiversity Center  
P.O. Box 9517, 2300 RA Leiden, Netherlands

<sup>3</sup>Fishery Academy of Kamasan  
Jl. Raya Bosnik No.1, Sumberker, Biak, Biak Numfor Regency Papua, Indonesia.  
Email: ludi001@brin.go.id

## Abstract

Papua, Indonesia is located in the Coral Triangle and is home to many marine creatures. Some areas of Papua, however, remain underexplored, particularly in terms of the diversity of mollusc and their seagrass bed habitat. This research aims to document benthic mollusc and seagrass species in Wondama Bay Regency and assess how habitat, temperature, and geographic location affect mollusc diversity and composition. Field data collection was conducted at nine sampling stations. A total of 197 benthic mollusc species was recorded, consisting of 53 species from 19 families of bivalves and 144 species from 40 families of gastropods. This suggests that mollusc species in Wondama Bay Regency still need further exploration which could reveal additional species. The bivalve *Tellina* sp. and gastropod *Mitrella scripta* were found in all nine sampling areas. Furthermore, nine species of seagrasses were recorded, with *Enhalus acoroides* and *Cymodocea rotundata* found at all sites. The number of seagrass species in our study is considerably high. The seagrass species richness ranged from three to eight species, while the percent cover varied from 13 % to 65 % across sites. This research found that temperature, seagrass species, and geographic location did not significantly impact the mollusc species composition among sites. Whereas, seagrass coverage showed as an important driver of mollusc community assemblages, highlighting the importance of seagrass bed protection for mollusk diversity. This finding may assist stakeholders in advancing marine conservation efforts and encouraging the sustainable utilization of marine resources particularly in Wondama Bay Regency, Papua.

**Keywords:** Bivalvia, Biodiversity, Gastropoda, Richness, Papua

## Introduction

Papua, located in the eastern part of Indonesia, at the center of the Coral Triangle region, is considered a pristine area of great interest for scientific exploration (Schultze-Westrum 2001; Palomares and Heymans 2006; Hoeksema 2007). Papua has diverse habitats, including coral reefs, seagrass beds, mangroves, rocky shores, soft sediments, and marine lake ecosystems (Mangubhai et al., 2012; Becking et al., 2024). The coastal areas of this region are home to many marine species (McKenna et al., 2002; Palomares and Heymans, 2006). One of the most diverse organisms in these tropical coastal regions is Mollusca, a group of soft-bodied animals (McKenna et al., 2002; Wells, 2002). Phylum Mollusca has various forms and sizes, with some living in shallow water areas, mainly benthic, meaning they live in or on the substrates (Dharma, 2005; Huber, 2010; Aji et al., 2018). Class Bivalvia and Gastropoda are the two most diverse and commonly found molluscs in coastal areas, especially

in seagrass ecosystems (Dharma, 2005; Huber, 2015; Aji and Widyastuti, 2020).

Seagrass is a flowering plant in shallow marine waters (Rahmawati et al., 2022). Seagrass beds, in particular, serve as important habitats for many marine organisms, including molluscs (Aji and Widyastuti, 2020; Paramasivam and Venkataraman, 2022). Seagrass beds provide services for marine animals, such as nursery areas, food, reproduction sites, and protection from predators (Rahmawati et al., 2022). Thus, seagrass availability is important for benthic mollusc communities (Rueda et al., 2009; Paramasivam and Venkataraman, 2022). Some molluscs are grazers or detritus feeders, primarily obtaining their food source from seagrass leaves (Rueda et al., 2009). Depending on the cover and species of seagrass, the mollusc species found in those seagrass beds may vary (Rueda et al., 2009; Aji and Widyastuti, 2020; Paramasivam and Venkataraman, 2022). Furthermore, geographic location and temperature can play a role in shaping

benthic biota community composition (Aji *et al.*, 2024). Seagrass and mollusc species composition differ between temperate, subtropical, and tropical areas due to temperature variations (Rueda *et al.*, 2008; Arbi, 2011; Islami *et al.*, 2018; Paramasivam and Venkataraman, 2022). Some mollusc species from class Bivalvia, such as those in the Mytilidae family, may survive extremely high temperatures (Huhn, 2016; Aji *et al.*, 2023). Hence, understanding the condition and community composition of molluscs and their seagrass habitats is essential for improving coastal conservation practices.

Seagrass in Indonesian waters consists of 16 species, with an overall cover average of around 42%, ranging from 5% to 81% (Rahmawati *et al.*, 2022). Specifically, based on the seagrass monitoring program held by the Indonesian government, in the Papua region there were reported eight species of seagrass in Raja Ampat, eight species in Cendrawasih Bay, eight species in Padaido islands, and six species in Biak island (Rahmawati *et al.*, 2022). Moreover, research on mollusc diversity in the Papua region is still limited compared to other parts of Indonesia (van der Meij *et al.*, 2009; Arbi, 2011; Eisenbarth *et al.*, 2018; Heryanto, 2018; Islami *et al.*, 2018), mainly due to the high costs and logistical challenges of conducting exploration in Papua. There have been few reports on the diversity and composition of molluscs in Papua's coastal ecosystem, such as in Raja Ampat (Wells, 2002) and Biak Island (Aji and Widyastuti, 2017, 2020). By exploring new geographic locations, researchers may discover mollusc species that are new to science or new records from specific areas. One of the underexplored regions for mollusc inventories is Wondama Bay Regency. Wondama Bay is part of Cendrawasih Gulf National Park in West Papua, Indonesia. While some studies have reported on the population and community structure of biota in Wondama Bay, such as whale shark (Suruan *et al.*, 2016), phytoplankton (Alianto *et al.*, 2018), reef fish (Madiyani *et al.*, 2018) and mangrove (Dharmawan and Widyastuti, 2017), there is limited information on mollusc and seagrass from this region.

The research aims to provide information on the diversity of molluscs in seagrass ecosystems of Wondama Bay. Specifically, our objective is to document the species list of molluscs and determine environmental factors, including habitat characteristics, temperature, and geographic location, that influence the diversity and composition of molluscs in seagrass ecosystems of Wondama Bay. This information may help stakeholders support marine conservation efforts and promote the sustainable use of marine resources.

## Materials and Methods


### *Sampling locations and characteristics*

Field data collection was conducted in Wondama Bay Regency at nine stations, with the distance between stations at least 3 km, to capture the spatial variation of Wondama Bay: Yende (Station 1), Kayob (Station 2), Nupu (Station 3), Sombokoro (Station 4), Batewar (Station 5), Anandoyari (Station 6), Imbursore (Station 7), Paraibo cape (Station 8), Ruruitua (Station 9) (Figure 1) during wet season from 9 to 19 February 2016. Geographically, Wondama Bay Regency (0° 15' - 3° 25' S and 132° 35' - 134° 45' E) borders Ransiki District of Manokwari Regency to the North, Yaur District of Nabire Regency to the East and South, and Kuri and Idor Districts of Bintuni Bay Regency to the West. Physical data, such as temperature and pH, were measured in situ using digital temperature and pH meters. Additionally, habitat characteristics, including substrate types (mud, muddy sand, sandy mud, rubble) and beach profiles of coastal areas on each station, were observed.

### *Seagrass and mollusc data collection*

Seagrass data was collected using the line and quadrat transect method, following the seagrass monitoring guidebook (Rahmawati *et al.*, 2017). Data collection occurred during low tide to facilitate more accessible sampling. A 100 m transect line was laid out perpendicular to the coastline, and square frames measuring 50 x 50 cm were placed at 10 m intervals from the shore. Seagrass species within the frames were identified, and their coverage percentage was recorded according to Rahmawati *et al.*, (2017).

For mollusc data collection, shore gleaning was conducted by walking around the seagrass beds and the beach, including mangrove and rocky shore habitats. Macro-molluscs (more than 1cm in size) were sampled qualitatively based on their occurrence (present/absence) at each site. Representative samples were collected for further identification in the laboratory and deposited as a reference collection at the Technical Implementation Unit for Marine Life Conservation – Indonesian Institute of Science, Biak, Papua. Morphological identification was performed using the guidebooks of Wilson *et al.*, (1993), Abbott and Dance (2000), Dharma (2005), and Huber (2010, 2015). The latest taxonomy and nomenclature provided by MolluscaBase (Molluscabase eds 2024) were adhered for classification.



**Figure 1.** Study area in (a) Wondama Bay Regency including nine sampling sites, (b) West Papua Province, and (c) Indonesia.

### Data analysis

Species accumulation curve with 999 permutations was conducted to evaluate whether most species was collected or not. After checking the normality assumptions, Pearson correlation ( $r$ ) tests were conducted to examine the relationships between mollusc richness and seagrass cover, mollusc richness and seagrass richness, mollusc richness and temperature, and seagrass richness and temperature using R v.4.0.2 (R Core Team 2023), with the results visualized through the ggplot2 package (Wickham, 2016). Moreover, the similarity in mollusc community composition between sites was evaluated using the presence/absence of species data for each site. The Jaccard index was calculated to measure this without applying transformations or standardization (Real and Vargas., 1996). This index was applied through pairwise comparisons between all sites. The similarity levels were determined by dividing the size of the intersection by the size of the union of the sample sets (Jaccard, 1908). Additionally, the composition of mollusc communities among sites was visualized using unconstrained

ordination non-metric multidimensional scaling (NMDS; Legendre and Gallagher, 2001) using the vegan package in R (Oksanen *et al.*, 2019). Finally, an envfit analysis with 999 permutations to obtain  $r^2$  and p-values (Oksanen *et al.*, 2019) was then performed to analyze the effect of explanatory variables (seagrass coverage, seagrass species, geographic location, and temperature) on the mollusc community assemblage across sites.

## Results and Discussion

### Characteristics and environmental condition

The habitat characteristics and environmental conditions of sites in the present study are presented in Table 1. Environmental parameter of temperature ranged between 26.8°C (St. 8) and 32°C (St.7). Ph ranged from approximately 7.36 (St. 9) to 8.1 (St. 4). According to Lee *et al* (2007), a temperature range between 23°C and 32°C supports maximum photosynthesis rates and optimal growth for tropical or subtropical seagrass species. Moreover, benthic molluscs in tropical regions can tolerate


temperatures as high as 36°C (Huhn, 2016; Aji *et al.*, 2023). Most substrate types in our sampling locations were sandy mud, mud, muddy sand, sand, and rubble. The substrate is commonly recognized as a characteristic for seagrass beds. Furthermore, the beach profiles of our study sites were mostly characterized by mangroves, coconut trees, rocky shores, villages, and docking spots. Most of the sites had mangroves, except at Station 1. According to Dharmawan and Widyastuti (2017), Wondama Bay had 18 species of true mangroves, with a relatively high canopy cover of 82±6 %, dominated by genus *Rhizophora*.

### Seagrass species diversity and coverage

This study recorded nine species of seagrasses (*Thalassia hemprichii*, *Enhalus acoroides*, *Cymodocea rotundata*, *Halodule uninervis*, *Halodule pinifolia*, *Halophila ovalis*, *Syringodium isoetifolium*, *Cymodocea serulata*, and *Halophila minor*) which belong into two families (Cymodoceaceae and Hydrocharitaceae) from the sampling locations (Figure 2, Table 1). The seagrass species *Enhalus acoroides* and *Cymodocea rotundata* were found at all nine sites. Rahmawati *et al* (2022) confirmed that those two seagrass species are widely distributed across Indonesian waters. However, *Halophila minor* was only found in station 5.

The highest number of seagrass species was found at Station 8, with eight species, while the lowest was found at Station 2, with only three species. From our understanding, a seagrass inventory in Wondama Bay has been reported by Tebay *et al.* (2020) at three sites in Aisandami village, where three seagrass species were found: *Thalassia hemprichii*, *Enhalus acoroides* and *Halophila ovalis*. The number of seagrass species in our study is considerably higher compared to other parts of Papua. Several reports on seagrass species from the Papua region found six species in Raja Ampat (Rizqi *et al.*, 2019), eight species in Padaido Islands (Aji *et al.*, 2018), eight species in Biak Island (Aji and Widyastuti, 2020), eight species in Salawati island, eight species in Batanta island (Rondonuwu *et al.*, 2019), and seven species in small islands of northern Papua (Nugraha *et al.*, 2021).

A higher diversity of seagrass species in a single location is believed to enhance the formation of a robust seagrass community, as each species plays a unique role. Some species are effective at trapping sediment and acting as barriers to currents, while others serve as substantial nursery grounds and food sources for many marine creatures as well as for megaherbivores such as dugongs and sea turtles (Rueda *et al.*, 2009; Nugraha *et al.*, 2021; Rahmawati *et al.*, 2022). Based on the characteristics of seagrass



**Figure 2.** Some seagrass species from Wondama Bay Regency, West Papua, Indonesia. Note: A. *Thalassia hemprichii*, B. *Halodule uninervis*, C. *Halophila ovalis*, D. *Halodule pinifolia*, E. *Syringodium isoetifolium*, F. *Cymodocea rotundata*, and G. *Cymodocea serulata*.

**Table 1.** Habitat characteristics and environmental parameters from nine sampling locations in Wondama Bay Regency. Note: Th= *Thalassia hemprichii*, Ea= *Enhalus acoroides*, Cr= *Cymodocea rotundata*, Hu= *Halodule uninervis*, Hp= *Halodule pinifolia*, Ho= *Halophila ovalis*, Si= *Syringodium isoetifolium*, Cs= *Cymodocea serulata*, Hm= *Halophila minor*.

| St | Name location          | Latitude   | Longitude   | Seagrass cover | Seagrass species               | Temp (°C) | pH   | Substrates         | Beach profiles           |
|----|------------------------|------------|-------------|----------------|--------------------------------|-----------|------|--------------------|--------------------------|
| 1  | Yende, Roon Island     | 02.38551 S | 134.53829 E | 38.92          | Th, Ea, Cr, Hu, Hp, Ho, Si     | 29.7      | 7.69 | sandy mud, mud     | coconut trees            |
| 2  | Kayob, Roon Island     | 02.38522 S | 134.51095 E | 12.86          | Ea, Cr, Hp                     | 30.2      | 7.71 | muddy sand         | mangroves, small village |
| 3  | Nupu, Windesi          | 02.43895 S | 134.25823 E | 45.56          | Th, Ea, Cr, Hu, Ho             | 30.0      | 7.6  | sandy mud          | mangroves, coconut trees |
| 4  | Sombokoro, Windesi     | 02.51179 S | 134.39609 E | 42.67          | Th, Ea, Cr, Cs, Hp, Ho, Si     | 31.0      | 8.1  | sandy mud          | mangroves                |
| 5  | Batewar, Windesi       | 02.55682 S | 134.33682 E | 35.38          | Th, Ea, Cr, Hu, Hp, Ho, Hm, Si | 28.0      | 7.63 | muddy sand, rubble | mangroves, nearby dock   |
| 6  | Anandoyari, Yop Island | 02.54339 S | 134.39269 E | 46.15          | Th, Ea, Cr, Hu, Hp, Ho, Si     | 31.2      | 7.97 | sandy mud, rubble  | coconut trees, mangroves |
| 7  | Imbursore, Yop Island  | 02.52088 S | 134.37366 E | 65             | Th, Ea, Cr, Hu, Hp, Ho, Si     | 32.0      | 7.75 | sand, rubble       | mangroves, rocky shores  |
| 8  | Paraibo, Yop Island    | 02.57487 S | 134.38280 E | 55.71          | Th, Ea, Cr, Cs, Hu, Hp         | 26.8      | 7.64 | sandy mud, rubble  | mangroves                |
| 9  | Ruruitua, Wasior       | 02.59666 S | 134.46537 E | 35.42          | Ea, Cr, Hp, Ho                 | 28.8      | 7.36 | sandy mud, rubble  | mangroves                |

habitats, life cycles, and meadow structures proposed by Kilminster *et al.* (2015). Small-sized seagrass species belong to opportunistic and colonizing groups—such as *Halophila ovalis*, *Cymodocea rotundata*, *Halodule uninervis*, *Halodule pinifolia*, and *Syringodium isoetifolium*—generally have rapid growth rates and short lifespans (Kilminster *et al.*, 2015; Rahmawati *et al.*, 2022). They may easily be uprooted from the substrate if conditions or disturbances become too severe. In contrast, larger seagrass species, such as *Enhalus acoroides* and *Thalassia hemprichii*, are considered persistent groups (Kilminster *et al.*, 2015). They tend to grow slowly but have strong adaptations to withstand pressure or extreme conditions. However, they require more time to recover than opportunistic and colonizing groups (Kilminster *et al.*, 2015; Nugraha *et al.*, 2021). Hence, a more diverse seagrass species in the ecosystem lead to more resilient ecosystem to disturbances.

In the present study, the average seagrass percent cover was 42%, with the highest seagrass cover recorded at Station 7 with 65%, whereas the lowest was at Station 2 with around 13%. For comparison, the percent cover of seagrass in other parts of Papua was reported to be 32% in Raja Ampat (Rizqi *et al.*, 2019), 42% in Padaido islands (Aji *et al.*, 2018), 36% in Biak island (Aji and Widyastuti, 2020), 32% in Salawati island, 28% in Batanta island (Rondonuwu *et al.*, 2019), and 59% in small islands of northern Papua (Nugraha *et al.*, 2021). According

to the Ministry of Environment Regulation no 200, the year 2004, seagrass cover in Wondama Bay (42%) is classified as being in fair condition, falling within the range of 30 to 59%. While, according to Rahmawati *et al.* (2014), it can be classified as medium (26-50%). It has been reported that seagrass cover in Indonesian waters declined from 42% in 2018 to around 34% in 2021 (Rahmawati *et al.*, 2022), likely due to anthropogenic pressures such as coastal development, human activities, tourism, and destructive fishing. Therefore, continued monitoring of seagrass conditions in Indonesian waters, especially in the Wondama Bay Regency, is needed for a more effective conservation approach to the seagrass ecosystem and its association with biota found in that ecosystem (Tebay *et al.*, 2020).

#### **Mollusc species composition and community structure**

This research is the first documentation on mollusc diversity in Wondama Bay. This study recorded a total of 197 benthic mollusc species from our sampling sites, with the species list presented in Table 2 and 3. Nevertheless, species accumulation curve did not approach an asymptote (Figure 3), indicating that the sampling effort ( $n=9$  stations) has not been sufficient to collect most species. Species diversity depends on the number of samples (Azovsky, 2011), suggesting that more species could be found in the Wondama Bay by increasing the number of sampling stations.

**Table 2.** The presence of mollusc Bivalvia from nine sites in Wondama Bay Regency

| Family         | Genus                | Station |   |   |   |   |   |   |   |   | Species number |
|----------------|----------------------|---------|---|---|---|---|---|---|---|---|----------------|
|                |                      | 1       | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |                |
| Arcidae        | <i>Anadara</i>       |         |   | + | + |   |   |   | + |   | 2              |
|                | <i>Tegillarca</i>    |         |   |   |   |   |   |   | + |   | 1              |
| Cardiidae      | <i>Fragum</i>        |         |   | + |   | + |   |   | + | + | 2              |
|                | <i>Vasticardium</i>  | +       | + | + | + | + | + | + | + |   | 6              |
| Chamidae       | <i>Chama</i>         |         |   |   |   |   |   |   | + |   | 1              |
| Cyrenoididae   | <i>Polymesoda</i>    |         | + |   | + |   |   |   |   |   | 1              |
| Glycymerididae | <i>Tucetona</i>      |         |   |   |   |   |   |   | + |   | 1              |
| Isognomonidae  | <i>Isognomon</i>     |         | + |   |   |   |   |   |   |   | 1              |
| Lucinidae      | <i>Codakia</i>       |         |   |   |   |   |   | + |   | + | 2              |
|                | <i>Anodontia</i>     |         |   | + |   |   |   | + |   |   | 1              |
|                | <i>Fimbria</i>       |         |   |   |   |   |   | + |   |   | 1              |
| Mactridae      | <i>Mactra</i>        |         | + | + |   |   | + |   |   |   | 2              |
| Malleidae      | <i>Malleus</i>       |         |   |   |   |   |   |   |   | + | 1              |
| Margaritidae   | <i>Pinctada</i>      |         |   |   |   |   |   | + | + |   | 1              |
| Mytilidae      | <i>Modiolus</i>      |         |   |   |   |   |   |   | + |   | 1              |
| Nuculanidae    | <i>Nuculana</i>      |         |   |   | + |   |   |   | + |   | 1              |
| Ostreidae      | <i>Crassostrea</i>   |         |   |   |   |   |   |   | + |   | 1              |
|                | <i>Ostrea</i>        |         |   | + |   |   |   |   |   |   | 1              |
| Pharidae       | <i>Pharella</i>      |         |   | + |   |   |   |   |   |   | 1              |
|                | <i>Siliqua</i>       |         | + |   |   |   |   |   |   |   | 1              |
| Psammobiidae   | <i>Asaphis</i>       |         |   | + |   |   |   |   |   |   | 1              |
|                | <i>Gari</i>          |         |   | + |   |   |   |   |   |   | 1              |
| Pteriidae      | <i>Pteria</i>        |         | + |   |   |   |   |   |   |   | 1              |
| Spondylidae    | <i>Spondylus</i>     |         |   | + |   |   | + | + | + | + | 2              |
| Tellinidae     | <i>Quidnipagus</i>   |         |   |   |   | + |   |   |   |   | 1              |
|                | <i>Scutarcopagia</i> | +       |   | + |   | + |   |   |   |   | 1              |
|                | <i>Tellina</i>       | +       | + | + | + | + | + | + | + | + | 1              |
|                | <i>Tellinella</i>    | +       |   | + | + | + | + |   |   | + | 1              |
|                | <i>Antigona</i>      |         |   | + |   |   |   | + |   |   | 1              |
| Veneridae      | <i>Dosinia</i>       | +       | + |   | + | + |   |   | + |   | 2              |
|                | <i>Gafrarium</i>     |         |   | + |   |   |   |   | + |   | 1              |
|                | <i>Lioconcha</i>     | +       | + |   | + | + | + |   |   | + | 4              |
|                | <i>Meretrix</i>      |         |   | + |   | + |   |   |   | + | 1              |
|                | <i>Pitar</i>         |         | + | + | + |   |   |   |   | + | 3              |
|                | <i>Timoclea</i>      |         |   |   | + | + |   |   |   |   | 1              |

The lowest mollusc diversity was found at station 1, with only 23 species, while the highest was found at stations 3 and 8, with 74 species (Figure 4). The species richness of molluscs from class Bivalvia was 53 species (Table 2), which is lower than that of Gastropoda with 144 species (Table 3). Several studies on the diversity of molluscan fauna in the Papua region has been reported such as seven gastropod species in mangrove forests of Nabire (Maitindom *et al.*, 2024), 63 gastropods in coastal area of Liki island (Sujarta *et al.*, 2022), 177 gastropods and 62 bivalves in Padaido islands (Aji *et al.*, 2018), 123 gastropods and 33 bivalves in Biak island (Aji and Widyastuti, 2020), 530 gastropods, and 159 bivalves in Raja Ampat (Wells, 2002). The high number of species recorded by Wells (2002) compared to the other regions in Papua is due to the use of a rapid assessment method across 44 sites, covering all habitats and employing diving activities in coral reef ecosystems. This suggests that mollusk species in Wondama Bay Regency still need further

exploration, particularly in coral reefs, which could reveal additional species.

The most commonly found bivalve species across the sampling sites (9 stations) of Wondama Bay was *Tellina* sp. and the second was *Tellinella virgata*, which was found in 6 stations. The substrate in our sampling areas is mostly fine sediment or sandy mud, suitable for those tellinid clams. The *Tellina* sp. and *Tellinella virgata* belong to the family Tellinidae (Huber, 2010). Tellinid clams specialize in deep burrowing and typically have narrow, thin shells. They usually position themselves horizontally in the sediment, lying on the left valve (Arruda *et al.*, 2003; Rueda *et al.*, 2009). They function as deposit feeders, supported by a long and mobile inhalant siphon that captures deposited particles and collects organic material from their surroundings (Arruda *et al.*, 2003; Rueda *et al.*, 2009). Among gastropods, *Mitrella scripta* was the only species found at all sites.

*Canarium urceus* occurred at eight stations, while other species found at seven stations included *Polinices aurantius*, *Nassarius albescens*, *Nassarius globosus*, and *Natica fasciata*. These widely distributed gastropods in Wondama Bay represent different trophic groups and feeding types within the ecosystem. The *Mitrella scripta*, *Polinices aurantius*, and *Natica fasciata* act as predators, *Canarium*

*urceus* as grazers and detritus feeders, and *Nassarius albescens* and *Nassarius globosus* as scavengers (Molluscabase eds, 2024). This indicates that molluscs play a role in the ecosystem as herbivores, omnivores, and carnivores, facilitating the transfer of energy from lower trophic levels, such as producers, into higher trophic levels (Rueda et al., 2009; Huber, 2010; Aji et al., 2023).

**Table 3A.** The presence of mollusc Gastropoda from nine sites in Wondama Bay Regency.

| Family         | Genus                 | Station |   |   |   |   |   |   |   |   | Species Number |
|----------------|-----------------------|---------|---|---|---|---|---|---|---|---|----------------|
|                |                       | 1       | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |                |
| Acteonidae     | <i>Pupa</i>           |         |   | + |   |   |   |   |   |   | 1              |
| Ancillariidae  | <i>Ancilla</i>        | +       |   |   |   |   |   |   |   |   | 1              |
| Angariidae     | <i>Angaria</i>        |         |   |   |   |   | + |   | + |   | 1              |
| Bullidae       | <i>Bulla</i>          |         |   |   |   |   |   |   | + | + | 1              |
| Bursidae       | <i>Tutufa</i>         |         |   |   |   |   |   |   | + |   | 1              |
| Cerithiidae    | <i>Cerithium</i>      | +       | + | + | + | + | + | + | + | + | 5              |
|                | <i>Pseudovertagus</i> |         |   |   |   |   |   | + |   |   | 1              |
|                | <i>Rhinoclavis</i>    |         |   | + | + |   | + |   | + |   | 3              |
| Chilodontaidae | <i>Euchelus</i>       |         |   |   |   |   |   |   | + |   | 1              |
|                | <i>Herpetopoma</i>    |         |   |   |   |   |   |   | + |   | 1              |
| Columbellidae  | <i>Mitrella</i>       | +       | + | + | + | + | + | + | + | + | 2              |
| Conidae        | <i>Conus</i>          |         |   |   |   |   | + |   | + |   | 5              |
| Costellariidae | <i>Vexillum</i>       | +       | + | + | + | + | + |   | + | + | 8              |
| Cymatiidae     | <i>Monoplex</i>       | +       |   |   |   | + |   |   |   |   | 2              |
| Cypraeidae     | <i>Cypraea</i>        |         |   |   | + |   |   | + | + |   | 2              |
|                | <i>Mauritia</i>       |         |   |   |   |   | + | + |   | + | 2              |
|                | <i>Monetaria</i>      |         |   | + |   |   | + | + | + | + | 1              |
|                | <i>Palmadusta</i>     |         |   |   |   |   |   | + |   |   | 1              |
| Ellobiidae     | <i>Ellobium</i>       |         |   | + | + |   |   |   |   |   | 1              |
|                | <i>Pythia</i>         |         |   |   |   |   | + |   |   |   | 1              |
|                | <i>Tralia</i>         |         |   |   |   | + |   |   |   |   | 1              |
| Fissurellidae  | <i>Amathina</i>       |         |   |   |   | + |   |   |   |   | 1              |
|                | <i>Emarginula</i>     |         |   | + |   |   |   |   |   |   | 1              |
| Haminoeidae    | <i>Aliculastrum</i>   |         |   | + |   | + |   |   | + | + | 1              |
|                | <i>Alys</i>           |         |   | + |   |   |   |   | + |   | 2              |
| Liotiidae      | <i>Liotinaria</i>     |         |   |   |   |   |   |   | + |   | 1              |
| Littorinidae   | <i>Littoraria</i>     | +       | + | + | + | + | + | + | + |   | 4              |
|                | <i>Littorina</i>      |         |   | + |   | + |   |   |   |   | 1              |
| Melongenidae   | <i>Thalessa</i>       |         |   |   |   |   |   |   |   | + | 1              |
| Mitridae       | <i>Imbricaria</i>     |         |   |   |   | + |   |   |   |   | 1              |
|                | <i>Nebularia</i>      |         |   |   |   |   |   | + |   |   | 1              |
|                | <i>Pterygia</i>       |         |   |   |   | + |   |   |   |   | 1              |
| Muricidae      | <i>Chicoreus</i>      |         | + | + |   |   |   |   |   |   | 1              |
|                | <i>Drupella</i>       |         |   | + |   |   |   |   |   |   | 1              |
|                | <i>Ergalatax</i>      |         |   |   |   |   |   |   | + |   | 1              |
|                | <i>Mancinella</i>     |         | + |   |   |   |   |   | + |   | 1              |
|                | <i>Morula</i>         |         |   |   |   |   |   |   | + | + | 1              |
|                | <i>Semiricinula</i>   |         | + | + |   |   |   |   | + |   | 2              |
|                | <i>Taurasia</i>       |         | + |   |   |   | + |   |   |   | 1              |
| Nassariidae    | <i>Nassarius</i>      | +       | + | + | + | + | + | + | + | + | 12             |
| Naticidae      | <i>Mammilla</i>       |         |   |   |   |   |   | + |   |   | 1              |
|                | <i>Natica</i>         |         |   | + | + | + | + | + | + | + | 3              |
|                | <i>Naticarius</i>     |         |   |   |   | + |   | + | + |   | 1              |
|                | <i>Notocochlis</i>    |         |   |   |   |   | + |   |   |   | 1              |
|                | <i>Polinices</i>      |         | + | + | + | + | + | + | + |   | 6              |

**Table 3B.** The presence of mollusc Gastropoda from nine sites in Wondama Bay Regency.

| Family            | Genus               | Station |   |   |   |   |   |   |   |   | Species number |
|-------------------|---------------------|---------|---|---|---|---|---|---|---|---|----------------|
|                   |                     | 1       | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |                |
| Neritidae         | <i>Nerita</i>       | +       | + | + |   |   | + | + | + |   | 10             |
|                   | <i>Neritodryas</i>  |         |   |   |   |   |   |   | + |   | 1              |
|                   | <i>Vittina</i>      |         |   | + |   |   |   |   |   |   | 1              |
| Olividae          | <i>Oliva</i>        |         |   | + | + | + | + | + | + |   | 7              |
| Ovulidae          | <i>Ovula</i>        |         |   |   |   |   |   |   | + |   | 1              |
| Phasianellidae    | <i>Phasianella</i>  |         |   | + |   | + |   |   | + |   | 1              |
| Pisaniidae        | <i>Cantharus</i>    | +       |   |   |   |   |   |   |   |   | 1              |
|                   | <i>Polia</i>        |         |   |   | + |   |   |   |   |   | 1              |
|                   | <i>Engina</i>       |         |   |   |   |   | + |   |   |   | 1              |
| Planaxidae        | <i>Fissilabia</i>   |         |   | + |   |   |   |   |   |   | 1              |
| Potamididae       | <i>Telescopium</i>  |         |   |   | + |   |   |   |   |   | 1              |
|                   | <i>Terebralia</i>   |         | + | + | + |   |   |   | + |   | 1              |
| Pseudomelatomidae | <i>Inquisitor</i>   |         |   |   |   | + |   |   |   |   | 1              |
| Pyramidellidae    | <i>Milda</i>        |         |   | + | + |   | + |   | + | + | 1              |
| Seraphsidae       | <i>Terebellum</i>   |         |   |   |   |   |   |   | + |   | 1              |
| Strombidae        | <i>Canarium</i>     | +       |   | + | + | + | + | + | + | + | 3              |
|                   | <i>Conomurex</i>    |         |   |   |   |   | + |   |   | + | 1              |
|                   | <i>Euprotomus</i>   |         | + |   | + |   |   |   |   |   | 1              |
|                   | <i>Gibberulus</i>   |         |   |   | + | + | + | + | + |   | 2              |
|                   | <i>Harpago</i>      |         |   |   |   |   |   | + |   |   | 1              |
|                   | <i>Lambis</i>       |         |   |   | + |   |   |   |   |   | 1              |
| Tegulidae         | <i>Tectus</i>       |         |   |   |   |   |   |   | + | + | 1              |
| Terebridae        | <i>Oxymeris</i>     |         |   | + |   |   | + |   | + |   | 3              |
|                   | <i>Terebra</i>      |         | + |   |   | + |   |   |   |   | 1              |
| Tonnidae          | <i>Tonna</i>        |         |   |   |   | + |   |   |   |   | 1              |
| Trochidae         | <i>Trochus</i>      |         |   |   |   |   |   |   |   | + | 1              |
|                   | <i>Monodonta</i>    |         |   | + |   |   |   | + | + |   | 2              |
|                   | <i>Rochia</i>       |         |   |   |   |   |   | + | + |   | 1              |
| Turbinidae        | <i>Astralium</i>    |         |   |   |   |   |   |   | + |   | 1              |
|                   | <i>Lunella</i>      |         |   | + |   |   |   |   | + |   | 1              |
|                   | <i>Turbo</i>        |         |   |   |   |   | + |   |   |   | 1              |
| Turridae          | <i>Unedogemmula</i> |         | + | + |   | + | + | + | + |   | 1              |
|                   | <i>Turris</i>       |         |   |   |   |   |   |   | + |   | 1              |
| Volutidae         | <i>Cymbiola</i>     |         |   |   |   |   | + |   | + | + | 1              |

Furthermore, it was noticed that class Bivalvia consisted of 19 families, represented by Arcidae, Cardiidae, Chamidae, Cyrenoididae, Glycymerididae, Isognomonidae, Lucinidae, Mactridae, Malleidae, Margaritidae, Mytilidae, Nuculanidae, Ostreidae, Pharidae, Psammobiidae, Pteriidae, Spondylidae, Tellinidae, and Veneridae. Meanwhile, Gastropoda consisted of 40 families, including Acteonidae, Ancillariidae, Angariidae, Bullidae, Bursidae, Cerithiidae, Chilodontidae, Columbellidae, Conidae,

Costellariidae, Cymatiidae, Cypraeidae, Ellobiidae, Fissurellidae, Haminoeidae, Liotiidae, Littorinidae, Melongenidae, Mitridae, Muricidae, Nassariidae, Naticidae, Neritidae, Olividae, Ovulidae, Phasianellidae, Pisaniidae, Planaxidae, Potamididae, Pseudomelatomidae, Pyramidellidae, Seraphsidae, Strombidae, Tegulidae, Terebridae, Tonnidae, Trochidae, Turbinidae, Turridae, and Volutidae. All of those mollusc families found in Wondama Bay are commonly found in tropical coastal areas and play

different roles in the ecosystems (Dharma, 2005; Arbi, 2011; Huber, 2015; Heryanto, 2018; Islami et al., 2018; Molluscabase eds, 2024).

As can be seen from Figure 5, there was no correlation between mollusc species richness and seagrass percent cover ( $r = 0.37$ ,  $p = 0.32$ ) and seagrass species richness ( $r = 0.035$ ,  $p = 0.93$ ) and temperature ( $r = -0.45$ ,  $p = 0.22$ ), nor between seagrass species richness and temperature ( $r = 0.078$ ,  $p = 0.84$ ). Next, the Non-metric multidimensional scaling (NMDS) ordination plot illustrates the variation in mollusc assemblage composition across sites (Figure 6). *Cymbiola vespertilio* and *Monetaria annulus* were likely indicator species at stations 8 and 9, while *Cerithium citrinum* was mainly found at stations 1, 2, and 5. Snail *Nerita* sp. was found only at stations 1 and 2,

and *Natica fasciata* was present at most sites but absent from stations 1 and 2. Additionally, envfit analysis indicated that temperature ( $r^2 = 0.2046$ ,  $p = 0.508$ ), seagrass species richness ( $r^2 = 0.5945$ ,  $p = 0.069$ ), geographic location UTM\_Northing ( $r^2 = 0.6225$ ,  $p = 0.053$ ), and UTM\_Easting ( $r^2 = 0.5739$ ,  $p = 0.086$ ) did not significantly impact mollusc species composition across sites. However, seagrass percent cover ( $r^2 = 0.7896$ ,  $p = 0.003$ ) was identified as a prominent driver of mollusc community assemblages.

The temperatures in Wondama Bay are relatively consistent across the sites and are suitable for benthic molluscs. Tropical mollusc species can even thrive in areas with temperatures as high as  $36^\circ\text{C}$  (Huhn, 2016; Aji et al., 2023). The impact of temperature is likely to affect species richness and composition of benthic organisms over a long monitoring




Figure 3. Species accumulation curve of mollusc species in Wondama Bay Regency.






Figure 4. The number of mollusc species from nine sampling areas in Wondama Bay Regency. Note: the total species number of Bivalvia and Gastropoda classes is displayed inside the bar graph.



**Figure 5.** Pearson correlations among (A) mollusc species richness and seagrass cover, (B) mollusc species richness and seagrass species richness, (C) mollusc species richness and temperature, and (D) seagrass species richness and temperature.



**Figure 6.** Non-metric multidimensional scaling (NMDS) ordination plot of mollusc assemblages from nine sites in Wondama Bay Regency, West Papua, Indonesia, based on Jaccard distances between sites. Mollusc species that have significant influence on reshaping the species composition among sites are shown *Cerithium citrinum* ( $r^2 = 0.6928$ ,  $p = 0.028$ ), *Nerita* sp. ( $r^2 = 0.7818$ ,  $p = 0.021$ ), *Natica fasciata* ( $r^2 = 0.7818$ ,  $p = 0.021$ ), *Monetaria annulus* ( $r^2 = 0.7893$ ,  $p = 0.006$ ), *Cymbiola vespertilio* ( $r^2 = 0.7787$ ,  $p = 0.029$ ), *Oliva guttata* ( $r^2 = 0.7226$ ,  $p = 0.046$ ). The arrows represent environmental (temperature), geographic location (UTM\_Northing and UTM\_Easting), and habitat characteristics (seagrass cover and seagrass species).

period with increasing temperature (Aji *et al.*, 2024). Temperature fluctuation influences benthic molluscs composition, though this is more common in temperate regions (Rueda *et al.*, 2008), as tropical areas have steady temperatures year-round. Similarly, geographic location did not influence mollusc composition, as the sites were too close. Wondama Bay is part of the Indo-West Pacific, where most mollusc species are widely distributed in tropical regions with similar climatic conditions, suggesting that the mollusc species found at our nine sites represent mollusc species from the Indo-West Pacific (Huber, 2010; Molluscabase eds, 2024).

Interestingly, only seagrass cover influenced mollusc composition, but not species richness. Additionally, the number of seagrass species did not determine mollusc richness and composition. This suggests that the number of mollusc species and their composition across our study sites could be influenced by human activities. Molluscs are a valuable source of income for local communities, providing food and materials for the ornamental and shellcraft industries (Islami *et al.*, 2018; Aji and Widyastuti, 2020). Thus, mollusc species richness may decrease because of the rising human consumption of protein from molluscs and the growing number of shell collectors. Benthic molluscs, such as gastropods and bivalves, are easily collected from their habitats, especially during low tide. As a result, large numbers of molluscs can be collected. Hence, if marine conservation programs are not properly managed, this could lead to a decline in mollusc diversity and reduce their population. Future research should consider measuring the impact of human activities in Wondama Bay, as these may influence mollusc species richness and composition within the ecosystems. Molluscs are important biota due to their role in the coastal ecosystems and their economic contribution to human life (Arbi, 2011; Eisenbarth *et al.*, 2018; Islami *et al.*, 2018).

## Conclusion

This study identified nine seagrass species in total, with *Enhalus acoroides* and *Cymodocea rotundata* present at all sampling sites. The mollusc species found in Wondama Bay were relatively diverse, totaling 197 species, including 53 bivalves and 144 gastropods. The most widespread bivalve species was *Tellina* sp., while *Mitrella scripta* was the most common gastropod found at all sites. Seagrass species, temperature, and geographic location had no significant influence on mollusc species composition across the sites. However, seagrass percent cover did affect mollusc community assemblages among sites. Future research should consider the impact of human activities, as this may

be a key factor influencing mollusc and seagrass composition in Wondama Bay Regency, West Papua. This present study collected samples in 2016 that may not represent the current conditions (the year 2025), as the distribution of molluscs may change due to seagrass and environmental changes as occurred. Climate change and anthropogenic (human activity) pressures are the two major stressors that have severely impacted coastal ecosystems, including seagrass beds. Nevertheless, our study could serve as the first assessment of molluscs diversity in the seagrass ecosystems of Wondama Bay. The results of this study can, therefore, be used as baseline data for monitoring seagrass ecosystems and their associated biota (e.g. molluscs) in the bay. Future sample collection during the wet and dry seasons is recommended to analyze seasonal variations in mollusc community compositions.

## Acknowledgement

We sincerely thank the local government of Wondama Bay Regency, West Papua. We also extend our appreciation to all the staff at the Technical Implementation Unit (TIU) for Marine Life Conservation in Biak, Papua, for their invaluable assistance. This work was fully supported and funded by TIU's annual research fund (DIPA-2016) for Marine Life Conservation Biak, Papua – Indonesian Institute of Science (LIPI).

## References

- Abbott, R.T., & Dance. S.P. 2000. Compendium of Seashells: A Full-Color Guide To More Than 4,200 Of The World's Marine Shells. Odyssey Publishing. 411 pp.
- Aji, L.P., Goud J., van der Steeg, S., Tapilatu, R., Maas, D.L., & Becking, L.E. 2023. The Diversity of Molluscan Faunas in marine Lakes of Raja Ampat, West Papua, Indonesia. *Contrib. Zool.*, 92: 1–40. <https://doi.org/10.1163/18759866-bja10047>
- Aji, L.P., Maas, D.L., Capriati, A., Ahmad, A., De Leeuw, C., & Becking, L.E. 2024. Shifts in Dominance of Benthic Communities Along a Gradient of Water Temperature and Turbidity in Tropical Coastal Ecosystems. *PeerJ*, 12: e17132. <https://doi.org/10.7717/peerj.17132>
- Aji, L.P., & Widyastuti, A. 2017. Molluscs Diversity in coastal Ecosystem of South Biak, Papua. *Oseanologi dan Limnologi di Indonesia*, 2: 25–37. <https://doi.org/10.14203/oldi.2017.v2i1.75>
- Aji, L.P., & Widyastuti, A. 2020. The Condition and Composition of Seagrass and Mollusca on Biak

- Island, Papua. *IOP Conf. Ser. Earth Environ. Sci.*, 404(1): p.012069. <https://doi.org/10.1088/1755-1315/404/1/012069>
- Aji, L.P., Widyastuti, A., & Capriati, A. 2018. Struktur Komunitas Moluska di padang Lamun Perairan Kepulauan Padaido dan Aimando Kabupaten Biak Numfor, Papua. *Oseanologi dan Limnologi di Indonesia*, 3: p.219. <https://doi.org/10.14203/oldi.2018.v3i3.184>
- Alianto, Hendri, & Suhaemi. 2018. Kelimpahan dan Kelompok Fitoplankton di Perairan Luar teluk Wondama, Propinsi Papua Barat. *J. Ilmu dan Teknologi Kelautan Tropis*, 10: 683–698. <https://doi.org/10.29244/jitkt.v10i3.19561>
- Arbi, U.Y. 2011. Struktur Komunitas Moluska di Padang Lamun Perairan Pulau Talise, Sulawesi Utara. *Oseanologi dan Limnologi di Indonesia*, 37: 71–88.
- Arruda, E.P., Domaneschi, O., & Amaral, A.C.Z 2003. Mollusc Feeding Guilds on Sandy Beaches in Sa˜o Paulo State, Brazil. *Mar. Biol.*, 143: 691–701. <https://doi.org/10.1007/s00227-003-1103-y>
- Azovsky, A.I. 2011. Species-Area and Species-Sampling Effort Relationships: Disentangling the Effects. *Ecography*, 34: 18-30. <https://doi.org/10.1111/j.1600-0587.2010.06288.x>
- Becking, L.E., Martinez, S.J., Aji, L.P., Ahmad, A., Alzate, A., Folkers, M., Lestari, D.F., Subhan, B., & Hoeksema B.W. 2024. Stony Corals And Their Associated Fauna Residing in Marine Lakes Under Extreme Environmental Conditions. *Diversity*, 16(5): p.295. <https://doi.org/10.3390/d16050295>
- Dharma, B. 2005. Recent and fossil Indonesian shells, ConchBooks. 424 pp.
- Dharmawan, I.W.E., & Widyastuti, A. 2017. Pristine Mangrove Community in Wondama Gulf, West Papua, Indonesia. *Mar. Res. Indonesia*, 42: 73–82. <https://doi.org/10.14203/mri.v42i2.175>
- Eisenbarth, J.H., Undap, N., Papu, A., Schillo, D., Dialao, J., Reumschiessel, S., Kaligis, F., Bara, R., Schaberle, T.F., Konig, G.M., Yonow, N., & Wagele, H. 2018. Marine Heterobranchia (Gastropoda, Mollusca) in Bunaken National Park, North Sulawesi, Indonesia - A follow-up Diversity Study. *Diversity*, 10: p.127. <https://doi.org/10.3390/d10040127>
- Heryanto. 2018. Benthic Molluscs Communities in the Intertidal Coast of Tanimbar Islands, West Southeast Mollucas. *J. Biologi Indonesia*, 14(1): 73–80. <https://doi.org/10.47349/jbi/14012018/73>
- Hoeksema, B.W. 2007. Delineation of the Indo-Malayan Centre Of Maximum Marine Biodiversity: the Coral Triangle, p. 117–178. In Willem Renema [ed.], Biogeography, Time, And Place: Distributions, Barriers, and Islands. Springer Netherlands.
- Huber, M. 2010. Compendium of Bivalves. A Full-Color Guide to 3,300 of the World's Marine Bivalves. A status on Bivalvia after 250 years of research. ConchBooks. 901 pp.
- Huber, M. 2015. Compendium of Bivalves 2. A full-Color Guide to The Remaining Seven Families. A Systematic Listing of 8.500 Bivalve Species and 10.500 Synonyms, ConchBooks. 907 pp.
- Huhn, M. 2016. The Relevance of Food Availability for the Tolerance to Environmental Stress in Asian Green Mussels, *Perna viridis*, from Coastal Habitats in Indonesia. Thesis. Christian-Albrechts-University. Kiel Germany. 136 pp.
- Islami, M. M., Ikhsani, I.Y., Indrabudi, T., & Pelupessy, I.A.H. 2018. Diversity, Composition and Utilization of Mollusc in Saparua Island, Center Molucca. *Widyariset*, 4: 173. <https://doi.org/10.14203/widyariset.4.2.2018.173-188>
- Jaccard, P. 1908. Nouvelles Recherches Sur La Distribution Florale. *Bull. Soc. Vaud. Sci. Nat.* 44: 223–270.
- Kilminster, K., McMahon, K., Waycott, M., Kendrick, G.A., Scanes P., McKenzie, L., O'Brien, K.R., Lyons, M., Ferguson, A., Maxwell, P., Glasby, T., & Udy, J. 2015. Unravelling Complexity in Seagrass Systems for Management: Australia as a microcosm. *Sci. Total Environ.* 534: 97–109. <https://doi.org/10.1016/j.scitotenv.2015.04.061>
- Lee, K.S., Park, S.R., & Kim, Y.K. 2007. Effects of Irradiance, Temperature, and Nutrients on Growth Dynamics of Seagrasses: A review. *J. Exp. Mar. Biol. Ecol.*, 350: 144–175. <https://doi.org/10.1016/j.jembe.2007.06.016>
- Legendre, P., & Gallagher, E.D. 2001. Ecologically Meaningful Transformations for Ordination of Species Data. *Oecologia*, 129: 271–280. <https://doi.org/10.1007/s004420100716>
- Madiyani, K.D.P., Triastuti, J., & Pursetyo. K.T. 2018. Inventory of the Tropical Coral Reef Fishes in Wondama Bay Regency, West Papua, Indonesia. *IOP Conf. Ser. Earth Environ. Sci.*, 137(1):

- p.012095. <https://doi.org/10.1088/1755-1315/137/1/012095>
- Maitindom, F.A., Mamangkey, G.F., Boneka, F.B., Lasut, M.T., Mantiri, R.O.S., & Bara, R. 2024. Mollusks Attached to Natural Mangrove Roots and Silvo-fisheries in Nabire Papua: to What Extent are They Different. *Platax*, 12: 224–235. <https://doi.org/10.35800/jip.v10i2.57758>
- Mangubhai, S., Erdmann, M.V., Wilson, J.R., Huffard, C.L., Ballamu, F., Hidayat, N.I., Hitipeuw, C., Lazuardi, M.E., Muhajir, Pada, D., Purba, G., Rotinsulu, C., Rumetna, L., Sumolang, K., & Wen, W. 2012. Papuan Bird's Head Seascape: Emerging Threats and Challenges in the Global Center of Marine Biodiversity. *Mar. Pollut. Bull.*, 64: 2279–2295. <https://doi.org/10.1016/j.marpolbul.2012.07.024>
- McKenna, S.A., Allen, G.R., & Suryadi, S. 2002. A Marine Rapid Assessment of the Raja Ampat Islands, Papua Province, Indonesia. RAP Bulletin of Biological Assessment 22. Conservation International.
- van der Meij, S.E.T., Moolenbeek, R.G., & Hoeksema, B.W. 2009. Decline of the Jakarta Bay Molluscan Fauna Linked to Human Impact. *Mar. Pollut. Bull.*, 59: 101–107. <https://doi.org/10.1016/j.marpolbul.2009.02.021>
- Molluscabase eds. 2024. Molluscabase. Available at <https://www.molluscabase.org>. Accessed 10 September 2024. MolluscaBase.
- Nugraha, A.H., Tasabaramo, I.A., Hernawan, U.E., Rahmawati, S., Putra, R.D., & Darus, R.F. 2021. Diversity, Coverage, Distribution and Ecosystem Services of Seagrass in Three Small Islands of Northern Papua, Indonesia: Liki Island, Meossu Island and Befondi Island. *Biodiversitas*, 22: 5544–5549. <https://doi.org/10.13057/biodiv/d221238>
- Oksanen, J.F., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., & Wagner, H. 2019. Vegan: community ecology package (version 2.5-6).
- Palomares, M.L.D., & Heymans, J.J. 2006. Historical ecology of the Raja Ampat Archipelago, Papua Province, Indonesia.
- Paramasivam, K., & Venkataraman, K. 2022. Comparison of Macro-Molluscan Assemblages in a Protected and a Non-Protected Tropical Seagrass Ecosystem. *Biotropica*, 54(6): 1491–1513. <https://doi.org/10.1111/btp.13153>
- R Core Team. 2023. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rahmawati, S., Irawan, A., Supriyadi, I.H., & Azkab, H. 2017. Panduan pemantauan penilaian kondisi padang lamun, 2nd ed. Coremap CTI Lembaga Ilmu Pengetahuan Indonesia.
- Rahmawati, S., Lisdayanti, E., Kusnadi, A., Rizqi, M.P., Putra, I.P., Irawan, A., Supriyadi, I.H., Prayudha, B., Suyarso, Alifatri, L.O., Iswari, M.Y., Anggraini, K., Hadiyanto, Hernawan, U.E., Ambo-Rappe, R., Choesin, D.N., Nugraha, A.H., Sjafrie, N.D.M., Riniatsih, I., Rifai, H., Fachriansyah, K., Manafi, M.R., Rustam, A., Ningsih, E., & Rahmadi, P. 2022. Status ekosistem lamun di Indonesia Tahun 2021, Pusat Riset Oseanografi, Organisasi Riset Kebumihan dan Maritim. Badan Riset dan Inovasi Nasional. 94 pp.
- Real, R., & Vargas, J.M. 1996. The Probabilistic Basis of Jaccard's Index of Similarity. *Syst. Biol.*, 45(3): 380–385
- Rizqi, M.P., Souhoka, J., Makatipu, P., Aji, L.P., Kusnadi, A., Nurdiansah, D., Patty, S., & Dzumalek, A.R. 2019. Monitoring kesehatan terumbu karang dan ekosistem terkait suaka alam perairan Raja Ampat Provinsi Papua Barat, Coremap CTI - Lembaga Ilmu Pengetahuan Indonesia. 68 pp.
- Rondonuwu, A.B., Lumingas, L.J.L., Bataragoa, N.E., Manengkey, H.K., Kondoy, K.I.F., Rangan, J.K., Makatipu, P., Nurdiansah, D., Tampanguma, B., & Kambey, A.G. 2019. Monitoring kesehatan terumbu karang dan ekosistem terkait lainnya di Pulau Batanta dan Pulau Salawati, Raja Ampat, Propinsi Papua Barat. Coremap CTI - Lembaga Ilmu Pengetahuan Indonesia. 131 pp.
- Rueda, J.L., Gofas, S., Urra, J., & Salas, C. 2009. A Highly Diverse Molluscan Assemblage Associated with Eelgrass Beds (*Zostera marina* L.) in the Alboran Sea: Micro-habitat Preference, Feeding Guilds and Biogeographical Distribution. *Sci. Mar.*, 73(4): 679–700. <https://doi.org/10.3989/scimar.2009.73n4679>
- Rueda, J.L., Urra, J., & Salas, C. 2008. Diel and Seasonal Variation of a Molluscan Taxocoenosis Associated with a *Zostera marina* Bed in Southern Spain (Alboran Sea). *Helgoland Mar. Res.*, 62: 227–240. <https://doi.org/10.1007/s10152-008-0111-1>
- Schultze-Westrum, T. 2001. West-Papua: Only the Village People Can Save Their Reefs and Rainforests. *Biodiversity*, 2: 15–19. <https://doi.org/10.1080/14888386.2001.9712531>

- Sujarta, P., Mailissa, M.G., Keiluhu, H.J., Hadisusanto, S., & Yuliana, S. 2022. Community Distribution and Utilization of Gastropods in the Coastal Area of Liki Island , Sarmi District, Papua, Indonesia. *Biodiversitas*, 23: 5001–5011. <https://doi.org/10.13057/biodiv/d231006>
- Suruan, S., Pranata, B., Tania, C., & Kamal, M.M. 2016. Photo ID-based Assessment of the Whale Shark (*Rhincodon typus*) Population in Kwatisore, Wondama Bay, West Papua, Indonesia. *QScience Proceedings (The 4th International Whale Shark Conference)*, <http://doi.org/10.5339/qproc.2016.iwsc4.61>
- Tebay, S., Boli, P., & Ainusi, J. 2020. Potensi Lamun di Kampung Aisandami Kabupaten Teluk Wondama dan Strategi Pengelolaanya. *Sumberdaya Akuatik Indopasifik*, 4: 111–128. <https://doi.org/10.46252/jsai-fpik-unipa>
- Wells, F. 2002. Molluscs of the Raja Ampat islands, Papua Province, Indonesia. In: A marine rapid assessment of the Raja Ampat islands, Papua Province, Indonesia. RAP Bulletin of Biological Assessment 22, Conservation International.
- Wickham, H. 2016. Package “ggplot2”: elegant graphics for data analysis.
- Wilson, B.R., Wilson, C., & Baker, P. 1993. Australian Marine Shells: Prosobranch Gastropods, part one, Vol.1. University of Western Australia Press. 408 pp.