

The Use of *Diopatra claparedii* Grube, 1878 (Onuphidae, Polychaeta) as a Bioindicator of Heavy Metal Contaminations in the Donan Creek, Cilacap, Central Java, Indonesia

Sri Lestari^{1*}, Windiariani Lestari², Eko Setio Wibowo³, Phuping Sucharitakul⁴, Joko Pamungkas⁵

¹Ecotoxicology Laboratory, Faculty of Biology, Jenderal Soedirman University

²Ecology Laboratory, Faculty of Biology, Jenderal Soedirman University

³Animal Physiology Laboratory, Faculty of Biology, Jenderal Soedirman University

Jl. Dr. Soeparno No.63 Grendeng. Kec. Purwokerto Utara, Banyumas, Jawa Tengah 53122 Indonesia

⁴Batavia Coast Maritime Institute

Separation Point Close 133, City of Greater Geraldton - Beachlands, Australia

⁵Directorate of Scientific Collection Management, National Research and Innovation Agency

Gedung B.J. Habibie, Jl. M.H. Thamrin No. 8, Jakarta Pusat 10340

Email: srilestari.bio@unsoed.ac.id

Abstract

The Donan Creek is a coastal ecosystem in Central Java that is strongly affected by industrial activities. This study represents the first attempt to use the polychaete worms *Diopatra claparedii* Grube, 1878 (Onuphidae) as a bioindicator of heavy metal contaminations. The work was conducted between July and September 2023 at four sampling sites, designated from the farthest to the closest to the Indian Ocean. Samples of *D. claparedii*, sediment and water were collected at each site during low tide. The samples were analyzed for Cd, Cr and Pb concentration using Atomic Absorption Spectrophotometry. In addition, several other environmental parameters were also measured to examine the water quality of the creek. Our study highlights the capability of *D. claparedii* to accumulate all the three heavy metals. The concentrations of Cd, Cr and Pb in the body tissues of the worms ranged from $29.743 \pm 0.011 \text{ mg.kg}^{-1}$ to $50.036 \pm 0.028 \text{ mg.kg}^{-1}$, from $21.718 \pm 0.024 \text{ mg.kg}^{-1}$ to $40.032 \pm 0.052 \text{ mg.kg}^{-1}$, and from $20.068 \pm 0.008 \text{ mg.kg}^{-1}$ to $44.030 \pm 0.000 \text{ mg.kg}^{-1}$, respectively. These amounts were so far the highest among all other aquatic animals (i.e. bivalves, crustaceans and fishes) used as bioindicators in the Donan Creek. The heavy metal concentrations in *D. claparedii* were significantly correlated with those in the sediment ($p < 0.05$). In general, the levels of the metals were considerably higher in the more industrial area than those of the less industrial area ($P < 0.05$). We further found that the concentrations of Cd, Cr and Pb as well as several other environmental parameters including BOD, COD, the total nitrate and phosphate in the water column of the creek did not meet the water quality standards stipulated by the Indonesian Government. These findings indicate that the ecosystem has been polluted by both heavy metals and organic matters. While environmental remediation is urgently required, a comprehensive review and improvement of industrial and domestic waste management practices in the region are equally essential to prevent further degradation.

Keywords: Annelida, marine pollution, monitoring, water contamination

Introduction

The Donan Creek is an estuary administratively located in Cilacap Regency, Central Java Province, Indonesia. The creek, which is connected to the Indian Ocean, is often considered the easternmost part of the Segara Anakan Lagoon (SAL) and plays a crucial role in the local coastal ecosystem. Along with the mangrove area, it provides essential habitats for many estuarine species (e.g. Nordhaus *et al.* 2009; Pamungkas, 2013) and holds economic importance for the local residents, particularly in transportation, fisheries, recreational and industrial activities (pers. obs.).

Due to intensive anthropogenic activities on land, the Donan Creek has been reportedly receiving various contaminants, mainly through sewages. Dzikowitzky *et al.* (2011) reported the occurrence of more than 50 contaminants – they were mostly polycyclic aromatic compounds (PACs) consisted primarily of polycyclic aromatic hydrocarbons (PAHs) – in the water, sediment and macrobenthic fauna inhabiting the eastern part of the SAL, including the Donan Creek. Previous studies also indicated that the eastern part of the lagoon was contaminated by heavy metals (e.g. Hernawati, 2014; Widjianto, 2014; Alam, 2015; Syakti *et al.*, 2015; Piranti *et al.*, 2020). The heavy metals, which typically come from

industries, pose a significant threat to the coastal water ecosystem as well as human health as they cannot be degraded by organisms – the metals accumulate in the environment, mainly settling at the bottom of waters to form more complex compounds (e.g. Arifin et al., 2012; Piwowarska et al., 2024; Shintianata et al., 2024; Shetaia et al., 2025).

Benthic animals living at the bottom of aquatic ecosystems are commonly used as bioindicators due to their ability to accumulate heavy metals in their body tissues, being sessile with limited mobility, and ease of collection (Mir et al., 2021). Bioindicator species are commonly used in biomonitoring contaminant exposure. Locally, the bivalves *Geloina expansa* (Mousson, 1849) or 'kerang totok' in the local name – the species is also known by the unaccepted species names *Geloina erosa* (e.g. Irawati et al., 2018) and *Polymesoda erosa* (e.g. Nordhaus et al., 2009) – have been used as a bioindicator for heavy metal contaminations in the eastern part of the SAL (Irawati et al., 2018) due to their widespread distribution in the lagoon and economic importance (the bivalves are consumed by the locals).

Another benthic species that has the potential for use as a bioindicator of heavy metal contaminations in the Donan Creek is the tubicolous polychaete worms *Diopatra claparedii* Grube, 1878 (Onuphidae). This species commonly occurs in several Southeast Asian countries and is often misidentified as *Diopatra neapolitana* Delle Chiaje, 1841 (e.g. Paxton, 2002; Idris and Arshad, 2013). The occurrence of this species in Indonesia, along with its economic importance, was first documented by Pamungkas et al. (2023) based on the specimens collected from the Donan Creek. Some biological aspects of this species have been studied by Wibowo et al. (2022; 2023; 2025). *Diopatra claparedii* is by far one of the biggest polychaete species in the SAL (pers. obs.), making it ideal for use as a bioindicator. While *D. neapolitana* has been reported to be a useful bioindicator for coastal contaminations (e.g. Mdaini et al., 2021; Arias et al., 2024) like some other polychaete species (e.g. Dean, 2008; Kies et al., 2020; Elías et al., 2021; Stefano et al., 2021), the potential of *D. claparedii* as an indicator species has not been investigated. In the present study, we used *D. claparedii* as an indicator species of heavy metal contaminations, especially Cd, Cr and Pb, and examined the water quality of the Donan Creek.

Materials and Methods

The present study was carried out in the Donan Creek, Cilacap, Central Java, between July and September 2023. Four sampling sites were designated from the furthest to the closest site from

the sea. Site 1 ($7^{\circ}39'54.6''S$, $109^{\circ}00'45.2''E$), Site 2 ($7^{\circ}40'46.3''S$, $109^{\circ}00'23.2''E$), Site 3 ($7^{\circ}42'22.5''S$, $108^{\circ}59'24.2''E$) and Site 4 ($7^{\circ}43'40.2''S$, $108^{\circ}59'15.2''E$) were situated near mangrove areas, a cement factory, an oil refinery and Sleko Harbour, respectively (Figure 1A). Site 1 and 4 are less industrial, whereas Site 2 and 3 are more industrial.

Sample collections

At each sampling site, samples of *D. claparedii*, water and sediment were collected three times during a low tide. The worms were collected by a local experienced worm digger from the bottom of the Donan Creek. The tip of the animal's tube was first palpated using one foot. The tube, with the animal inside, was then taken out of the sediment using bare hands and was put on a thin wall container. The details on how a local digger collected the worms were documented by Pamungkas (2023) – see <https://www.youtube.com/watch?v=1cBQx1yDOqU>. On the boat, the worms obtained were removed from their tube by massaging the posterior part of the tube anterior wards. The worms were placed in a smaller thin wall container (Figure 2A). Water samples were taken using 1 L plastic bottles, and approximately 1 kg of sediment samples were taken using hands covered with plastic gloves and were put in plastic bags. Both samples were labelled and stored in a cool box for further analyses in the laboratory.

Species identification and other bioindicators

The worms collected from the field were identified in the laboratory under both stereo and compound microscopes using the identification keys of Paxton (2002), Budaeva and Fauchald (2011), Idris and Arshad (2013), as well as the description of the local species by Pamungkas et al. (2023). This identification step was essential to ensure if the worms obtained were *D. claparedii* (Figure 2B). While the presence of spiralled branchiae and peristomial cirri confirmed the identity of the genus, the presence of distinct pectinate chaetae with funnel-like combs confirmed the identity of the species.

For comparison, we also compiled a list of all the species that have been used as bioindicators of heavy metal contaminations in the Donan Creek, along with the information about their capacity to accumulate the metals from relevant literature.

Environmental parameter measurements

Three heavy metals, i.e. cadmium (Cd), chromium (Cr) and lead (Pb), were measured from the body of *D. claparedii*, as well as the water and the sediment of the Donan Creek. These three elements,




Figure 1. Map of the study area. Red circles with numbers indicate sampling sites (basic map was provided by Erwin R. Ardli)



Figure 2. A. Worm removal from the tube. B. Morphology of a living *Diopatra claparedii* collected from the Donan Creek

along with arsenic (As) and mercury (Hg), are globally recognized as priority metals of public health concern due to their high toxicity (Tchounwou *et al.*, 2012). Worm samples were first dried in the sun then air dried overnight. The drying process was then continued in the oven at a temperature of 60°C for 24h. Thereafter, a 5 g worm sample was ashed using a furnace at a temperature of 600°C for 4 h. The ash obtained was then cooled in a desiccator. The worm ash was put into a digestion flask and was added with 5 mL of concentrated HNO₃ and 2 mL of H₂O₂ until the ash was completely dissolved. The solution was then filtered using a Whatman grade 42 ashless filter paper (Badan Standarisasi Nasional, 2021). Using a 50 mL volumetric flask, the filtrate was diluted with distilled water to 50 mL. The filtrate was analyzed for Cd, Cr and Pb concentration using Atomic Absorption Spectrophotometry (AAS) with wavelengths of 228.8, 357.9 and 283.3 nm, respectively.

Sediment samples were first dried in the sun then air dried overnight. The drying process was then continued in the oven at a temperature of 105°C for 24 h. Thereafter, the sediment was crushed in a porcelain cup, then sieved. A 20 g sediment sample was put into an Erlenmeyer tube added with 500 mL of distilled water, then was centrifuged at 2000 rpm for 30 min. The water phase was removed and the precipitate was dried in the oven at a temperature of 105°C for 24 h. Afterwards, a 1 g sediment sample was put in a glass beaker added with 6 mL of aqua regia, then heated on a hotplate at a temperature of 103°C until the solution was almost dry (Badan Standarisasi Nasional, 2021). After the sample was cooled, 1 mL of concentrated HNO₃ was added. The solution was then filtered using a Whatman grade 42 ashless filter paper. Using a 10 mL volumetric flask, the filtrate was diluted with distilled water to 10 mL. The filtrate was analyzed for Cd, Cr and Pb concentration using AAS with wavelengths of 228.8, 357.9 and 283.3 nm, respectively.

A 50 mL water sample was taken and put into a 100 mL glass beaker. The sample was added with 10 mL of concentrated HNO₃ and 5 mL of 20% HCl then was heated on a hotplate at a temperature of 120°C until the solution became clear. Afterwards, the solution was filtered using a Whatman grade 42 ashless filter paper. Using a 50 mL volumetric flask, the filtrate was diluted with distilled water to 50 mL (Badan Standarisasi Nasional, 2019). The filtrate was analyzed for Cd, Cr and Pb concentration using AAS with wavelengths of 228.8, 357.9 and 283.3 nm, respectively.

Bioconcentration factor (BCF), *i.e.* the uptake of the heavy metals from both the sediment and the water by the worms was calculated using the

equation $BCF = C_{dc}/C_{s \text{ or } w}$. Here, C_{dc} is the heavy metal concentration in *D. claparedii* (mg.kg⁻¹), whereas C_s and C_w are the heavy metal concentrations in the sediment (mg.kg⁻¹) and the water (mg.L⁻¹), respectively. BCF values smaller, equal and bigger than 1 indicate that the worms contain less, equal and higher metals than the environment, respectively. The BCF is an important concept in the environmental risk assessment since it gives quantitative information regarding the ability of a contaminant to be taken up by organisms from the water (Wang, 2016). In addition, concentration factor (CF), *i.e.* the uptake of the heavy metals from the water by the sediment, was calculated using the equation $CF = C_s/C_w$. CF values smaller, equal and bigger than 1 indicate that the sediment contains less, equal and higher metals than the water, respectively.

Various environmental parameters were measured to examine the water quality of the Donan Creek. The parameters included temperature, conductivity, salinity, pH, total dissolved solids (TDS), total suspended solids (TSS), dissolved oxygen (DO), biochemical oxygen demand (BOD), chemical oxygen demand (COD), total nitrate and total phosphate. Water quality parameters such as temperature, conductivity, salinity, pH, TDS, TSS, BOD and COD are essential to understand in relation to heavy metal analysis because they strongly influence the behavior, mobility, and bioavailability of metals in aquatic systems. all the data obtained were then compared to the Water Quality Standards for Marine Biota set in the Government Regulation No. 22 Year 2021, Appendix VIII.

Statistical analysis

As the data obtained are not normally distributed (the Shapiro-Wilk test was performed to inspect the data normality), the non-parametric Spearman correlation analysis was used to evaluate the relationships among the heavy metal concentrations in the body tissues of *D. claparedii*, the water and the sediment. The Mann-Whitney U test was also performed to compare the concentrations of the metals between the less industrial area (Sites 1 and 4) and the more industrial area (Sites 2 and 3). The statistical analyses were conducted using R version 4.4.0. coupled with RStudio version 2024.04.1+748.

Results and Discussion

This study demonstrated that the concentrations of Cd, Cr and Pb in the body of *D. claparedii*, as well as in the water and the sediment of the Donan Creek, varied among four sampling sites. However, they were generally higher in the more

industrial area than those in the less industrial area ($P < 0.05$; Table 1; Figure 3).

The worms collected from Site 2 (near the cement factory) accumulated the highest amount of Cd ($50.036 \pm 0.028 \text{ mg.kg}^{-1}$), Cr ($40.032 \pm 0.052 \text{ mg.kg}^{-1}$) and Pb ($44.030 \pm 0.000 \text{ mg.kg}^{-1}$), followed by those collected from Site 3 (near the oil refinery). At the latter site, the amounts of Cd, Cr and Pb in the body of the worms were $33.050 \pm 0.000 \text{ mg.kg}^{-1}$, $38.050 \pm 0.047 \text{ mg.kg}^{-1}$ and $40.053 \pm 0.005 \text{ mg.kg}^{-1}$, respectively. Lower levels of heavy metals were found at Sites 1 and 4, which were less industrial (Table 1 and Figure 3).

In the water column, the levels of Cd and Cr gradually increased from Site 1 to 4. Specifically, Cd increased from $2.808 \pm 0.008 \text{ mg.L}^{-1}$ to $6.723 \pm 0.007 \text{ mg.L}^{-1}$, and from $0.561 \pm 0.025 \text{ mg.L}^{-1}$ to $2.042 \pm 0.053 \text{ mg.L}^{-1}$ for Cr. Slightly different from these two metals, the level of Pb increased from Site 1 ($0.601 \pm 0.010 \text{ mg.L}^{-1}$) to 3 ($2.751 \pm 0.018 \text{ mg.L}^{-1}$), then declined to $1.845 \pm 0.010 \text{ mg.L}^{-1}$ at Site 4 (Table 1 and Figure 3).

In the sediment, the metal concentrations increased from Site 1 to 3, then decreased at Site 4. The highest amounts of Cd ($78.181 \pm 0.058 \text{ mg.kg}^{-1}$), Cr ($48.117 \pm 0.022 \text{ mg.kg}^{-1}$) and Pb ($72.899 \pm 0.010 \text{ mg.kg}^{-1}$) were hence recorded at Site 3, whereas the lowest concentrations were at Site 1 (Table 1 and Figure 3).

The higher concentrations of the heavy metals in the more industrial area seem to be associated with the presence of the cement factory (near Site 2) and the oil refinery (near Site 3) that regularly discharge effluent into the creek (pers. obs). Typically, Cd, Cr and Pb resulted from the cement industry are sourced from limestone, clay, fly ash, additives such as steel slag, and fossil fuels, whereas in the oil refinery industry, Cd comes from crude oil or catalysts in the reforming, hydrocracking and desulfurization processes; Cr comes from pipe materials, tanks or metal catalysts; and Pb comes from heavy oil fractions and anti-knock additives in gasoline.

As can be seen in Table 1 and Figure 3, the amounts of Cd, Cr and Pb in the body of *D. claparedii* ranged from $29.743 \pm 0.011 \text{ mg.kg}^{-1}$ to $50.036 \pm 0.028 \text{ mg.kg}^{-1}$, from $21.718 \pm 0.024 \text{ mg.kg}^{-1}$ to $40.032 \pm 0.052 \text{ mg.kg}^{-1}$, and from $20.068 \pm 0.008 \text{ mg.kg}^{-1}$ to $44.030 \pm 0.000 \text{ mg.kg}^{-1}$, respectively. These findings show that the worms are resistant to the three metals and possess the ability to absorb metallic elements similar to the European species *D. neapolitana* (e.g. Freitas et al., 2012; Arias et al., 2024). The concentrations of metals accumulated in the body of *D. claparedii* were found to be much

higher than those in other groups of aquatic animals inhabiting the Donan Creek such as bivalves, crustaceans and fishes (Table 2). These suggest that the species has the highest ability to absorb metal contaminations from the surrounding environment.

This study further showed that the heavy metal concentrations in the body of *D. claparedii* had a strong positive correlation with those of the sediment of the Donan Creek ($R = 0.76$; $P = 9.6e-08$; Figure 4A), but a weak positive correlation with those of the water ($P > 0.05$; Figure 4B). The strong positive correlation between the metals in the body of *D. claparedii* and those in the sediment can be attributed to the life and the feeding habits of aquatic animals (e.g. Arifin et al., 2012). The adult *D. claparedii* is benthic infaunal organisms dwelling in the sediment of the Donan Creek for their entire lives (Pamungkas et al., 2023). The worms are continuously exposed to heavy metal contaminants that settle in the sediment (the amount of the metals in the water column positively correlated ($R = 0.45$; $P = 0.0059$; Figure 4C) with that in the sediment), resulting in the accumulation of metals in their body tissues. The metals also enter the worms' body through their dietary uptake. While *Diopatra* worms are known to be omnivorous – they feed on aquatic organisms living on the tube cap, on neighbouring tube caps, and within nearby sediments (Berke, 2022) – several studies have showed that a number of aquatic animals in the Donan Creek were reportedly contaminated with heavy metals (see Table 2).


Understanding how *D. claparedii* resists the toxic effects of heavy metals requires a further investigation. However, previous studies show that some polychaete species have developed detoxification mechanisms. For example, they may produce metal-binding proteins such as metallothioneins (MTs) or metallothionein-like proteins (MTLPs), which chelate and neutralize toxic metals within their tissues (Zhao et al., 2020). They may also regulate their internal metal concentrations by inhibiting the uptake of metals or promoting their excretion (Sun and Zhou, 2007). Despite this, morphological abnormalities in *D. claparedii* inhabiting the Donan Creek – the abnormalities are likely to be linked to heavy metal exposure – were documented by Pamungkas et al. (2025), indicating that the deleterious effects of the metals may not be entirely avoidable.

This study further showed that the BCFs (dc-s) were close to 1, i.e. ranging from 0.547 to 1.770, suggesting that the concentrations of the heavy metals in both the body of the worms and the sediment were nearly equal (values below 1 may be linked to the young age of the animals). In general, these findings demonstrate that *D. claparedii* can be used as a bioindicator. The BCFs (dc-w) ranged from

Table 1. Environmental parameters at each sampling site

Parameter	Site ¹				GR ²
	1	2	3	4	
Heavy metals					
Cd in worm (mg.kg ⁻¹)	32.064 ± 0.037	50.036 ± 0.028	33.050 ± 0.000	29.743 ± 0.011	-
in sediment (mg.kg ⁻¹)	27.989 ± 0.008	57.875 ± 0.006	78.181 ± 0.058	27.081 ± 0.057	-
in water (mg.L ⁻¹)	2.808 ± 0.008	4.512 ± 0.012	6.379 ± 0.008	6.723 ± 0.007	0.001
Cr in worm (mg.kg ⁻¹)	22.065 ± 0.046	40.032 ± 0.052	38.050 ± 0.047	21.718 ± 0.024	-
in sediment (mg.kg ⁻¹)	20.983 ± 0.104	47.818 ± 0.011	48.117 ± 0.022	37.838 ± 0.100	-
in water (mg.L ⁻¹)	0.561 ± 0.025	0.981 ± 0.003	1.923 ± 0.010	2.042 ± 0.053	0.005
Pb in worm (mg.kg ⁻¹)	20.068 ± 0.008	44.030 ± 0.000	40.053 ± 0.005	24.748 ± 0.007	-
in sediment (mg.kg ⁻¹)	16.441 ± 0.032	48.27 ± 0.010	72.899 ± 0.010	13.979 ± 0.073	-
in water (mg.L ⁻¹)	0.601 ± 0.010	1.030 ± 0.001	2.751 ± 0.018	1.845 ± 0.010	0.008
Others					
Temperature (°C)	29.233 ± 0.971	29.967 ± 0.586	29.633 ± 1.35	29.267 ± 0.551	28-32
Conductivity (µS.cm ⁻¹)	41.9 ± 7.996	44.667 ± 13.65	47.5 ± 10.259	46.733 ± 13.683	-
Salinity (ppt)	22.967 ± 2.001	24.7 ± 7.856	25.2 ± 4.453	26.133 ± 7.727	up to 34
pH	7.623 ± 0.186	7.75 ± 0.105	7.693 ± 0.047	7.773 ± 0.107	7-8.5
TDS (mg.L ⁻¹)	22.233 ± 6.161	26.333 ± 10.42	24.933 ± 7.06	23.1 ± 7.206	-
TSS (mg.L ⁻¹)	20.667 ± 3.215	20.667 ± 1.528	20.333 ± 3.215	20.667 ± 2.309	80
DO (mg.L ⁻¹)	5.693 ± 0.508	8.083 ± 2.144	7.937 ± 1.842	8.02 ± 1.647	> 5
BOD (mg.L ⁻¹)	393 ± 136.121	381.667 ± 190.306	352.667 ± 195.677	291 ± 110.964	20
COD (mg.L ⁻¹)	605.667 ± 231.738	596.333 ± 301.291	537.667 ± 295.392	450.333 ± 171.844	-
Total nitrate (mg.L ⁻¹) ³	< 0.5/< 0.04/< 0.2	< 0.5/< 0.04/< 0.2	< 0.5/< 0.04/< 0.2	< 0.5/< 0.04/< 0.2	0.06
Total phosphate (mg.L ⁻¹) ⁴	< 0.04/< 0.5/< 0.06	< 0.04/< 0.5/< 0.07	0.3/< 0.5/< 0.1	< 0.4/< 0.5/< 0.06	0.015

¹ The data are presented as the average value ± standard deviation; values in bold have exceeded the threshold set by the Indonesian Government. ² GR = Government Regulation No. 22 Year 2021, Appendix VIII. ^{3,4} The three values at each sampling site measured in July, August, September, respectively

**Figure 3.** Concentrations of heavy metals at each sampling site

10.634 to 42.748, indicating that the concentrations of the metals in the body of the worms were considerably higher than those of the water column. Similarly, the CFs (s-w) ranged from 7.577 to 48.761,

showing that the metal concentrations in the sediment were much higher than those of the water column (Table 3). Lower heavy metal concentrations in the water column appear to be related to their fate

Table 2. Heavy metal concentrations in the body of aquatic animals inhabiting the Donan Creek

Species ¹	Common name	Local name	Heavy metals (mg.kg ⁻¹) ³			Reference
			Cd	Cr	Pb	
Bivalves						
<i>Geloina expansa</i> (Mousson 1849)	Broad geloina	Kerang totok	0.0645	0.0495	-	Annisa (2023)
<i>Geloina expansa</i> (Mousson 1849)	Broad geloina	Kerang totok	0.117 ± 0.043	-	0.130 ± 0.059	Irawati et al. (2018)
Crustaceans						
<i>Alpheus</i> sp. ²	Snapping/pistol shrimp	<i>Udang pletok</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Macrobrachium equidens</i> (Dana 1852) ²	Rough river prawn	<i>Udang wuku</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Metapenaeus</i> sp. ²	-	<i>Udang jahe</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Metapenaeus</i> sp. ²	-	<i>Udang peci</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Oratosquilla fabricii</i> (Holthuis 1941) ²	Mantis shrimp	<i>Udang sikat</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Oratosquillina interrupta</i> (Kemp 1911) ²	Blackspot mantis shrimp	<i>Udang sikat</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Parapenaeopsis coromandelica</i> Alcock 1906 ²	Coromandel shrimp	<i>Udang jahe</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Penaeus monodon</i> Fabricius 1798 ²	Asian tiger shrimp	<i>Udang topus/windu</i>	Ud	-	Ud	Tumisem and Pusapawiningtiyas (2011)
<i>Scylla serrata</i> (Forskål 1775)	Mangrove crab	<i>Kepiting bakau</i>	0.030	-	< 0.001	Sastranegara et al. (2021)
Fishes						
<i>Mugil</i> sp.	Flathead grey mullet	<i>Ikan belanak</i>	0.2115	-	-	Afifah (2019)
<i>Mugil cephalus</i> Linnaeus 1758	Flathead grey mullet	<i>Ikan belanak</i>	-	-	2.05	Ningrum (2006)
<i>Sillago sihama</i> (Forskål 1775)	Northern whiting	<i>Ikan rejung</i>	0.564	-	9.194	Cahyani et al. (2016)
Polychaetes						
<i>Diopatra claparedii</i> Grube 1878	-	<i>Lur umah</i>	50.036 ± 0.028	40.032 ± 0.052	44.030 ± 0.000	This study

¹ Based on the accepted species name on the World Register of Marine Species (www.marinespecies.org)

² The authors only provided local name; the scientific name was given by a crustacean taxonomist, Rena T. Hernawati, based merely on the photograph of the animal provided in the literature. Taxonomic investigations based on specimens collected from the Donan Creek may be required to verify the identification.

³ Highest value in the animal provided in the literature; the symbol '-' indicates that the heavy metal was not measured; 'Ud' indicates undetected value measured using the thresholds 0.025 mg.kg⁻¹ (Pb) and 0.01 mg.kg⁻¹ (Cd); values in bold are the highest.

Table 3. Concentration factors at each sampling site

Factor*	Site 1**			Site 2***			Site 3***			Site 4**		
	Cd	Cr	Pb	Cd	Cr	Pb	Cd	Cr	Pb	Cd	Cr	Pb
BCF (dc-s)	1.146	1.052	1.221	0.865	0.837	0.912	0.549	0.791	0.549	1.770	0.574	1.770
BCF (dc-w)	11.420	39.318	33.410	11.089	40.821	42.748	14.561	19.790	14.561	13.413	10.634	13.413
CF (s-w)	9.969	37.389	27.372	12.826	48.761	46.864	26.502	25.026	26.502	7.577	18.527	7.577

* BCF = bioconcentration factor; CF = concentration factor; dc = *Diopatra claparedii*; s = sediment; w = water; ** Less industrial

*** More industrial

in the water environment: the metals sink and accumulate in the sediment, or are absorbed by aquatic organisms.

This study further found that the levels of the three heavy metals in the water column of the Donan

Creek at all sampling sites, i.e. ranging between $0.561 \pm 0.025 \text{ mg.L}^{-1}$ and $6.723 \pm 0.007 \text{ mg.L}^{-1}$, have significantly exceeded the threshold levels set by the Indonesian Government (Table 1). These findings are consistent with the results of the study conducted by Piranti et al. (2020) in the same area in 2018 – they




Figure 4. Spearman correlation analyses. R is the Spearman's correlation coefficient. The result is considered significant if $P < 0.05$. Solid red linear lines indicate positive correlation. Grey shading represents standard error

nonetheless reported lower concentrations of Cd (0.013–0.022 mg.L⁻¹), Cr (0.0165–0.0283 mg.L⁻¹) and Pb (0.02–0.03 mg.L⁻¹) – yet contradict Sastranegara *et al.* (2021). The latter authors, whose the investigation was done in 2015, suggested that the concentrations of Cd (0.001 mg.L⁻¹) and Pb (0.003 mg.L⁻¹) in the Donan Creek still met the national water quality standard. While the differences are in part because the sampling sites were not exactly the same, the data actually showed that the water quality of the creek has changed over the past decade: the coastal ecosystem has now been polluted by the metals, most likely from the effluent discharged by nearby factories.

Furthermore, whereas most of the other environmental parameters such as temperature, conductivity, salinity, pH, TDS, TSS and DO still met the national water quality standard, the high levels of both BOD (i.e. from 291 ± 110.964 mg.L⁻¹ to 393 ± 136.121 mg.L⁻¹) and COD (i.e. from 450.333 ± 171.844 mg.L⁻¹ to 605.667 ± 231.738 mg.L⁻¹) indicated high organic pollutions that occurred in the Donan Creek, most likely from anthropogenic activities on land. These findings are in line with what Dsikowitzky *et al.* (2011) documented that the eastern part of the SAL, including the Donan Creek, has been mainly contaminated by PAHs. The excessive amounts of both nitrate and phosphate in the present study (i.e. up to nearly 0.5 mg.L⁻¹), in particular, may accelerate eutrophication in the coastal ecosystem. These pollutants are mainly sourced from the locals' residential and agricultural areas as well as harbours and traditional markets situated right along the creek (see Figure 1), which discharge waste into the creek on a daily basis (pers. obs.).

Conclusion

The present study sheds light on the capability of *D. claparedii* to absorb heavy metal contaminants

from the environment. Being resistant to metal exposure, the species is a useful bioindicator in coastal waters and is by far a more effective species in accumulating metals than other aquatic organisms such as bivalves, crustaceans, and fishes. Further studies are nevertheless required to investigate where the metals are accumulated the most in the body of the species (e.g. digestive tract, skin, or even the tube – the analysis of heavy metal concentrations in the tube has been particularly challenging as the tube cannot be dried using the standard procedure). Our study also provides critical insights into the contamination occurring in the Donan Creek. The concentrations of three heavy metals, *i.e.* Cd, Cr and Pb, as well as several other environmental parameters including BOD, COD, and total nitrate and phosphate in the water column of the creek, did not meet the water quality standards set by the Indonesian Government. This suggests that the ecosystem has been polluted by both heavy metals and organic matter. As these metals are particularly toxic and can cause serious diseases in humans, environmental remediation is required. Furthermore, both industrial and domestic waste management systems in the area need to be thoroughly evaluated.

Acknowledgements

We would like to thank Muji Pramono for assisting us with the worm collections. The identification of crustacean species in Tumisem and Pusapawiningtiyas (2011) by Rena T. Hernawati (Research Center for Biosystematics and Evolution) is also appreciated. Constructive critical comments from an anonymous reviewer improved the quality of this manuscript, and are appreciated. Our gratitude is expressed to the Research and Community Service of Jenderal Soedirman University for the research grant provided under the *Riset Dasar* funding scheme (MoU no. 27.255/UN23.37/PT.01.03/II/2023). This research project was conducted in collaboration with the Research Centre for Biosystematics and

Evolution, National Research and Innovation Agency (MoU no. 92/V/KS/08/2022 and T/502/UN23.6/HK.06.00/2022).

References

Afifah, N.N., 2019. Akumulasi logam berat kadmium (Cd) pada media air, sedimen, dan ikan Belanak (*Mugil* sp.) di Sungai Donan, Cilacap. Thesis, Jenderal Soedirman University.

Alam, M., 2015. Kajian logam berat merkuri pada *Metapenaeus monocerus* yang tertangkap dan persepsi masyarakat sekitar Sungai Donan, Cilacap. Thesis, Jenderal Soedirman University.

Alcock, A., 1906. Catalogue of the Indian decapod Crustacea in the collections of the Indian Museum. Part III. Macrura. Fasciculus I. The prawns of the Penaeus group. Calcutta: Trustees of the Indian Museum.

Annisa, A.C.N., 2023. Analisis kandungan logam berat kadmium (Cd) dan kromium (Cr) pada Kerang Totok (*Polymesoda erosa*) di perairan Segara Anakan, Cilacap. Thesis, Jenderal Soedirman University.

Arias, A., Woodin, S.A. & Paxton, H., 2024. An introduction to Diopatra, the amazing ecosystem engineering polychaete. *Biology*, 12(7): 1027. <https://doi.org/10.3390/biology12071027>

Arifin, Z., Puspitasari, R. & Miyazaki, N., 2012. Heavy metal contamination in Indonesian coastal marine ecosystems: A historical perspective. *Coast. Mar. Sci.*, 35(1): 227-233. <https://doi.org/10.15083/00040657>

Badan Standarisasi Nasional, 2019. Cara uji kadar logam terlarut dan logam total secara Spektrofotometer Serapan Atom (SSA) (SNI 6989-84:2019). Jakarta.

Badan Standarisasi Nasional, 2021. Cara uji kadar logam dalam contoh uji limbah padat, sedimen dan tanah dengan metode destruksi asam menggunakan Spektrofotometer Serapan Atom (SSA) - nyala / Inductively Coupled Plasma Optical Emission Spectrometric (ICP-OES) (SNI 8910:2021). Jakarta.

Berke, S.K., 2022. A review of Diopatra ecology: current knowledge, open questions, and future threats for an ecosystem engineering polychaete. *Biology*, 11(10): 1485. <https://doi.org/10.3390/biology11101485>

Budaeva, N. & Fauchald, K., 2011. Phylogeny of the *Diopatra* generic complex with a revision of Paradiopatra Ehlers, 1887 (Polychaeta: Onuphidae). *Zool. J. Linn. Soc.*, 163(2): 319-436. <https://doi.org/10.1111/j.1096-3642.2011.00701.x>

Cahyani, N., Batu, D.T.F.L. & Sulistiono, 2016. Kandungan logam berat Pb, Hg, Cd, dan Cu pada daging ikan rejung (*Sillago sihama*) di estuari Sungai Donan, Segara Anakan timur, Cilacap, Jawa Tengah. *J. Pengolahan Hasil Perikanan Indonesia*, 19(3): 267-276. <https://doi.org/10.17844/jphpi.v19i3.14544>

Dana, J.D., 1852. *Conspectus crustaceorum*, etc. Conspectus of the Crustacea of the exploring expedition under Capt. C. Wilkes, U.S.N. Macroura. *Proc. Acad. Nat. Sci. Philadelphia*, 6: 10-28.

Dean, H.K., 2008. The use of polychaetas (Annelida) as indicator species of marine pollution: a review. *Rev. Biol. Trop.*, 56(4): 11-38.

Delle Chiaje, S., 1841. Descrizione e notomia degli animali invertebrati della Sicilia citeriore osservati vivi negli anni 1822-1830. Naples: Batteli and Co.

Dsikowitzky, L., Nordhaus, I., Jennerjahn, T.C., Khrycheva, P., Sivatharshan, Y., Yuwono, E. & Schwarzbauer, J., 2011. Anthropogenic organic contaminants in water, sediments and benthic organisms of the mangrove-fringed Segara Anakan Lagoon, Java, Indonesia. *Mar. Pollut. Bull.*, 62(4): 851-862. <https://doi.org/10.1016/j.marpolbul.2011.02.023>

Elías, R., Villamar, F., Laverde Castillo, J.J., Brauko, K.M., Araki Braga, M., Cunha Lana, P.D., Díaz Díaz, O., Méndez, N., Muniz, P., Cabanillas, R. & Gutiérrez Rojas, C., 2021. Polychaetes as biological indicators in Latin America and the Caribbean. *Mar. Fish. Sci.*, 34(1): 37-107. <https://doi.org/10.47193/mafis.3412021010301>

Fabricius, J.C., 1798. *Supplementum entomologiae systematicae*. Hafniae: Proft et Storck.

Forskål, P., 1775. *Descriptiones Animalium, Avium, Amphibiorum, Piscium, Insectorum, Vermium; quae in Itinere Orientali Observavit Petrus Forskål. Post Mortem Auctoris editit Carsten Niebuhr. Adjuncta est materia *Medica Kahirina*. Hafniae: Möller.*

Freitas, R., Pires, A., Quintino, V., Rodrigues, A.M. & Figueira, E., 2012. Subcellular partitioning of elements and availability for trophic transfer: Comparison between the bivalve *Cerastoderma edule* and the polychaete *Diopatra*

neapolitana. *Est. Coast. Shelf Sci.*, 99: 21-30. <https://doi.org/10.1016/j.ecss.2011.11.039>

Grube, A.E., 1878. Annulata Semperiana. Beiträge zur Kenntniss der Annelidenfauna der Philippinen nach den von Herrn Prof. Semper mitgebrachten Sammlungen. *Mém. Acad. Imp. Sci. St.-Pétersbourg* (série 7), 25(8): 1-300.

Hernawati, 2014. Kajian akumulasi logam berat kadmium pada keping bakau dan persepsi masyarakat terhadap pencemaran lingkungan di Segara Anakan, Cilacap. Thesis, Jenderal Soedirman University.

Holthuis, L.B., 1941. Biological results of the Snellius Expedition XII. The Stomatopoda of the Snellius Expedition. *Temminckia*, 6: 241-294.

Idris, I. & Arshad, A., 2013. Checklist of polychaetous annelids in Malaysia with redescription of two commercially exploited species. *Asian J. Anim. Vet. Adv.*, 8(3): 409-436. <https://doi.org/10.3923/ajava.2013.409.436>

Irawati, Y., Lumbanbatu, D.T.F. & Sulistiono, S., 2018. Logam berat kerang totok (*Geloina erosa*) di timur Segara Anakan dan barat Sungai Donan, Cilacap. *J. Pengolahan Hasil Perikanan Indonesia*, 21(2): 232-242. <https://doi.org/10.17844/jphpi.v21i2.22842>

Kemp, S., 1911. Preliminary descriptions of new species and varieties of Crustacea Stomatopoda in the Indian Museum. *Rec. Indian Mus.*, 6: 93-100.

Kies, F., Kerfouf, A., Elegbede, I., Matemilola, S., De Los Rios Escalante, P., Khorchani, A. & Savari, S., 2020. Assessment of the coastal and estuarine environment quality of western Algeria using the bioindicator Polychaeta; the genus *Nereis*. *J. Mater. Environ. Sci.*, 11(9): 1472-1481.

Linnaeus, C., 1758. *Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata*. Stockholm: Laurentii Salvii.

Mdaini, Z., Telahigue, K., Hajji, T., Rabeh, I., El Cafsi, M.H., Tremblay, R. & Gagné, J.P., 2021. Comparative biomarker responses to urban pollution in three polychaete species: *Perinereis cultrifera*, *Diopatra neapolitana*, and *Marphysa sanguinea* from the lagoon of Tunis. *Environ. Monit. Assess.*, 193(3): 119. <https://doi.org/10.1007/s10661-021-08911-8>

Mousson, A., 1849. Die land- und süsswasser-Mollusken von Java. Zurich: Friedrich Schulthess.

Mir, Z.A., Arafat, M.Y. & Bakhtiyar, Y., 2021. Benthic macroinvertebrates as bioindicators of water quality in freshwater bodies. In: Dar, G.H., Hakeem, K.R., Mehmood, M.A. & Qadri, H. (eds.) *Freshwater pollution and aquatic ecosystems*. New York: Apple Academic Press, pp. 165-184.

Ningrum, P.Y., 2006. Kandungan logam berat timbal (Pb) serta struktur mikroanatomik branchia, hepar, dan musculus ikan belanak (*Mugil cephalus*) di perairan Cilacap. Thesis, Sebelas Maret University.

Nordhaus, I., Hadipudjana, F.A., Janssen, R. & Pamungkas, J., 2009. Spatio-temporal variation of macrobenthic communities in the mangrove-fringed Segara Anakan lagoon, Indonesia, affected by anthropogenic activities. *Reg. Environ. Change*, 9(4): 291-313. <https://doi.org/10.1007/s10113-009-0097-5>

Pamungkas, J., 2013. Polychaete community composition of the Segara Anakan mangrove forest, Cilacap, Central Java, Indonesia. Thesis, University of Bremen.

Pamungkas, J., 2023. First record of *Diopatra claparedii* Grube, 1878 (Onuphidae, Polychaeta) from Indonesian waters. <https://www.youtube.com/watch?v=1cBQx1yDOqU&t=52s> [Accessed 30 May 2024].

Pamungkas, J., Atang & Wibowo, E.S., 2023. First record of *Diopatra claparedii* Grube, 1878 (Onuphidae, Polychaeta) from Indonesian waters, with some notes on its economic importance. *Treubia*, 50(1): 11-20. <https://doi.org/10.14203/treubia.v50i1.4485>

Pamungkas, J., Wibowo, E.S., Alam, M. & Lestari, S., 2025. Pollutants may have caused morphological abnormalities in some polychaete species (Annelida) collected from Cilacap, Central Java, Indonesia. *Trop. Life Sci. Res.*, 36(1): 297. <https://doi.org/10.21315/tlsr.2025.36.1.14>

Paxton, H., 2002. *Diopatra* Audouin and Milne Edwards (Polychaeta: Onuphidae) from Thailand. *Spec. Publ. Phuket Mar. Biol. Cent.*, 24: 101-114.

Piranti, A.S., Rahayu, D.R.U.S., Ardli, E.R., Setyaningrum, N., Widyartini, D.S. & Insan, I., 2020. Water quality status of Segara Anakan,

Cilacap, Indonesia, for biota life. *IOP Conf. Ser. Earth Environ. Sci.*, 593(1): 012038. <https://doi.org/10.1088/1755-1315/593/1/012038>

Piwowarska, D., Kiedrzyńska, E. & Jaszczyszyn, K., 2024. A global perspective on the nature and fate of heavy metals polluting water ecosystems, and their impact and remediation. *Crit. Rev. Environ. Sci. Technol.*, 54: 1436-1458. <https://doi.org/10.1080/10643389.2024.2309845>

Sastranegara, M.H., Lestari, W., Sudiana, E. & Nasution, E.K., 2021. Distribution and accumulation of heavy metals from waters and sediments to *Scylla serrata* in Segara Anakan, Cilacap. *IOP Conf. Ser. Earth Environ. Sci.*, 746(1): 012026. <https://doi.org/10.1088/1755-1315/746/1/012026>

Shetaia, S.A., Wang, Y., Marriner, N., Li, M., Zhao, X., Chen, J., Salem, A. & Chen, Z., 2025. Fate of heavy metals in the water-scarce Nile Delta lagoons: a post-Aswan Dam perspective. *Environ. Pollut.*, 126754. <https://doi.org/10.1016/j.envpol.2024.126754>

Shintianata, D., Lubis, A.A., Sugiharto, U., Andini, F.P., Meiliastri, S.S. & Elias, M.S. 2024. Tracing Heavy Metal Dynamics in Mangrove Sediments: A Study from Ujung Kulon National Park. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, 29 (4): 519-529 doi: 10.14710/ik.ijms.29.4.519-529

Stefano, R., Buono, D. & Santovito, A., 2021. Polychaetes as bioindicators of environmental pollution: Impact of bisphenol A on the reproduction rate of *Ophryotrocha diadema* (Åkesson, 1976) (Eunicida: Dorvilleidae). *Zool. Ecol.*, 31(1): 61-65. <https://doi.org/10.3551/321658005.2021.1.8>

Sun, F.H. & Zhou, Q.X., 2007. Metal accumulation in the polychaete *Hediste japonica* with emphasis on interaction between heavy metals and petroleum hydrocarbons. *Environ. Pollut.*, 149(1): 92-98. <https://doi.org/10.1016/j.envpol.2006.12.013>

Syakti, A.D., Demelas, C., Hidayati, N.V., Rakasiwi, G., Vassalo, L., Kumar, N., Prudent, P. & Doumenq, P., 2015. Heavy metal concentrations in natural and human-impacted sediments of Segara Anakan Lagoon, Indonesia. *Environ. Monit. Assess.*, 187(1): 4079. <https://doi.org/10.1007/s10661-014-4079-9>

Tchounwou, P.B., Yedjou, C.G., Patlolla, A.K. & Sutton, D.J., 2012. Heavy metal toxicity and the environment. In: Luch, A. (ed.) Molecular, clinical and environmental toxicology. Basel: Springer, pp. 133-164. https://doi.org/10.1007/978-3-7643-8340-4_6

Tumisem & Puspawiningtyas, E., 2011. Analisis kadar logam dan cara mudah mengenali udang yang terakumulasi logam: studi kasus tentang udang di Sungai Donan, Cilacap, Jawa Tengah. *J. Manusia dan Lingkungan*, 18(2): 111-126. <https://doi.org/10.22146/jml.18436>

Wang, W.X., 2016. Bioaccumulation and biomonitoring. In: Blasco, J., Chapman, P.M., Campana, O. & Hampel, M. (eds.) *Marine ecotoxicology: current knowledge and future issues*. London: Academic Press, pp. 99-119. <https://doi.org/10.1016/B978-0-12-803371-5.00004-7>

Wibowo, E.S., Puspitasari, I.G.A.A.R., Atang & Pamungkas, J., 2022. Biological aspects of *Diopatra* sp. (Onuphidae, Polychaeta) collected from mangrove habitats of Jeruklegi, Cilacap Regency. *Depik*, 11(3): 299-305. <https://doi.org/10.13170/depik.11.3.27195>

Wibowo, E.S., Pitriani, M.R., Rachmawati, F.N., Susilo, U., Setiyono, E., Sucharitakul, P. & Pamungkas, J., 2023. Sexual maturity and macronutrient contents in *Diopatra* sp. (Onuphidae, Polychaeta) maintained at different salinity levels. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, 28(1): 89-98. <https://doi.org/10.14710/ik.ijms.28.1.89-98>

Wibowo, E.S., Setiyono, E., Simanjuntak, S.B.I., Susilo, U., Sucharitakul, P., Apriyanti, Y. & Pamungkas, J., 2025. Biological aspects of *Diopatra claparedii* Grube, 1878 (Onuphidae, Polychaeta) maintained at different salinity levels. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, 30(1): 45-56. <https://doi.org/10.14710/ik.ijms.30.1.45-56>

Widiyanto, B., 2014. Kajian kandungan logam berat tembaga pada udang penaeid dan persepsi masyarakat terhadap pencemaran lingkungan di Segara Anakan, Cilacap. Thesis, Jenderal Soedirman University.

Zhao, H., Wang, M., Pang, M., Sun, J., Huang, Y., Gao, F., Yang, D., Zhou, Y. & Wang, B., 2020. Quantitative immunodetection of metalloprotein II in *Perinereis aibuhitensis* after heavy metal exposure. *Mar. Pollut. Bull.*, 160: 111550. <https://doi.org/10.1016/j.marpolbul.2020.111550>