

Manuskrip_JIL_ImeldaPurba

by Cek Turnitin

Submission date: 19-Mar-2024 12:14PM (UTC+0700)

Submission ID: 2324520759

File name: Manuskrip_JIL_ImeldaPurba.docx (92.98K)

Word count: 5712

Character count: 32086

Subjective Health Complaints in Women Farmers Exposed to Pesticides in Agricultural Area

Imelda Gernauli Purba^{1,2}, Haryoto Kusnopranto³

¹Doctoral Program in Public Health, Faculty of Public Health, Universitas Indonesia; e-mail: imelda@fkm.unsri.ac.id.

²Environmental Health Program, Faculty of Public Health, Universitas Sriwijaya

³Department of Environmental Health, Faculty of Public Health, Universitas Indonesia

ABSTRAK

Pestisida adalah bahan kimia yang semakin banyak digunakan terutama di bidang pertanian untuk mengendalikan hama tanaman. Pestisida tidak hanya memberikan manfaat tetapi juga menimbulkan dampak buruk bagi manusia, terutama jika tidak ditangani dengan baik. Wanita di daerah pertanian yang terpajan pestisida memiliki risiko mengalami gangguan kesehatan, terkait dengan keterlibatannya dalam kegiatan pertanian seperti penyemprotan, pencampuran pestisida, pencucian peralatan dan pakaian menyemprot. Penelitian ini bertujuan untuk menganalisis faktor risiko keluhan kesehatan subjektif akibat pajanan pestisida pada petani wanita. Jenis penelitian ini adalah observasional dengan pendekatan cross sectional. Populasi penelitian adalah petani wanita yang berdomisili di Kecamatan Sekayu, Musi Banyuasin Sumatera Selatan, dengan jumlah sampel 136, diambil dengan teknik Cluster. Pengumpulan data primer melalui wawancara dan observasi menggunakan kuesioner dan checklist. Analisis data menggunakan *Chi-Square Test*. Hasil penelitian menunjukkan bahwa keluhan kesehatan subjektif yang dialami oleh petani perempuan antara lain kelelahan, cemas, sakit kepala, pandangan kabur, mual, nafsu makan menurun, kelemahan otot, dan kejang otot. Beberapa variabel yang berhubungan dengan keluhan kesehatan subjektif antara lain umur ($p=0,05$), masa kerja ($p=0,002$), jumlah jenis pestisida ($p=0,000$), serta cara penyimpanan pestisida ($p=0,021$). Petani perempuan sebaiknya membatasi paparan pestisida dengan mengurangi lama kontak dengan pestisida, menghindari pencampuran berbagai jenis pestisida, dan menyimpan pestisida sesuai dengan tempat penyimpanannya.

Kata kunci: Pestisida, Petani wanita, Keluhan kesehatan subjektif

ABSTRACT

Pesticides are chemicals that are increasingly used especially in agriculture to control plant pests. Pesticides not only provide benefits but also cause adverse effects on humans, especially if not handled properly. Women in agricultural areas are a group at risk of health problems due to pesticide exposure, related to their involvement in agricultural activities such as spraying, mixing of pesticide, washing equipment and spraying clothes. This research aims to analyze the risk factors for subjective health complaints due to pesticide exposure in female farmers. This type of research was observational with a cross sectional approach. The population of the study was female farmers who resided in Sekayu District, Musi Banyuasin South Sumatra, Indonesia with a sample size of 136 taken by Cluster Sampling. Primary data collection through interviews and observations using questionnaires and checklists. Data were analyzed using Chi-Square Test. The results showed that the subjective health complaints experienced by female farmers included fatigue, anxiety, headache, blurred vision, nausea, decreased appetite, muscle weakness, and muscle spasms. Several variables were associated with subjective health complaints including age ($p=0.05$), working period ($p=0.002$), number of types of pesticides ($p=0.000$), and storage method of pesticides ($p=0.021$). Female farmers should limit pesticide exposure through reducing the length of contact with pesticides, avoid mixing different types of pesticides, and store pesticides according to storage.

Keywords: Pesticides, Female farmers, Subjective health complaints

Citation: Purba, Imelda Gernauli., Haryoto Kusnopranto, (2024). Subjectif Health Complaints Due to Pesticide Exposure Among Female Farmers in Agricultural Areas. *Jurnal Ilmu Lingkungan*, xx(x), xx-xx, doi:10.14710/jil.xx.x.xxx-xx

1. Introduction

Women play an important role in the agricultural sector, accounting for 39% globally (UN Woman,

2020) and 24% of women in Indonesia work in the agricultural (Kementerian Pemberdayaan Perempuan dan Perlindungan Anak, 2018). Involvement in

agricultural activities exposes Women to pesticides. Agricultural pesticide use is increasing, in 2019 there were 4.2 million tons (Mt) of active ingredients, while worldwide pesticide application per cropland area was 2.7 kg/ha. Total pesticide use increased in 2010 by more than 50 percent compared to the 1990s, with pesticide use per cropland area increasing from 1.8 to 2.7 kg/ha. In 2019 Asia used the most pesticides in the agricultural sector (2.2 Mt), both in terms of total and per ha of agricultural land (3.7 kg/ha) (FAO, 2021). Pesticides are used by 53.2% of people in Asia and 29.3% of people in the Americas (Zhang et al., 2019).

Pesticide exposure is quite high when mixing and applying pesticides, using doses that are not as directed, mixing several types of pesticides, not using complete personal protective equipment, and improper application techniques (Filippi et al., 2021). Exposure to Organophosphate pesticides can cause a decrease in blood cholinesterase levels (Nambunmee et al., 2021)(Utami et al., 2021). This is due to the inhibition of the acetylcholine enzyme breaking down acetic acid and choline in the central nervous system. Organophosphate poisoning symptoms include increased salivation, diarrhea, vomiting, muscle tremors, gastrointestinal disturbances, and confusion (Hughes et al., 2021). Pesticide poisoning affects 500,000-1,000,000 people and kills 5,000-20,000 people worldwide each year with two-thirds of these incidents occurring in developing countries (Hadian et al., 2019). Studies in Tanzania mentioned organophosphate pesticide poisoning of 28.7% of which occurred in adolescent women (Lekei et al., 2020). Poor agricultural practices in developing countries are a major source of dermal and oral pesticide poisoning (Kumari & John, 2019).

Sekayu District, located in Musi Banyuasin Regency, South Sumatra, is an area where most of the population relies on agriculture as a source of income. Women play an important role in the agricultural or plantation sector in Sekayu Sub-district. Women are at risk of pesticide exposure through agricultural activities, especially mixing, spraying, cleaning spraying equipment, and washing spraying clothes. The risk of exposure is even greater due to incomplete use of personal protective equipment and poor storage of pesticides in the home. Pesticide exposure can cause various health problems such as symptoms of pesticide poisoning. A preliminary survey of female farmers in Sekayu sub-district, Musi Banyuasin, revealed subjective complaints related to pesticide exposure. Starting from this problem, so the aim of this research was to analyze the risk factors for subjective health complaints due to pesticide exposure in female farmers.

2. Methods

This research was conducted after first passing the ethical review by the ethics commission of FKM Unsri, the respondents were given informed consent and signed it before the interview and observation. The design of this study was cross sectional to analyze the

relationship between pesticide exposure and effects at the same time. The population of this study were all female farmers who live in Sekayu District, Musi Banyuasin. The sample of this study is part of the female farmers who live in Sekayu District Musi Banyuasin, totaling 136 people. The sampling technique used cluster sampling. The sample criteria were female farmers aged 17-45 years, not pregnant, and did not have a history of certain diseases. The type of data in this study is primary data obtained through interviews and observations including age, length of exposure, number of pesticide types, and storage methods. Data processing was carried out with the stages of editing, coding then entry and data cleaning. Data analysis was performed using the Chi-square statistical test at a 5% level of significance.

3. Result

3.1. Relationship between Age, Working Period, Number of Pesticide Types with Fatigue

Complaints of fatigue were more likely to be experienced by respondents aged ≥ 32 years (80.6%) than those aged < 32 years, but there was no significant relationship between age and complaints of fatigue ($p=0.270$). Respondents with a working period as a farmer ≥ 5 years were more likely to experience complaints of fatigue than respondents with a working period < 5 years, meaning that working period was associated with complaints of fatigue. The working period was significantly related to complaints of fatigue, where the working period was a protective factor against complaints of fatigue, meaning that the working period reduces the risk of 0.696 for complaints of fatigue. The number of types of pesticides once sprayed was related to complaints of fatigue. Respondents who used ≥ 2 types of pesticides per spraying and experienced complaints of fatigue were greater in proportion than those who used < 2 types. The number of pesticide types ($p=0.000$) was significantly associated with fatigue, where the number of pesticide types was a protective factor against fatigue. The number of < 2 types of pesticides reduced the risk of 0.578 for the occurrence of complaints of fatigue. Pesticide storage method ($p=1.000$) was not significantly associated with complaints of fatigue in respondents, the complete data is presented in table 1 below.

3.2. Relationship between Age, Working Period, Number of Pesticide Types, and Storage with Complaints of Restlessness

Complaints of restlessness tended to be experienced by respondents aged ≥ 32 years (40.3%) compared to those aged < 32 years, there was a significant relationship between age and complaints of restlessness ($p = 0.05$). Respondents with working period as a farmer ≥ 5 years were more likely to experience complaints of anxiety (35.6%) than respondents with working period < 5 years, meaning that working period was associated with complaints

of anxiety, but not significantly related ($p=0.250$). The number of types of pesticides sprayed at one time was associated with restlessness. Respondents who used ≥ 2 types of pesticides at one time and experienced complaints of anxiety were more likely than those who used < 2 types (37.5%). The number of pesticide types ($p=0.037$) was significantly associated with agitation, where the number of pesticide types was a protective factor against agitation. The number < 2 types of pesticides reduced the risk of restlessness by 0.467. Pesticide storage method ($p=0.021$) was not significantly associated with agitation. Poor storage of pesticides was a risk factor for agitation among respondents. The complete data is presented in table 2 below.

3.3. Relationship between Age, Working Period, Number of Pesticide Types with Headache Complaints

The results showed that headache complaints tended to be experienced by respondents aged ≥ 32 years (58.2%) compared to those aged < 32 years, but there was no significant relationship between age and headache complaints ($p=0.593$). Female farmers with working period ≥ 5 years were more likely to experience headache complaints (59.8%) than those with working period < 5 years, but there was no significant relationship between working period and headache complaints ($p=0.206$). Female farmers who used ≥ 2 types of pesticides had a higher proportion of headache complaints (60.4%) compared to those who used < 2 types. The number of pesticide types ($p=0.085$) was not significantly associated with headache complaints. Pesticide storage method was not significantly associated with headache ($p=0.805$). The complete data is presented in table 3 below.

3.4. Relationship between Age, Working Period, Number of Pesticide Types, Storage with Complaints of Blurred Vision

The proportion of respondents with complaints of blurred vision was greater in the age group ≥ 32 years (31.3%) than those aged < 32 years (18.8%). The relationship between age and complaints of blurred vision was not significant ($p=0.137$). Female farmers with working period ≥ 5 years were more likely to experience complaints of blurred vision (29.9%) than respondents with working period < 5 years (16.3%) meaning that there was a relationship between working period with complaints of blurred vision but not significantly related ($p=0.122$). The number of types of pesticides sprayed once was associated with complaints of blurred vision. Respondents who used ≥ 2 types of pesticides and experienced complaints of blurred vision had a higher proportion (26%) than those who used < 2 types (22.5%). The number of pesticide types ($p=0.828$) was not significantly associated with complaints of blurred vision. Pesticide storage method ($p=0.668$) was not significantly associated with complaints of blurred vision. The complete data is presented in table 4 below.

44

3.5 Relationship between Age, Working Period, Number of Pesticide Types with Complaints of Increased Saliva Production

Female farmers with complaints of increased saliva production tend to occur in the age group ≥ 32 years (9%) compared to those aged < 32 years (4.3%), there was no significant relationship between age and complaints of increased saliva production ($p=0.462$). Working period was not significantly associated with complaints of increased saliva production ($p=0.1000$). The number of pesticide types was associated with complaints of increased saliva production. Respondents who used ≥ 2 types of pesticides and experienced complaints of increased saliva production, the proportion was greater (8.3%) than those who used < 2 types (2.5%). The number of pesticide types ($p=0.385$) was not significantly associated with complaints of increased saliva production. Female farmers with poor pesticide storage methods had a higher proportion of complaints of increased saliva production (7.4%), than female farmers with good storage methods (4.8%). Pesticide storage method was not significantly associated with complaints of increased saliva production. The complete data is presented in table 5 below.

3.6. Relationship between Age, Working Period, Number of Pesticide Types, Storage with Nausea Complaints

Nausea tended to be experienced by respondents aged ≥ 32 years (11.9%) compared to those aged < 32 years (7.2%). This shows that there was an association between age and nausea complaints, but there was no significant relationship (0.523). Respondents with working period ≥ 5 years were more likely to experience complaints of nausea (12.6%) than respondents with working period < 5 years (4.1%). Working period was not significantly associated with nausea ($p=0.185$). The number of pesticide types sprayed was associated with nausea. Respondents who used ≥ 2 types of pesticides and experienced nausea were more likely (10.4%) than those who used < 2 types (7.5%). The number of pesticide types ($p=0.836$) was not significantly associated with nausea among female farmers. Pesticide storage method ($p=1.000$) was not significantly associated with nausea among female farmers. The complete data is presented in table 6 below.

3.7. Relationship between Age, Working Period, Number of Pesticide Types, Storage with Complaints of Decreased Appetite

Complaints of reduced appetite among female farmers were more prevalent among those aged ≥ 32 years (16.4%) than those aged < 32 years (7.2%). The data showed an association between age and complaints of reduced appetite, but did not prove a significant relationship ($p=0.163$). Respondents with

working period ≥ 5 years were more likely to experience complaints of reduced appetite (13.8%) than respondents with working period < 5 years (8.2%) meaning that working period was associated with complaints of reduced appetite, but did not show a significant relationship ($p=0.483$). The number of pesticide types was not significantly associated with complaints of decreased appetite ($p=0.295$). Female farmers with poor pesticide storage methods were more likely to experience complaints of decreased appetite (14%) compared to farmers with good pesticide storage methods (4.8%). Pesticide storage method was not significantly associated with complaints of reduced appetite among respondents ($p=0.160$). The complete data is presented in table 7 below.

3.8. Relationship between Age, Working Period, Number of Pesticide Types with Muscle Weakness Complaints

The proportion of complaints of muscle weakness was greater in respondents aged ≥ 32 years (13.4%) than those aged < 32 years. Age was not significantly associated with complaints of muscle weakness ($p=0.742$). Working period of female farmers was not significantly associated with complaints of muscle weakness ($p=0.684$). Number of pesticide types once sprayed was associated with complaints of muscle weakness. The number of pesticide types ($p=0.295$) was not significantly associated with complaints of muscle weakness. Pesticide storage method was not significantly associated with complaints of muscle weakness ($p=0.406$). The complete data is presented in table 8 below.

3.9. Relationship between Age, Working Period, Number of Pesticide Types, Storage with Muscle Spasm Complaints

Female farmers aged ≥ 32 years had a higher proportion of muscle spasm complaints (16.4%) than those aged < 32 years. Age of respondents was not significantly associated with muscle spasm complaints ($p=0.270$). Respondents with a working period as a farmer ≥ 5 years were more likely to experience muscle spasm complaints (14.9%) than respondents with a working period < 5 years (8.2%). Working period was not significantly associated with complaints of spasm in respondents ($p=0.380$). The number of types of pesticides was significantly associated with complaints of muscle spasms ($p=0.393$). Pesticide storage method ($p=0.674$) was not significantly associated with complaints of muscle spasms. The complete data is presented in table 9 below.

4. Discussion

Pesticides enter the human body through the skin, mouth, eyes, and respiratory system, so acute illnesses associated with scientifically confirmed pesticides include headache, abdominal pain, vomiting, skin rashes, difficulty breathing, eye

irritation, sneezing, seizures, and coma (Kalyabina et al., 2021). Organophosphates are one of the active ingredient classes of insecticides that are widely used in the agricultural sector. Symptoms and signs due to Organophosphate exposure that can occur are headache, hypersecretion, muscle twitching, nausea, diarrhea, vomiting, tachycardia/bradycardia, bronchospasm/bronchorrhea respiratory depression, seizures, decreased consciousness (EPA, 2013). Symptoms and signs of Organophosphate poisoning can be grouped into three, namely muscarinic effects, nicotinic effects and central nervous system. Muscarinic effects consist of miosis, bradycardia, hypotension, bronchorrhea, salivation, emesis, diarrhea, abdominal pain, disturbed urinary frequency, and heart rhythm disturbances. Nicotinic effects consist of symptoms of muscle fasciculation, muscle weakness, muscle paralysis, respiratory insufficiency, pale face, sweating. Central nervous system effects include anxiety, agitation, tremor, changes in consciousness, hallucinations, convulsions, respiratory center inhibition, respiratory insufficiency, hypothermia (Voicu et al., 2010).

5 Acetylcholine (AChE) is a nerve conductor present in the entire central nervous system, autonomic nerves and somatic nervous system. Acetylcholine acts on sympathetic and parasympathetic ganglion, parasympathetic receptors, muscle nerve nodes, conducting nerve cells and suprarenal gland medulla. Organophosphates have a mechanism of action of AChE inhibition, where they inactivate AChE by phosphorylating the serine hydroxyl group located on the active group of AChE. With various organophosphate insecticides, an irreversibly inhibited enzyme is formed. Signs and symptoms of intoxication become more prolonged and persistent. Without intervention, toxicity will persist until a sufficient amount of new AChE is formed, 20-30 days, to efficiently destroy excess Ach (Satoh, 2006).

The results showed that female farmers experienced complaints of fatigue, anxiety, headache, blurred vision, increased salivation, nausea, decreased appetite, muscle weakness, and muscle spasms. Research result of (Warno Utomo et al., 2019) mentioned that the complaints experienced by horticultural farmers who use pesticides include itching, sore eyes, body weakness, blurred vision, nausea and dizziness and tightness in the chest. The research (Purba et al., 2023) mentioned several complaints experienced by crop farmers in North Dempo District who were exposed to pesticides were easily tired, restless, headache, blurred vision, increased salivation, frequent nausea, muscle weakness, skin itching, shortness of breath and coughing.

Fatigue is one of the most common complaints among female farmers in connection with exposure to Organophosphate pesticides through agricultural activities in Sekayu District, Musi Banyuasin. Complaints of easy fatigue are experienced by both

age groups both in the younger and older groups with almost the same proportion so that no significant relationship between age and complaints of easy fatigue is proven. In contrast to the variable of age, the variables of working period and number of pesticide types were associated with complaints of fatigue. Working period is the length of time (in years) farmers work as farmers, which is a factor that determines the intensity of pesticide exposure obtained by each farmer. The number of types of pesticides mixed in one spraying certainly determines the toxicity that will be experienced by farmers, because there can be additive effects or synergistic effects. The way pesticides are stored in the home environment is generally poor, this can be the cause of the meaningless relationship between the way pesticides are stored and complaints of fatigue. Poor storage methods will increase the risk of pesticide exposure for female farmers as well as other residents.

Agitation was experienced by 31.6% of the female farmers in this study. The number of types of pesticides is associated with complaints of restlessness in female farmers, where the number of types of pesticides is a protective factor against complaints of restlessness. The use of < 2 types of pesticides can reduce the risk of anxiety by 0.467 times. Complaints of feeling restless are included in the muscarinic effect and are classified as a mild level of poisoning (Jeyaratnam, J. and Koh, 2009). Pesticide storage method is a risk factor for anxiety, where respondents with poor pesticide storage methods have a 2 times greater risk of experiencing anxiety than respondents with good pesticide storage methods. Poor storage methods allow a greater risk of exposure to women farmers, for example, unqualified storage room ventilation causes room temperature to increase so that pesticides evaporate and may be inhaled by women farmers and family members. Careless placement of pesticides can contaminate the home environment.

Headache complaints were experienced by 51% of the respondents in this study, where headache is one of the typical symptoms of pesticide poisoning. Headache complaints were also experienced by farmers spraying horticulture in Meru County, Kenya (Marete et al., 2021). While in Buleleng Bali, headache was the most common complaint experienced by horticultural farmers (51.7%) (Minaka et al., 2016). Headache complaints experienced by female farmers are not related to exposure risk factors including length of service, number of types of pesticides, and how pesticides are stored. However, it is important to limit these exposure risk factors, because long-term exposure can cause complaints related to the effects of pesticides including headaches.

The complaint of blurred vision experienced by respondents in this study was not related to age, working period, number of types of pesticides, and how pesticides were stored. Blurred vision is a muscarinic effect and one of the symptoms of

organophosphate poisoning which is classified as moderate poisoning (Jeyaratnam, J. and Koh, 2009).

The meaninglessness of the relationship between working period and complaints of blurred vision can occur because blood cholinesterase concentrations can recover when resting from pesticide exposure, this causes no symptoms of poisoning in the form of blurred vision. The majority of respondents use more than two types of pesticides in one spraying. This is done with the assumption that it will be more effective in eradicating pests so that they can get increased yields. Effects on the environment and human health have not been a consideration for respondents in the use of pesticides.

Increased saliva production is a muscarinic effect with symptoms of severe intoxication (Jeyaratnam, J. and Koh, 2009). Respondents who experienced increased saliva production in this study were only 6.6%. The small proportion of increased saliva production symptoms resulted in no significant relationship with age, working period, number of types of pesticides, and how to store pesticides. In theory, the number of types of pesticides affects the toxicity they cause. Research on farmers in Brebes proved a significant relationship between the number of types of pesticides with poisoning (Istianah & Yuniastuti, 2017). Research on farmers in Jember also proved a significant relationship between the number of active ingredients of pesticides used with the number of health complaints experienced by farmers (Sofiana et al., 2022).

Nausea is a symptom of moderate muscarinic effects (Jeyaratnam, J. and Koh, 2009). Frequent symptoms of nausea also occurred in respondents of previous related studies (Sofiana et al., 2022). The results of statistical analysis in this study did not prove a significant relationship, age, working period, number of types of pesticides, and how to store pesticides with symptoms of frequent nausea. Complaints of muscle weakness are not significantly related to, working period, and the number of types of pesticides. Muscle weakness is one of the nicotinic effects of moderate intoxication symptoms (Jeyaratnam, J. and Koh, 2009).

The limitation of this study is that there is no examination of female farmers' cholinesterase levels as a biomarker of the effects of pesticides, or metabolite examination to prove the presence of pesticides in the body.

Table 1. Relationship between age, working period, number of pesticide types with fatigue complaints

Variables		Easily Tired				Total		P value	PR (95% CI)
		Yes		No.		n	%		
		n	%	N	%				
Age	< 32 Years	49	71	20	29	69	100	0,270	0,881 (0,728-1,067)
	≥ 32 Years	54	80,6	13	19,4	67	100		
Working period	< 5 Years	29	59,2	20	40,8	49	100	0,002	0,696 (0,543-0,892)
	≥ 5 Years	74	85,1	13	14,9	87	100		
Number of Pesticide Types	< 2 Types	20	50	20	50	40	100	0,000	0,578 (0,420-0,796)
	≥ 2 Types	83	86,5	13	13,5	96	100		
Storage	Bad	71	75,5	23	24,5	94	100	1,000	0,991 (0,808-1,216)
	Good	32	76,2	10	23,8	42	100		

Table 2. Relationship between age, working period, number of pesticide types, and storage with complaints of agitation

Variables		Restlessness				Total		P value	PR (95% CI)
		Yes		No.		n	%		
		n	%	N	%				
Age	< 32 Years	16	23,2	53	76,8	69	100	0,050	0,575 (0,342-0,967)
	≥ 32 Years	27	40,3	40	59,7	67	100		
Working period	< 5 Years	12	24,5	37	75,5	49	100	0,250	0,687 (0,390-1,212)
	≥ 5 Years	31	35,6	56	64,4	87	100		
Number of Pesticide Types	< 2 Types	7	17,5	33	82,5	40	100	0,037	0,467 (0,227-0,959)
	≥ 2 Types	36	37,5	60	62,5	96	100		
Storage	Bad	36	38,3	58	61,7	94	100	0,021	2,298 (1,115-4,736)
	Good	7	16,7	35	83,3	42	100		

Table 3. Relationship between age, working period, number of pesticide types with headache complaints

Variables		Headache				Total		P value	PR (95% CI)
		Yes		No.		n	%		
		N	%	n	%				
Age	< 32 Years	36	52,2	33	47,8	69	100	0,593	0,896 (0,662-1,214)
	≥ 32 Years	39	58,2	28	41,8	67	100		
Working period	< 5 Years	23	46,9	26	53,1	49	100	0,206	0,785 (0,557-1,108)
	≥ 5 Years	52	59,8	35	40,2	87	100		
Number of Pesticide Types	< 2 Types	17	42,5	23	57,5	40	100	0,085	0,703 (0,474-1,044)
	≥ 2 Types	58	60,4	38	39,6	96	100		
Storage	Bad	53	56,4	41	43,6	94	100	0,805	1,076 (0,767-1,510)
	Good	22	52,4	20	47,6	42	100		

Table 4. Relationship between age, working period, number of pesticide types, storage with complaints of blurred vision

Variables		Blurred vision				Total		P value	PR (95% CI)
		Yes		No.		n	%		
		n	%	n	%				
Age	< 32 Years	13	18,8	56	81,2	69	100	0,137	0,601 (0,328-1,100)
	≥ 32 Years	21	31,3	46	68,7	67	100		
Working period	< 5 Years	8	16,3	41	83,7	49	100	0,122	0,546 (0,268-1,112)
	≥ 5 Years	26	29,9	61	70,1	87	100		
Number of Pesticide Types	< 2 Types	9	22,5	31	77,5	40	100	0,828	0,864 (0,444-1,286)
	≥ 2 Types	25	26	71	74	96	100		
Storage	Bad	22	23,4	72	76,6	94	100	0,668	0,819 (0,449-1,496)
	Good	12	28,6	30	71,4	42	100		

Table 5. Relationship between Age, Working period, Number of pesticide types with Complaints of Increased Saliva Production

Variables		Increased Saliva Production				Total		P value	PR (95% CI)
		Yes		No.		n	%		
		n	%	N	%				
Age	< 32 Years	3	4,3	66	95,7	69	100	0,462	0,486 (0,127-1,863)
	≥ 32 Years	6	9	61	91	67	100		
Working period	< 5 Years	3	6,1	46	93,9	49	100	1,000	0,888 (0,232-3,394)
	≥ 5 Years	6	6,9	81	93,1	87	100		
Number of Pesticide Types	< 2 Types	1	2,5	39	97,5	40	100	0,385	0,300 (0,039-2,321)
	≥ 2 Types	8	8,3	88	91,7	96	100		
Storage	Bad	7	7,4	87	92,6	94	100	0,835	1,564 (0,339-7,213)
	Good	2	4,8	40	95,2	42	100		

Table 6. Relationship between age, working period, number of pesticide types, storage with nausea complaints

Variables	Nausea				Total		P value	PR (95% CI)	
	Yes		No.		n	%			
	n	%	n	%					
Age	< 32 Years	5	7,2	64	92,8	69	100	0,523	0,607 (0,209-1,761)
	≥ 32 Years	8	11,9	59	88,1	67	100		
Working period	< 5 Years	2	4,1	47	95,9	49	100	0,185	0,323 (0,075-1,398)
	≥ 5 Years	11	12,6	76	78,7	87	100		
Number of Pesticide Types	< 2 Types	3	7,5	37	92,5	40	100	0,836	0,720 (0,209-2,479)
	≥ 2 Types	10	10,4	86	89,6	96	100		
Storage	Bad	9	9,6	85	90,4	94	100	1,000	1,005 (0,328-3,082)
	Good	4	9,5	38	90,5	42	100		

Table 7. Relationship between age, working period, number of pesticide types, storage and complaints of decreased appetite

Variables	Decreased Appetite				Total		P value	PR (95% CI)	
	Yes		No.		n	%			
	N	%	N	%					
Age	< 32 Years	5	7,2	64	92,8	69	100	0,163	0,441 (0,162-1,202)
	≥ 32 Years	11	16,4	56	83,6	67	100		
Working period	< 5 Years	4	8,2	45	91,8	49	100	0,483	0,592 (0,202-1,736)
	≥ 5 Years	12	13,8	75	86,2	87	100		
Number of Pesticide Types	< 2 Types	7	17,5	33	82,5	40	100	0,295	1,867 (0,747-4,667)
	≥ 2 Types	9	9,4	87	90,6	96	100		
Storage	Bad	14	14,9	80	85,1	94	100	0,160	3,128 (0,744-13,151)
	Good	2	4,8	40	95,2	42	100		

Table 8. Relationship between age, working period, number of pesticide types with complaints of weak muscles

Variables	Muscle Weakness				Total		P value	PR (95% CI)	
	Yes		No		n	%			
	n	%	n	%					
Age	< 32 Years	7	10,1	62	89,9	69	100	0,742	0,755 (0,298-1,912)
	≥ 32 Years	9	13,4	58	85,6	67	100		
Working period	< 5 Years	7	14,3	42	85,7	49	100	0,684	1,381 (0,548-3,478)
	≥ 5 Years	9	10,3	78	89,7	87	100		
Number of Pesticide Types	< 2 Types	7	17,5	33	82,5	40	100	0,295	1,867 (0,747-4,667)
	≥ 2 Types	9	9,4	87	90,6	96	100		
Storage	Bad	13	13,8	81	86,2	94	100	0,406	1,936 (0,582-6,438)
	Good	3	7,1	39	92,9	42	100		

Table 9. Relationship between age, working period, number of pesticide types, storage with muscle spasm complaints

Variables	Muscle Spasms				Total		P value	PR (95% CI)	
	Yes		No		N	%			
	n	%	n	%					
Age	< 32 Years	6	8,7	63	91,3	69	100	0,270	0,530 (0,208-1,351)
	≥ 32 Years	11	16,4	56	83,6	67	100		
Working period	< 5 Years	4	8,2	45	91,9	49	100	0,380	0,546 (0,188-1,584)
	≥ 5 Years	13	14,9	74	85,1	87	100		
Number of Pesticide Types	< 2 Types	7	17,5	33	82,5	40	100	0,393	1,680 (0,688-4,102)
	≥ 2 Types	10	10,4	86	89,6	96	100		
Storage	Bad	13	13,8	81	86,2	94	100	0,674	1,452 (0,503-4,191)
	Good	4	9,5	38	90,5	42	100		

4. Conclusion

Health complaints experienced by female farmers include fatigue, anxiety, headache, blurred vision, nausea, decreased appetite, muscle weakness, and muscle spasms. Age (p=0.05), working period (p=0.002), number of types of pesticides (p=0.000), how to store pesticides (p=0.021) have a significant effect on the onset of subjective health complaints subjective health complaints. Female farmers should

limit pesticide exposure through reducing the length of contact with pesticides, avoid mixing different

types of pesticides, and store pesticides according to storage requirements.

REFERENCES

- EPA. (2013). *Organophosphate Insecticides* (pp. 45–55). EPA. https://www.epa.gov/sites/default/files/documents/rmpp_6thed_ch5_organophosphates.pdf
- FAO. (2021). *FAOSTAT: Pesticide Use*. <http://www.fao.org/faostat/en/#data/RP>
- Filippi, I., Lucero, P., Bonansea, R. I., Lerda, D., Butinof, M., Fernandez, R. A., Wunderlin, D. A., Ame, M. V., & Munoz, S. E. (2021). Validation of exposure indexes to pesticides through the analysis of exposure and effect biomarkers in ground pesticide applicators from Argentina. *Heliyon*, 7(9), e07921. <https://doi.org/10.1016/j.heliyon.2021.e07921>
- Hadian, Z., Eslamizad, S., & Yazdanpanah, H. (2019). Pesticide residues analysis in Iranian fruits and vegetables by gas chromatography-mass spectrometry. *Iranian Journal of Pharmaceutical Research*, 18(1), 275–285.
- Hughes, D., Thongkum, W., Tudpor, K., Turnbull, N., Yukalang, N., Sychareun, V., Van Vo, T., Win, L. L., Watkins, A., & Jordan, S. (2021). Pesticides use and health impacts on farmers in thailand, vietnam, and lao pdr: Protocol for a survey of knowledge, behaviours and blood acetyl cholinesterase concentrations. *PLoS ONE*, 16(9 September), 1–17. <https://doi.org/10.1371/journal.pone.025813>
- Istianah, & Yuniastuti, A. (2017). Hubungan masa kerja, lama menyempromt, jenis pestisida, penggunaan APD dan pengelolaan pestisida dengan kejadian keracunan pada petani di brebes. *Public Health Perspective Journal*, 2(2), 117–123.
- Jeyaratnam, J. and Koh, D. (2009). *Praktik Kedokteran Kerja'e*. EGC.
- Kalyabina, V. P., Esimbekova, E. N., Kopylova, K. V., & Kratasyuk, V. A. (2021). Pesticides: formulants, distribution pathways and effects on human health – a review. *Toxicology Reports*, 8, 1179–1192. <https://doi.org/10.1016/j.toxrep.2021.06.004>
- Kementerian Pemberdayaan Perempuan dan Perlindungan Anak, & B. P. S. (2018). *Profil Perempuan Indonesia*.
- Kumari, D., & John, S. (2019). Health risk assessment of pesticide residues in fruits and vegetables from farms and markets of Western Indian Himalayan region. *Chemosphere*, 224, 162–167. <https://doi.org/10.1016/j.chemosphere.2019.02.091>
- Lekei, E., Ngowi, A. V., Kapeleka, J., & London, L. (2020). Acute pesticide poisoning amongst adolescent girls and women in northern Tanzania. *BMC Public Health*, 20(1), 1–8. <https://doi.org/10.1186/s12889-020-8374-9>
- Marete, G. M., Lalah, J. O., Mputhia, J., & Wekesa, V. W. (2021). Pesticide usage practices as sources of occupational exposure and health impacts on horticultural farmers in Meru County, Kenya. *Heliyon*, 7(2), e06118. <https://doi.org/10.1016/j.heliyon.2021.e06118>
- Minaka, I. A. D. A., Sawitri, A. A. S., & Wirawan, D. N. (2016). Hubungan Penggunaan Pestisida dan Alat Pelindung Diri dengan Keluhan Kesehatan pada Petani Hortikultura di Buleleng, Bali. *Public Health and Preventive Medicine Archive*, 4(1), 74. <https://doi.org/10.15562/phpma.v4i1.60>
- Nambunmee, K., Kawiya, T., Neitzel, R. L., & Seeprasert, P. (2021). Pesticide Spraying and Reduced Cholinesterase Activity among Hill Tribe Farmers in Thailand. *Journal of Health and Pollution*, 11(31), 1–12. <https://doi.org/10.5696/2156-9614-11.31.210908>
- Purba, I. G., Trisnaini, I., & Razak, R. (2023). Keluhan Kesehatan Subjektif Akibat Paparan Pestisida pada Petani Palawijaya di Kecamatan Dempo Utara Pagar Alam. *Jurnal Kesehatan Lingkungan Indonesia*, 22(3), 282–293. <https://doi.org/10.14710/jkli.22.3.282-293>
- Satoh, T. (2006). No Title. In *Global Epidemiology of Organophosphate and Poisonings*. In: GUPTA, R. (ed.) *Toxicology of Organophosphate and Carbamate Compounds*. Burlington: Elseviers.
- Sofiana, K. D., Indreswari, L., Firdaus, J., Prasetyo, A., Pralampita, P. wijang, & Supangat, S. (2022). Analisis Penggunaan Jumlah Bahan Aktif Pestisida dan Banyaknya Keluhan Masalah Kesehatan Pada Petani di Wilayah Agroindustri Jember. *Buletin Poltanesa*, 23(1), 140–146. <https://doi.org/10.51967/tanesa.v23i1.1209>
- UN Woman. (2020). *Generation Equality. Research and Data Section, UN Women: Milan*. unwo.men/eDNK50yyBhA.
- Utami, T. P., Lestari, M., Novrikasari, N., Purba, I. G., Sitorus, R. J., Nandini, R. F., & Fujianti, P. (2021). Penurunan Kadar Enzim Kolinesterase Tenaga Sprayer di Perkebunan Kelapa Sawit. *Jurnal Kesehatan Lingkungan Indonesia*, 20(1), 27–33. <https://doi.org/10.14710/jkli.20.1.27-33>
- Voicu, V. A., Thiermann, H., Rădulescu, F. Ş., Mircioiu, C., & Miron, D. S. (2010). The Toxicokinetics and Toxicodynamics of Organophosphonates versus the Pharmacokinetics and Pharmacodynamics of Oxime Antidotes: Biological Consequences. *Basic and Clinical Pharmacology and Toxicology*, 106(2), 73–85. <https://doi.org/10.1111/j.1742-7843.2009.00486.x>
- Warno Utomo, S., Fauzi, R., & Kusnopranto, H. (2019). Relationship of determination pesticide doses with horticultural farmers health complaints in Cikajang, Garut. *IOP Conference Series: Earth and Environmental Science*, 314(1). <https://doi.org/10.1088/1755-1315/314/1/012033>
- Zhang, Y., Dong, T., Hu, W., Wang, X., Xu, B., Lin, Z., Hofer, T., Stefanoff, P., Chen, Y., Wang, X., & Xia, Y. (2019). Association between exposure to a mixture of phenols, pesticides, and phthalates

and obesity: Comparison of three statistical models. *Environment International*, 123, 325-336. <https://doi.org/10.1016/j.envint.2018.11.076>

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

9%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to kopusat.turnitin@gmail.com Student Paper	3%
2	ejournal.undip.ac.id Internet Source	2%
3	jurnal.stikeskesdam4dip.ac.id Internet Source	2%
4	Submitted to Badan PPSDM Kesehatan Kementerian Kesehatan Student Paper	1%
5	Satoh, Tetsuo. "Global Epidemiology of Organophosphate and Carbamate Poisonings", Toxicology of Organophosphate & Carbamate Compounds, 2006. Publication	1%
6	Fitria Saftarina, Jamsari Jamsari, Masrul Masrul, Yuniar Lestari. "The Risk Factors and Pesticide Poisoning among Horticultural Farmers: A Pilot Study in Indonesia", Open Access Macedonian Journal of Medical Sciences, 2022 Publication	1%

7	Rehab SA Shehata, Noha A Radwan, Huny M Bakry, Aisha Seif Eldin. "Assessment of serum CK-MM level as a bioindicator for work-related musculoskeletal injuries among Emergency hospital workers", Toxicology and Industrial Health, 2022 Publication	1 %
8	ijsrp.org Internet Source	1 %
9	ijhes.com Internet Source	1 %
10	www.slideshare.net Internet Source	1 %
11	Nabila Husna, Tri Niswati Utami. "Identifikasi Postur Janggal, Gerakan Berulang Dan Masa Kerja Terhadap Kejadian MSDs Pekerja Pemanen Kelapa Sawit PT Astra Agro Lestari Tbk", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2023 Publication	<1 %
12	www2.epa.gov Internet Source	<1 %
13	Submitted to University of the Philippines Los Banos Student Paper	<1 %

- | | | |
|----|--|------|
| 14 | Suryapratap Ray, Sanjana Thanjan Shaju. "Bioaccumulation of pesticides in fish resulting toxicities in humans through food chain and forensic aspects", Environmental Analysis Health and Toxicology, 2023
Publication | <1 % |
| 15 | journal.uin-alauddin.ac.id
Internet Source | <1 % |
| 16 | Ilyas Ibrahim, Sahrir Sillehu, Heru Santoso Wahito Nugroho. "Risk Factors for Pesticide Poisoning in Horticultural Farmers in the Agricultural Area of West Seram Regency, Indonesia", National Journal of Community Medicine, 2023
Publication | <1 % |
| 17 | ejournal.unib.ac.id
Internet Source | <1 % |
| 18 | Submitted to iGroup
Student Paper | <1 % |
| 19 | repository.unair.ac.id
Internet Source | <1 % |
| 20 | Dominique Monnet, Antoine P. Brézin, Gary N. Holland, Fei Yu, Alfred Mahr, Lynn K. Gordon, Ralph D. Levinson. "Longitudinal Cohort Study of Patients With Birdshot Chorioretinopathy. I. | <1 % |

Baseline Clinical Characteristics", American Journal of Ophthalmology, 2006

Publication

21

cronfa.swan.ac.uk

Internet Source

<1 %

22

www.ejournal.poltekkesaceh.ac.id

Internet Source

<1 %

23

D. Kumari, S. John. "Health Risk Assessment of Pesticide Residues in Fruits and Vegetables from Farms and Markets of Western Indian Himalayan Region", Chemosphere, 2019

Publication

<1 %

24

Novi Dwi Putri, Trisniarty Adjeng Moelyati, Choriyah Choiriyah. "Influence of Brand Equity, Quality, and Price on Consumer Satisfaction of Ades Brand Bottled Drinking Water in The Sekayu City Sub-District Community Musi Banyuasin District", International Journal of Business, Management and Economics, 2023

Publication

<1 %

25

Nurhikmah, Onny Setiani, Yusniar H Darundiati. "RELATIONSHIP BETWEEN PESTICIDE EXPOSURE AND HEMOGLOBIN LEVEL AND ERYTHROCYTE AMOUNT IN HORTICULTURAL FARMERS IN THE DISTRICT OF PAAL MERAH, JAMBI CITY", International Journal of Research -GRANTHAALAYAH, 2018

<1 %

26

dyuthi.cusat.ac.in

Internet Source

<1 %

27

edepot.wur.nl

Internet Source

<1 %

28

scitepress.org

Internet Source

<1 %

29

www.hindawi.com

Internet Source

<1 %

30

www.nature.com

Internet Source

<1 %

31

"Biotechnological Innovations for Environmental Bioremediation", Springer Science and Business Media LLC, 2022

Publication

<1 %

32

Pauwelijn Verhoestraete, Marijn Carpentier, Jan Donck, Hans Vandekerckhove. "Dilated cardiomyopathy and rhabdomyolysis caused by hypopituitarism: a challenging diagnosis", Acta Cardiologica, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On