

Manuscript JIL Ursada et al.pdf

by sarah yas

Submission date: 06-May-2025 01:35AM (UTC-0500)

Submission ID: 2667073445

File name: Manuscript_JIL_Ursada_et_al.pdf (972.15K)

Word count: 7640

Character count: 41098

Black Soldier Fly Larval Bioconversion of Fruit and Vegetable Waste: Carbon-Nitrogen Conversion and Estimation of Greenhouse Gas Emissions

Riang Ursada¹, Ahmed Zuhdi¹, Arseto Yekti Bagastyo¹, Arry Febrianto¹

¹Department of Environmental Engineering, Faculty of Civil, Planning, and Geo Engineering, Institut Teknologi Sepuluh Nopember, Kampus ITS Sukolilo, Surabaya 60111, Indonesia; e-mail: bagastyo@enviro.its.ac.id

ABSTRAK

Sampah buah dan sayur menyumbang sekitar 36% dari total sampah makanan di Indonesia. Akumulasi sampah ini di tempat pembuangan akhir (TPA) berkontribusi terhadap peningkatan emisi gas rumah kaca (GRK) serta pencemaran lingkungan akibat produksi lindi. Pengolahan menggunakan larva *Black Soldier Fly* (BSF) menawarkan alternatif yang efisien dan berkelanjutan, dengan emisi GRK yang lebih rendah dibandingkan metode komposting maupun digesti anaerobik. Penelitian ini bertujuan untuk mengevaluasi efisiensi biokonversi larva BSF dalam mengurangi sampah buah dan sayur (yakni kol, kangkung, bayam, sawi, semangka, melon, pisang, dan pepaya) melalui analisis transformasi karbon (C) dan nitrogen (N) menjadi biomassa larva dan residu. Komposisi substrat divariasikan dengan rasio sayur terhadap buah sebesar 90:10, 80:20, 70:30, dan 60:40 (b/b), serta diuji pada kepadatan larva 1, 2, dan 4 larva/cm². Hasil menunjukkan bahwa konversi karbon dan nitrogen menjadi biomassa larva masing-masing berkisar antara 7,92–17,59% dan 4,96–21,69%, sedangkan konversi ke residu berkisar antara 22,53–63,75% untuk karbon dan 18,12–80,78% untuk nitrogen. Substrat dengan rasio sayur:buah 90:10 menghasilkan efisiensi konversi tertinggi. Nilai *Approximate Digestibility* (AD), *Efficiency of Conversion of Digested Food* (ECD), dan *Efficiency of Conversion of Ingested Food* (ECI) masing-masing berada pada kisaran 32,44–74,71%, 17,68–42,96%, dan 8,09–18,64%. Tingkat kelangsungan hidup larva mencapai 95,61%. Selain itu, biokonversi dengan larva BSF menghasilkan emisi GRK terendah di antara semua metode pengolahan sampah yang dibandingkan, yaitu sebesar 102,27 g CO₂ ek/kg sampah. Temuan ini menunjukkan potensi signifikan larva BSF dalam pengelolaan sampah buah dan sayur sekaligus meminimalkan dampak lingkungan, serta menjadi dasar bagi optimalisasi lebih lanjut dalam sistem pemanfaatan sampah yang berkelanjutan.

Kata kunci: Karbon, Nitrogen, Larva black soldier fly, Kepadatan larva, Biokonversi

ABSTRACT

Fruit and vegetable waste accounts for approximately 36% of total food waste in Indonesia. The accumulation of this waste in landfills contributes to increasing greenhouse gas (GHG) emissions and environmental pollution through leachate production. Treatment using Black Soldier Fly (BSF) larvae offers an efficient and sustainable alternative, with lower GHG emissions compared to composting or anaerobic digestion. This study aimed to evaluate the bioconversion efficiency of BSF larvae in reducing fruit and vegetable waste (specifically cabbage, water spinach, spinach, mustard greens, watermelon, melon, banana, and papaya) by analyzing the transformation of carbon (C) and nitrogen (N) into larval biomass and residue. Substrate compositions were varied with vegetable-to-fruit ratios of 90:10, 80:20, 70:30, and 60:40 (w/w), and larval densities of 1, 2, and 4 larvae/cm² were tested. Results showed that carbon and nitrogen conversion into larval biomass ranged from 7.92–17.59% and 4.96–21.69%, respectively, while conversion into residue ranged from 22.53–63.75% for carbon and 18.12–80.78% for nitrogen. The substrate with a 90:10 vegetable-to-fruit ratio produced the highest conversion efficiency. The values of Approximate Digestibility (AD), Efficiency of Conversion of Digested Food (ECD), and Efficiency of Conversion of Ingested Food (ECI) ranged from 32.44–74.71%, 17.68–42.96%, and 8.09–18.64%, respectively. The larval survival rates reached 95.61%. Furthermore, BSF bioconversion generated the lowest GHG emissions among all compared waste treatment methods, with a value of 102.27 g CO₂ eq/kg of waste. These findings demonstrate the significant potential of BSF larvae in managing fruit and vegetable waste while minimizing environmental impact, providing a foundation for further optimization in sustainable waste valorization systems.

Keywords: Carbon, Nitrogen, Black soldier fly larvae, Larval density, Bioconversion

Citation: Ursada, R., Zuhdi, A., Bagastyo, A. Y., Febrianto, A. (2025). Black Soldier Fly Larval Bioconversion of Fruit and Vegetable Waste: Carbon-Nitrogen Conversion and Estimation of Greenhouse Gas Emissions. Jurnal Ilmu Lingkungan, xx(x), xx-xx, doi:10.14710/jil.xx.x.xxx-xx

66

1. INTRODUCTION

Indonesia, with a population of approximately 270 million, generated an estimated 31 million tons of municipal solid waste in 2023, of which 40.6% was food waste (Kementerian Lingkungan Hidup dan Kehutanan Sistem Informasi Pengelolaan Sampah Nasional, 2024). Fruit and vegetable waste are the largest contributors to food waste in the country, accounting for 16% and 20%, respectively (Kementerian Perencanaan Pembangunan Nasional, 2021). A significant portion of this organic waste is directly disposed of in landfills, resulting in environmental pollution, greenhouse gas (GHG) emissions, and a shortened landfill lifespan (Akbar, 2016; Lutviyanti et al., 2022; Puger, 2018). Conventional treatment methods such as composting and anaerobic digestion also produce considerable GHG emissions, highlighting the need for more efficient and sustainable waste management alternatives.

An emerging method for managing fruit and vegetable waste is bioconversion using *Hermetia illucens* (Black Soldier Fly, BSF) larvae. These larvae can efficiently consume organic waste and convert it into valuable biomass. BSF larval bioconversion offers several advantages, including lower GHG emissions compared to composting and anaerobic digestion. In fact, total GHG emissions from BSF-based treatment have been reported to be up to 90% lower than those from composting processes (Ermolaev et al., 2019; Mertenat et al., 2019; Pang, Hou, Chen, et al., 2020).

Understanding the factors that influence the efficiency of BSF larval bioconversion is essential to support the development of waste treatment technologies using this method. Bioconversion efficiency can be evaluated based on the larvae's ability to reduce substrate volume and generate larval biomass during the treatment process. Additionally, the conversion of carbon (C) and nitrogen (N) into larval biomass, as well as the losses of C and N throughout bioconversion, serve as key indicators of process performance. The higher the proportion of C and N converted into larval biomass, the more efficient the bioconversion process.

Previous studies have shown that BSF larvae can convert approximately 27–35.5% of carbon and 63.5–75% of nitrogen from food waste with a C/N ratio ranging from 21 to 10 (Jin et al., 2022). A mixture of food waste and straw with a high C/N ratio of 127.3 resulted in carbon and nitrogen conversion into larval biomass ranging from 1.95–13.41% and 5.4–18.93%, respectively (Pang, Hou, Chen, et al., 2020). Meanwhile, bioconversion of a mixture of pig manure and corn stalks with a C/N ratio between 35 and 15 yielded carbon conversion of 4.17–6.61% and nitrogen conversion of 17.45–23.73% (Pang, Hou, Nowar, et al., 2020).

Larval density—defined as the number of larvae per unit surface area—is another critical factor influencing BSF bioconversion efficiency. Larval

63

density plays a significant role in determining the larvae's ability to reduce organic substrate (Purnomo et al., 2021). For optimal larval biomass production, an ideal density is typically within the range of 1.2–5 larvae/cm², with a feeding rate not exceeding 95 mg/larva/day. Excessively high densities may hinder larval movement and increase competition for the substrate, while overly low densities may result in suboptimal substrate consumption and reduced conversion efficiency (Purnomo et al., 2021). Therefore, this study aims to evaluate the carbon and nitrogen conversion in BSF larval bioconversion of fruit and vegetable waste using various substrate compositions and larval densities.

2. METHODS & MATERIALS

2.1. Preliminary research

Preliminary analysis was performed to determine moisture, organic carbon, and nitrogen (organic and inorganic) content of fruit and vegetable waste used as BSF larvae feed. Waste was collected from temporary storage sites near traditional markets or directly from vendors in Surabaya, then sorted, bagged, and transported to the research site. The selected fruit and vegetable waste, categorized as food loss, consisted of relatively fresh, minimally decayed organic material such as leaves, stems, peels, flesh, and seeds. Vegetable waste included cabbage, water spinach (*Ipomoea aquatica*), spinach (*Amaranthus* spp.), and mustard greens (*Brassica juncea*), while fruit waste comprised watermelon, melon, banana, and papaya. Composition varied based on waste availability.

After collecting, the fruit and vegetable waste was shredded until it reached a slurry-like consistency. It was then portioned according to the substrate compositions listed in Table 1. The prepared waste was stored in sealed containers at a temperature of approximately 4°C in the cold storage to maintain freshness prior to use in the bioconversion experiments.

Tabel 1. Research Variable

Substrate composition (%)	Larval density (larvae/cm ²)	Code	Feeding rate (mg/larvae/day)
Vegetable (100)	1	A1	35
	2	A2	
	4	A4	
Vegetable:Fruit (90:10)	1	B1	
	2	B2	
	4	B4	
Vegetable:Fruit (80:20)	1	C1	
	2	C2	
	4	C4	
Vegetable:Fruit (70:30)	1	D1	
	2	D2	
	4	D4	
Vegetable:Fruit (60:40)	1	E1	
	2	E2	
	4	E4	

2.2. Black Soldier Fly Larval Rearing

BSF larval rearing was conducted over two different durations, namely 8 days and 12 days, at the BSF Larvae Facility of the Environmental Engineering Department, Institut Teknologi Sepuluh Nopember. Five-day-old larvae (5 days after oviposition, or 5 DOL) were used as the initial stock and obtained from the Wonorejo Composting Center in Surabaya. The substrates, prepared according to the designated composition variables, were weighed and applied based on a feeding rate of 35 mg/larva/day. Feeding was carried out every two days throughout the rearing period.

The BSF larval rearing reactors were cylindrical plastic containers with a top diameter of 15 cm, bottom diameter of 12 cm, and height of 14 cm. These reactors were placed inside rectangular plastic trays measuring 59 cm in length, 39 cm in width, and 14 cm in height. Each tray held six individual rearing reactors. The reactors were covered with lids and placed on designated racks to prevent larval escape, as illustrated in Figure 1.




Figure 1. BSF Larval Rearing Reactors

At the end of the rearing period, samples of BSF larvae and residual substrate were collected. Separation was performed using a sieve, allowing the larvae to crawl through the mesh and fall into a collection container below, while the residual substrate remained on top of the sieve. The separated BSF larvae and residual substrate were then collected in zip-lock bags. The harvested larvae were rinsed with water to remove dirt and debris, then gently dried using tissue paper. The cleaned samples were stored in a refrigerator or freezer at approximately 4°C, prior to further analysis.

2.3. Analysis

The moisture content was determined using the oven method according to SNI 01-2891-1992. Organic carbon (C) content was determined using the gravimetric method (Adams et al., 1951). Nitrogen content was analyzed by measuring Total Kjeldahl Nitrogen (TKN) according to EPA-NERL: 351.3, which provides values for both organic nitrogen and ammonia (EPA, 1978). Additionally, spectrophotometric analysis was performed to

determine the concentrations of nitrate and nitrite according to EPA-NERL: 351.3 (EPA, 1978).

The relationship between food conversion efficiency and the capacity for carbon (C) and nitrogen (N) conversion during the bioconversion process was analyzed. This analysis allowed for an understanding of how variation in substrate composition and larval density affects the overall efficiency of the BSF bioconversion process in terms of C and N utilization. Waste reduction index (WRI), survival rate (SR), and food conversion efficiencies, including efficiency conversion of ingested food (ECI), approximate digestibility (AD), and efficiency conversion of digested food (ECD), were also calculated. The calculation formulas can be seen in Equations 1 to 7 (Meneguz et al., 2018; Permana & Putra, 2018; Waldbauer, 1968).

$$C \text{ conversion (\%)} = \frac{C \text{ mass adult larvae}}{C \text{ mass total substrate}} \times 100 \quad (1)$$

$$N \text{ conversion (\%)} = \frac{N \text{ mass adult larvae}}{N \text{ mass total substrate}} \times 100 \quad (2)$$

$$AD (\%) = \frac{\text{total substrate (g)} - \text{residue (g)}}{\text{total substrate (g)}} \times 100 \quad (3)$$

$$WRI (\%) = \frac{\frac{\text{total substrate (g)} - \text{residue (g)}}{\text{total substrate (g)}}}{\text{days of trial (d)}} \times 100 \quad (4)$$

$$ECD (\%) = \frac{\text{larvae weight (g)}}{\text{total substrate (g)} - \text{residue (g)}} \times 100 \quad (5)$$

$$ECI (\%) = ECD \times AD \times 100 \quad (6)$$

$$SR (\%) = \frac{\text{final number of live larvae}}{\text{initial number of live larvae}} \times 100 \quad (7)$$

GHG emissions were estimated by calculating the loss of carbon and nitrogen (as nitrate and nitrite) during the BSF larval bioconversion process. These losses were expressed in terms of atomic weight of carbon and nitrogen, then converted to GHG emissions using the ratio of atomic weight to molecular weight of the respective gases. Carbon loss was used to estimate emissions of CO₂ and CH₄, while nitrate and nitrite losses were used to estimate N₂O emissions. The ratio of CO₂ to CH₄ emissions applied was 39416.06:1, based on a study that used fruit and vegetable waste as substrate (Komakech et al., 2023). The estimated GHG emissions from BSF bioconversion were then compared with those from composting, anaerobic digestion, and landfilling, using GHG emission factors that were reported (EPA, 2024; IPCC, 2006; Komakech et al., 2023).

3. RESULTS AND DISCUSSION

3.1. Substrate characteristics

As shown in Table 2, the moisture content of the feed substrate ranged from 87.73% to 89.56%, which is within the optimal range for BSF larval growth. BSF larvae can thrive on substrates with moisture content

ranging from 70% to 90% (Cheng et al., 2017; Salam et al., 2022). Additionally, the carbon-to-nitrogen (C/N) ratio of the feed substrate ranged from 14 to 20, which is considered ideal for both substrate reduction and BSF larval growth (Beesigamukama et al., 2021; Jin et al., 2022; Lu et al., 2021).

Table 2. Substrate Characteristics

Measurand	Substrate composition				
	A	B	C	D	E
Moisture content (%)	87.73	88.34	88.95	89.23	89.56
Ash (%)	11.28	10.59	10.19	9.56	9.06
Carbon (%)	51.58	51.98	52.22	52.58	52.87
Nitrogen (%)	3.45	3.17	3.05	2.85	2.62
C/N ratio (%)	14.93	16.4	17.11	18.44	20.21

The moisture content, carbon content, and C/N ratio of the substrate compositions increased with a higher proportion of fruit in the substrate. This is because fruit mixtures typically have higher moisture content compared to vegetable mixtures, and fruits contain higher levels of sugar, contributing to a greater carbon content than the vegetable mixtures. Conversely, the ash and nitrogen content increased in substrates with a higher proportion of vegetables, as vegetable mixtures tend to have higher nitrogen content (Saputra & Nuryanti, 2018; U.S. Department of Agriculture, 2024).

3.2. Carbon and nitrogen content in BSF larvae

As shown in Figure 2, the carbon content in BSF larvae from the rearing experiments ranged from 46.75% to 50.63%. The lowest carbon content was observed in the larvae reared on A1 over 12 days, while the highest carbon content was found in the larvae reared on E4 over 8 days. The carbon content values in this study are comparable to those found in previous studies, which reported carbon content values of 47.61% and 51.56% for vegetable and fruit substrates, and 47.83% for a mixed vegetable and fruit substrate (Andari et al., 2021; Barrera et al., 2023). The carbon content in BSF larvae increased with both higher larval density and the addition of more fruit in the substrate.

The elevated carbon content may be attributed to the higher fat content in the larvae. Excess carbohydrates in the diet are converted into fats through the larvae's metabolic processes (Carpentier et al., 2024). Increased larval density has been shown to elevate the fat content in BSF larvae (Barragan-Fonseca et al., 2018). Additionally, the high fructose content in fruits, a simple sugar, also contributed to the higher carbon content in the larvae. The presence of sugars in the larvae feed facilitates the larvae's ability to obtain carbohydrates as a carbon source (Agustin et al., 2023; Prahasuti, 2011).

As shown in Figure 3, the nitrogen content in BSF larvae ranged from 1.26% to 4.29%. The lowest nitrogen content was observed in larvae reared on B1 over 12 days, while the highest nitrogen content was found in larvae reared on A2 over 8 days. The nitrogen

content in the BSF larvae from this study was lower than values reported in previous research, which showed nitrogen content ranging from 5% to 7% when reared on pig manure substrates, 7% on a wheat and starch powder substrate, and 6.1% to 7.1% on food waste substrates (Jin et al., 2022; Parodi et al., 2020, 2022).




Figure 2. Carbon Content of BSF Larvae




Figure 3. Nitrogen Content of BSF Larvae

The lower nitrogen content in the BSF larvae from this study may be due to the low protein content in the substrates, as well as the high fiber content. Low protein levels in the larval feed during rearing can result in reduced nitrogen content in the larvae (Aldis et al., 2024). Furthermore, fiber is indigestible to BSF larvae, and high fiber content in the feed can reduce the overall nutrient density available for larval growth, resulting in lower nitrogen content in the larvae (Froonickx et al., 2023; Gold et al., 2018).

3.3. Carbon and nitrogen content in residues

As shown in Figure 4, the carbon content in the residues ranged from 42.76% to 47.89% for the 8-day and 12-day rearing periods. The lowest carbon content in the residues was observed in A1 after 12 days of rearing, while the highest carbon content was found in B2 after 12 days of rearing. A higher carbon weight in the residue leads to a higher conversion of carbon into residues, while a lower carbon weight results in a lower conversion of carbon into residue. There was a decrease in carbon content in the substrate after it was consumed by the BSF larvae, indicating that some of the carbon was converted into biomass by the larvae. Additionally, the carbon content in the residues tended to be lower at a larval density of 1 larva/cm², suggesting that more carbon

was converted into BSF biomass and/or lost at lower larval densities. The carbon content in the residues of this study was comparable to values obtained in previous studies, which reported carbon content in residues of 41.40% to 50.14% for mixed vegetable and fruit waste substrates (Oemar et al., 2023).

As shown in Figure 5, the nitrogen content in the residues ranged from 1.67% to 4.48%. The lowest nitrogen content in the residues was observed in C2 after 12 days of rearing, while the highest nitrogen content in the residues was found in A1 after 8 days of rearing. The nitrogen content in the residues from this study was similar to previous research on mixed vegetable and fruit waste residues, which ranged from 1.303% to 3.701% (Oemar et al., 2023).




Figure 4. Carbon Content of Residue




Figure 5. Nitrogen Content of Residue

An increase in nitrogen content in the residues, compared to the initial nitrogen content of the substrate, may be due to the decomposition of organic matter and nitrogen mineralization by microorganisms, which produces nitrogen. Additionally, the biochemical activity in the BSF larvae's digestive system contributes to the nitrogen present in the residues (L. Purnamasari et al., 2021; Sarpong et al., 2019).

3.4. Carbon and nitrogen conversion

The percentage of carbon converted into BSF larva biomass during the 8-day and 12-day rearing periods ranged from 10.52% to 17.59% and 7.92% to 13.76%, respectively. In contrast to previous studies using food waste, which reported carbon conversion rates ranging from 27% to 35.5% (Jin et al., 2022). Bioconversion of food waste and straw mixtures

yields carbon conversion from 1.95% to 13.41% (Pang, Hou, Chen, et al., 2020). In the bioconversion of pig manure and corn stalk mixtures, the carbon conversion rate to BSF larva biomass ranges from 4.17% to 6.61% (Pang, Hou, Nowar, et al., 2020). The carbon conversion into BSF larva biomass was higher during the 8-day rearing period compared to the 12-day period because the larvae in the 12-day rearing had entered the pre-pupal phase, during which they consume little or no food and mostly move around seeking shaded, dry, and safe areas to pupate (Dortmans et al., 2021; Lievens et al., 2021; Putra & Ariesmayana, 2020; Wardhana, 2017).

As shown in Figures 6 and 7, the rearing of BSF larvae at a density of 1 larva/cm² was more efficient in converting carbon into larval biomass, but the carbon converted into residue was low. This could be attributed to the low density, which allows the larvae to move more freely, causing more carbon as an energy source to be wasted (Parodi et al., 2022; Parra Paz, Carrejo, & Rodríguez, 2015).

According to Figures 8 and 9, the percentage of nitrogen converted into BSF larva biomass during the 8-day and 12-day rearing periods ranged from 9.21% to 19.95% and 4.96% to 21.69%, respectively. The percentage of nitrogen converted into residue during the 8-day and 12-day rearing periods ranged from 30.68% to 80.78% and 18.12% to 62.82%, respectively. The nitrogen conversion into BSF larva biomass in this study differed from studies using vegetable and fruit waste substrates (Lalander et al., 2019) and food waste substrates (Jin et al., 2022), where nitrogen conversion into larval biomass was reported to be 34.3% and 35% to 75%, respectively. However, the nitrogen conversion values in this study were similar to those observed in studies using food waste and straw mixtures (Pang, Hou, Chen, et al., 2020) and pig manure and corn stalk mixtures (Purnomo et al., 2021), where nitrogen conversion into larva biomass ranged from 5.4% to 18.93% and 17.45% to 23.73%, respectively. The high fiber content in the substrate may have caused the lower nitrogen conversion in the BSF larvae because the larvae are unable to digest fiber, and the high fiber content in the substrate could reduce the overall nutritional density for larval growth (Froonickx et al., 2023; Gold et al., 2018).




Figure 6. Conversion of Substrate Carbon to BSF Larvae Carbon




Figure 7. Conversion of Substrate Carbon to BSF Residue Carbon




Figure 8. Conversion of Substrate Nitrogen to BSF Larvae Nitrogen




Figure 9. Conversion of Substrate Nitrogen to Residue Nitrogen

3.5. Larval moisture content

No significant differences in moisture content were observed across the variations in larval density and substrate composition. The moisture content of BSF larvae reared for 8 days ranged from 67.68% to 70.97%. The lowest moisture content was found in variation D2, while the highest was observed in variation B1. For larvae reared over 12 days, moisture content ranged from 62.27% to 65.66%, with the lowest value in variation C2, and the highest in variation A1.

The BSF larvae moisture content in this study, ranging from 67.68–70.97% (8-day rearing) and 62.27–65.66% (12-day rearing), is similar to the range reported by previous studies using fruit and vegetable substrates, such as 63.37–67.41% (Saragi, 2015). However, these values are relatively lower compared to the findings of other studies, which

reported moisture contents of 78.90% using vegetable substrates (Amri et al., 2023) and 78.90–81.95% using fruit and vegetable substrates (D. K. Purnamasari et al., 2023). Lower moisture content in BSF larvae may indicate a higher fat content, as suggested by (Amri et al., 2023). The decline in moisture content from day 8 to day 12 is associated with the physiological transition of the larvae into the prepupal stage, during which water content typically decreases (D. K. Purnamasari et al., 2023).

3.6. Bioconversion performance

Substrate reduction was calculated based on the dry weight of the initial feed and resulting residue to assess the consumption efficiency of BSF larvae (Meneguz et al., 2018). After 8 days of rearing, substrate reduction ranged from 32.44% to 74.71%, with the highest reduction observed in B1 and the lowest in E4. For the 12-day rearing period, reduction ranged from 38.44% to 74.03%, with the highest in A1 (68.25%) and B1 (66.55%). Overall, lower larval density resulted in higher substrate reduction due to improved feed accessibility (Parra Paz, Carrejo, & Gómez Rodríguez, 2015). Substrates with higher vegetable content also showed greater reduction compared to fruit-dominant ones (Amin et al., 2024; Triwandani et al., 2023). Additionally, substrate reduction decreased in the 12-day rearing due to reduced feed intake by larvae entering the prepupal stage (Dortmans et al., 2021; Lievens et al., 2021; Putra & Ariesmayana, 2020). The waste reduction index ranged from 4 to 9.4% dry matter.

Approximate Digestibility (AD) is defined as the proportion of feed that is digested relative to the total amount provided (Permana & Putra, 2018; Waldbauer, 1968). The AD values observed in the 12-day rearing period were consistently lower than those recorded at 8 days as can be seen on Figure 10. This decline is likely due to a reduction in larval feeding activity, as many individuals had entered the prepupal stage by day 12, where feeding is known to diminish (Dortmans et al., 2021; Lievens et al., 2021; Putra & Ariesmayana, 2020). The highest AD was recorded in substrate composition B1. Lower larval densities generally resulted in higher AD values, likely due to reduced competition and better feed accessibility (Parra Paz, Carrejo, & Gómez Rodríguez, 2015). Conversely, higher densities and low survival rates may have led to uneaten feed, reducing overall digestibility. Compared to previous findings, the AD values in this study were lower than the BSF larvae fed kitchen waste values 87.7% (Mahmood et al., 2021).

Efficiency of Conversion of Digested Food (ECD) represents the larvae's ability to transform digested feed into body mass. Higher ECD values indicate more efficient bioconversion processes (Permana & Putra, 2018; Waldbauer, 1968). Figure 11 shown ECD values, the highest ECD during the 8-day rearing was achieved with substrate composition E2. In the 12-day rearing, the highest ECD (26.97%) was recorded in substrate A2. These findings align with ECD values

between 11–56% using fruit and vegetable substrates (Rofi et al., 2021). Notably, ECD values in this study were generally higher than BSF reared on kitchen waste and coffee waste values 16.60% and 2.71–5.00% respectively (Mahmood et al., 2021; Permana & Putra, 2018), suggesting favorable conversion efficiency under the tested conditions.




Figure 10. Approximate Digestibility




Figure 11. Efficiency of Conversion of Digested Food




Figure 12. Efficiency of Conversion of Ingested Food
Efficiency of Conversion of Ingested Food (ECI) measures the proportion of ingested feed that is utilized for energy production and biomass formation. It is influenced by feed digestibility and the larvae's metabolic efficiency (Permana & Putra, 2018; Waldbauer, 1968). ECI values can be seen on Figure 12. The highest value was observed in substrate B1,

while the lowest was found in substrate D4. At this stage, ECI values for larval densities of 1, 2, and 4 larvae/cm² ranged from 12.28–18.64%, 13.58–17.12%, and 10.64–16.73%, respectively. For the 12-day rearing, highest ECI values is observed in substrate C1. The trend showed that lower larval densities generally supported higher ECI values. High ECI reflects efficient conversion of ingested feed into larval biomass and energy, highlighting the importance of optimizing both substrate composition and rearing conditions.

Collectively, these results emphasize the significance of larval density and substrate composition in influencing BSF larvae's feeding performance. Optimal combinations of vegetable-rich substrates and lower larval densities consistently supported improved digestibility and conversion efficiencies, offering valuable insights for enhancing the sustainability and productivity of BSF bioconversion system. As a summary, the characteristics of the larvae, prepupae, and residue for the best reduction (B1) are presented in Table 3.

Table 3. Larvae, Prepupae, and Residue Characteristics

Measurand	Larvae	Prepupae	Residue
Moisture content (%)	70.79	64.92	92.66
Ash (%)	15.59	15.49	22.67
Carbon (%)	49.08	49.14	44.96
Nitrat (%)	0.0015	0.014	0.007
Nitrit (%)	0.007	0.004	0.004
N organic (%)	2.432	1.242	3.438

3.7. Estimates of GHG emissions

The carbon and nitrogen losses during the BSF larval bioconversion process are primarily released in gaseous forms. Carbon is emitted as carbon dioxide (CO₂) and methane (CH₄), generated through larval metabolism, respiration, and substrate decomposition processes. Meanwhile, nitrogen is lost as ammonia (NH₃) and nitrous oxide (N₂O). NH₃ is released through volatilization of organic nitrogen, while N₂O is formed via nitrification and denitrification of nitrite and nitrate compounds (Froonickx et al., 2023).

The estimated GHG emissions include CO₂, CH₄, and N₂O, while NH₃ was not considered due to its exclusion as a greenhouse gas. According to the emission estimates presented in Table 4, BSF bioconversion generated the lowest GHG emissions among all compared waste treatment methods, with a value of 102.27 g CO₂ eq/kg of waste. These findings suggest that BSF larval bioconversion is a more environmentally friendly approach for processing fruit and vegetable waste when compared to composting, anaerobic digestion, and landfilling.

Table 4. Estimates of GHG emissions

Processing Method (Emission Factor Source)	Total Emission (g CO ₂ eq)				
	CO ₂	CH ₄	N ₂ O	Total	Emisi (g CO ₂ eq/kg waste)
Bioconversion of BSF Larvae	2429.41	0.56	120.65	2550.62	102.27
Composting (Komakech et al., 2023)	2181.31	342.53	454.99	2978.83	119.42
Composting (IPCC, 2006)	NA	2494.36	1783.96	NA	NA
Composting (EPA, 2024)	NA	NA	NA	2993.23	120.00

Processing Method (Emission Factor Source)	Total Emission (g CO ₂ eq)				
	CO ₂	CH ₄	N ₂ O	Total	Emisi (g CO ₂ eq/kg waste)
Anaerobic (Komakech et al., 2023)	1526.80	1055.20	454.99	3036.98	121.75
Anaerobic (IPCC, 2006)	NA	498.87	NA	NA	NA
Anaerobic (EPA, 2024)	NA	NA	NA	3741.53	150.00
Landfilling (IPCC, 2006)	NA	8190.84	NA	NA	NA
Landfilling (EPA, 2024)	NA	NA	NA	7732.50	310.00

NA: not available

4. CONCLUSION

This study demonstrates that Black Soldier Fly (BSF) larval bioconversion offers an effective and environmentally friendly approach for managing fruit and vegetable waste. Carbon conversion into BSF larval biomass ranged from 7.92% to 17.59%, while carbon conversion into residue ranged from 22.53% to 63.75%. Lower larval densities favored higher carbon conversion into larval biomass, though this was accompanied by reduced carbon retention in residue. Nitrogen conversion into larval biomass ranged from 4.96% to 21.69%, and into residue from 18.12% to 80.78%, with substrate composition and larval density showing no significant effect on nitrogen partitioning. Among the compared waste treatment methods, BSF bioconversion resulted in the lowest greenhouse gas emissions (102.27 g CO₂ eq/kg waste), highlighting its potential as a sustainable alternative to composting, anaerobic digestion, and landfilling.

ACKNOWLEDGEMENT

This research was funded by the Indonesian Endowment Fund for Education (LPDP) through the Ministry of Education, Culture, Research, and Technology of Indonesia, and administered under the Indonesia-NTU Singapore Institute of Research for Sustainability and Innovation (INSPIRASI) program (Grant No. 6636/E3/KL.02.02/2023).

REFERENCES

- Adams, R. C., Bennett, F. M., Dixon, J. K., Lough, R. C., Maclean, F. S., & Martin, G. I. (1951). The utilization of organic wastes in N.Z. *New Zealand Engineering*, 6(11), 396–424. <https://search.informit.org/doi/10.3316/informit.359233295091848>
- Agustin, H., Warid, W., & Musadik, I. M. (2023). Kandungan Nutrisi Kasgot Larva Lalat Tentara Hitam (*Hermetia illucens*) Sebagai Pupuk Organik. *Jurnal Ilmu-Ilmu Pertanian Indonesia*, 25(1), 12–18. <https://doi.org/10.31186/jipi.25.1.12-18>
- Akbar, R. A. (2016). Pengaruh Paparan CH₄ dan H₂S Terhadap Keluhan Gangguan Pernapasan Pemulung di TPA Mrican Kabupaten Ponorogo. *Journal of Industrial Hygiene and Occupational Health*, 1(1), 1–14. <https://doi.org/10.21111/jihoh.v1i1.603>
- Aldis, R. E., Muhlisin, M., Zuprizal, Z., Sasongko, H., Hanim, C., & Anas, M. Al. (2024). Black soldier fly larvae meal supplementation in a low protein diet reduced performance, but improved nitrogen efficiency and intestinal morphology of duck. *Animal Bioscience*, 37(4), 678–688. <https://doi.org/10.5713/ab.23.0259>
- Amin, U. K., Lando, A. T., & Djamaluddin, I. (2024). Potential of Black Soldier Fly Larvae in Reduction Various Types Organic Waste. *Ecological Engineering & Environmental Technology*, 25(9), 190–201. <https://doi.org/10.12912/27197050/190639>
- Amri, N. N., Ardiasani, S., Elvira, N., Putri, L. S. E., & Sahari, B. (2023). Mortality, Development and Nutrient Content of Black Soldier Fly Larvae (*Hermetia illucens* L.) Maintained on Different Types of Household Organic Waste. *IOP Conference Series: Earth and Environmental Science*, 1220(1), 012018. <https://doi.org/10.1088/1755-1315/1220/1/012018>
- Andari, G., Ginting, N. M., & Nurdiana, R. (2021). Black Soldier Fly Larvae (*Hermetia illucens*) as a Waste Reduction Agent and an Alternative Livestock Feed. *Jurnal Ilmiah Peternakan Terpadu*, 9(3), 246. <https://doi.org/10.23960/jiptv9i3.p246-252>
- Barragan-Fonseca, K. B., Dicke, M., & van Loon, J. J. A. (2018). Influence of larval density and dietary nutrient concentration on performance, body protein, and fat contents of black soldier fly larvae (*Hermetia illucens*). *Entomologia Experimentalis et Applicata*, 166(9), 761–770. <https://doi.org/10.1111/eea.12716>
- Barrera, J., Salinas, E. R., Rivera, A. R., Malalis, J. S., Bagaforo, Engr. R. O., Araya, J. P., & Sulay, Dr. D. L. D. (2023). Productivity and Nutritive Value of Black Soldier Fly (*Hermetia illucens*) Larvae Fed with Different Waste Substrates. *International Journal of Multidisciplinary: Applied Business and Education Research*, 4(2), 588–599. <https://doi.org/10.11594/ijmaber.04.02.25>

- Beesigamukama, D., Mochoge, B., Korir, N. K., K.M. Fiaboe, K., Nakimbugwe, D., Khamis, F. M., Subramanian, S., Wangu, M. M., Dubois, T., Ekesi, S., & Tanga, C. M. (2021). Low-cost technology for recycling agro-industrial waste into nutrient-rich organic fertilizer using black soldier fly. *Waste Management*, 119, 183–194. <https://doi.org/10.1016/j.wasman.2020.09.043>
- Carpentier, J., Martin, C., Luttenschlager, H., Deville, N., Ferrara, D., Purcaro, G., Blecker, C., Francis, F., & Caparros Megido, R. (2024). Common soluble carbohydrates affect the growth, survival, and fatty acid profile of black soldier fly larvae *Hermetia illucens* (Stratiomyidae). *Scientific Reports*, 14(1), 28157. <https://doi.org/10.1038/s41598-024-75730-5>
- Cheng, J. Y. K., Chiu, S. L. H., & Lo, I. M. C. (2017). Effects of moisture content of food waste on residue separation, larval growth and larval survival in black soldier fly bioconversion. *Waste Management*, 67, 315–323. <https://doi.org/10.1016/j.wasman.2017.05.046>
- Dortmans, B. M. A., Egger, J., Diener, S., & Zurbrugg, C. (2021). Black Soldier Fly Biowaste Processing-A Step-by-Step Guide, 2 nd Edition. In *Bibliographic*.
- EPA. (1978). *EPA-NERL: 351.3 Nitrogen, Kjeldahl, Total (Colorimetric; Titrimetric; Potentiometric)*. U.S. Environmental Protection Agency.
- EPA. (2024). *Emission Factors for Greenhouse Gas Inventories*. U.S. Environmental Protection Agency.
- Ermolaev, E., Lalander, C., & Vinnerås, B. (2019). Greenhouse gas emissions from small-scale fly larvae composting with *Hermetia illucens*. *Waste Management*, 96, 65–74. <https://doi.org/10.1016/j.wasman.2019.07.011>
- Froonickx, L., Berrens, S., Broeckx, L., & Van Miert, S. (2023). The potential of black soldier fly to recycle nitrogen from biowaste. *Current Opinion in Green and Sustainable Chemistry*, 44, 100864. <https://doi.org/10.1016/j.cogsc.2023.100864>
- Gold, M., Tomberlin, J. K., Diener, S., Zurbrugg, C., & Mathys, A. (2018). Decomposition of biowaste macronutrients, microbes, and chemicals in black soldier fly larval treatment: A review. *Waste Management*, 82, 302–318. <https://doi.org/10.1016/j.wasman.2018.10.022>
- IPCC. (2006). *2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme* (S. Eggleston, L. Buendia, K. Miwa, T. Ngara, & K. Tanabe, Eds.; Vol. 1). IGES, Japan.
- Jin, N., Liu, Y., Zhang, S., Sun, S., Wu, M., Dong, X., Tong, H., Xu, J., Zhou, H., Guan, S., & Xu, W. (2022). C/N-Dependent Element Bioconversion Efficiency and Antimicrobial Protein Expression in Food Waste Treatment by Black Soldier Fly Larvae. *International Journal of Molecular Sciences*, 23(9), 5036. <https://doi.org/10.3390/ijms23095036>
- Kementerian Lingkungan Hidup dan Kehutanan Sistem Informasi Pengelolaan Sampah Nasional. (2024, July 20). *Capaian Kinerja Pengelolaan Sampah*. <https://sipsn.menlhk.go.id/sipsn/>
- Kementerian Perencanaan Pembangunan Nasional. (2021). *Study Report Food Loss and Waste in Indonesia: Supporting the Implementation of Circular Economy and Low Carbon Development*.
- Komakech, A. J., Rubagumya, I., Kizito, S. S., Zziwa, A., Kabenge, I., & Menya, E. (2023). *Quantifying greenhouse gas emissions from three fruit and vegetable waste management technologies using the static chamber*. <https://doi.org/10.21203/rs.3.rs-3312640/v1>
- Lalander, C., Diener, S., Zurbrugg, C., & Vinnerås, B. (2019). Effects of feedstock on larval development and process efficiency in waste treatment with black soldier fly (*Hermetia illucens*). *Journal of Cleaner Production*, 208, 211–219. <https://doi.org/10.1016/j.jclepro.2018.10.017>
- Lievens, S., Poma, G., De Smet, J., Van Campenhout, L., Covaci, A., & Van Der Borgh, M. (2021). Chemical Safety of Black Soldier Fly Larvae (*Hermetia illucens*), Knowledge Gaps and Recommendations for Future Research: a Critical Review. *Journal of Insects as Food and Feed*, 7(4), 383–396. <https://doi.org/10.3920/JIFF2020.0081>

- Lu, Y., Zhang, S., Sun, S., Wu, M., Bao, Y., Tong, H., Ren, M., Jin, N., Xu, J., Zhou, H., & Xu, W. (2021). Effects of Different Nitrogen Sources and Ratios to Carbon on Larval Development and Bioconversion Efficiency in Food Waste Treatment by Black Soldier Fly Larvae (*Hermetia illucens*). *Insects*, 12(6), 507. <https://doi.org/10.3390/insects12060507>
- Lutviyani, A., Firdausi, F. F., & Hanim, H. (2022). Tinjauan Limbah Makanan Terhadap Lingkungan dalam Perspektif Islam dan Sains. *Prosiding Konferensi Integrasi Interkoneksi Islam Dan Sains* 4, 49–53.
- Mahmood, S., Zurbrugg, C., Tabinda, A. B., Ali, A., & Ashraf, A. (2021). Sustainable Waste Management at Household Level with Black Soldier Fly Larvae (*Hermetia illucens*). *Sustainability*, 13(17), 9722. <https://doi.org/10.3390/su13179722>
- Meneguz, M., Schiavone, A., Gai, F., Dama, A., Lussiana, C., Renna, M., & Gasco, L. (2018). Effect of rearing substrate on growth performance, waste reduction efficiency and chemical composition of black soldier fly (*Hermetia illucens*) larvae. *Journal of the Science of Food and Agriculture*, 98(15), 5776–5784. <https://doi.org/10.1002/jsfa.9127>
- Mertenat, A., Diener, S., & Zurbrugg, C. (2019). Black Soldier Fly biowaste treatment – Assessment of global warming potential. *Waste Management*, 84, 173–181. <https://doi.org/10.1016/j.wasman.2018.11.040>
- Oemar, T., Purwaningrum, P., Ruhayat, R., & Ashardiono, F. (2023). Potential of Black Soldier Fly (BSF) in Reducing Municipal Food Loss and Waste (FLW) at Taman Sari District, West Jakarta. *Indonesian Journal of Urban and Environmental Technology*, 132–144. <https://doi.org/10.25105/urbanenvirotech.v6i2.16932>
- Pang, W., Hou, D., Chen, J., Nowar, E. E., Li, Z., Hu, R., Tomberlin, J. K., Yu, Z., Li, Q., & Wang, S. (2020). Reducing greenhouse gas emissions and enhancing carbon and nitrogen conversion in food wastes by the black soldier fly. *Journal of Environmental Management*, 260, 110066. <https://doi.org/10.1016/j.jenvman.2020.110066>
- Pang, W., Hou, D., Nowar, E. E., Chen, H., Zhang, J., Zhang, G., Li, Q., & Wang, S. (2020). The influence on carbon, nitrogen recycling, and greenhouse gas emissions under different C/N ratios by black soldier fly. *Environmental Science and Pollution Research*, 27(34), 42767–42777. <https://doi.org/10.1007/s11356-020-09909-4>
- Parodi, A., Yao, Q., Gerrits, W. J. J., Mishyna, M., Lakemond, C. M. M., Oonincx, D. G. A. B., & Van Loon, J. J. A. (2022). Upgrading ammonia-nitrogen from manure into body proteins in black soldier fly larvae. *Resources, Conservation and Recycling*, 182, 106343. <https://doi.org/10.1016/j.resconrec.2022.106343>
- Parra Paz, A. S., Carrejo, N. S., & Gómez Rodríguez, C. H. (2015). Effects of Larval Density and Feeding Rates on the Bioconversion of Vegetable Waste Using Black Soldier Fly Larvae *Hermetia illucens* (L.), (Diptera: Stratiomyidae). *Waste and Biomass Valorization*, 6(6), 1059–1065. <https://doi.org/10.1007/s12649-015-9418-8>
- Parra Paz, A. S., Carrejo, N. S., & Rodríguez, C. H. (2015). Effects of Larval Density and Feeding Rates on the Bioconversion of Vegetable Waste Using Black Soldier Fly Larvae *Hermetia illucens* (L.), (Diptera: Stratiomyidae). *Waste and Biomass Valorization*, 6(6), 1059–1065. <https://doi.org/10.1007/s12649-015-9418-8>
- Permana, A. D., & Putra, J. E. N. R. E. (2018). Growth of Black Soldier Fly (*Hermetia illucens*) Larvae Fed on Spent Coffee Ground. *IOP Conference Series: Earth and Environmental Science*, 187(1), 012070. <https://doi.org/10.1088/1755-1315/187/1/012070>
- Prahastuti, S. (2011). Konsumsi Fruktosa Berlebihan Dapat Berdampak Buruk Bagi Kesehatan Manusia. *Maranatha Journal of Medicine and Health*, 10(2).
- Puger, I. G. N. (2018). Sampah Organik, Kompos, Pemanasan Global, dan Penanaman Aglaonema di Pekarangan. *Agro Bali (Agricultural Journal)*, 1, 127–136.
- Purnamasari, D. K., Erwan, Syamsuhaidi, Sumiati, Wiryawan, I. K. G., Maslami, V., & Kurniyati.

- (2023). Kandungan Nutrisi Setiap Fase Siklus Black Soldier Fly (BSF) yang Dibudidayakan Menggunakan Sampah Organik. *Jurnal Ilmu Dan Teknologi Peternakan Indonesia (JITPI) Indonesian Journal of Animal Science and Technology*, 9(2), 111–121. <https://doi.org/10.29303/jitpi.v9i2.182>
- Purnamasari, L., Muhlison, W., & Sucipto, I. (2021). Biokonversi limbah ampas tahu dan limbah sayur dengan menggunakan agen larva Black Soldier Fly (*Hermetia illucens*). 105–111. <https://doi.org/10.25047/animpro.2021.13>
- Purnomo, B. C., Nurjuzuli, N., & Suhartono, S. (2021). Pengaruh Luas Penampang Wadah Terhadap Besarnya Reduksi Volume Sampah Organik Rumah Tangga Menggunakan Larva Lalat Bsf (Black Soldier Fly). *Jurnal Sehat Mandiri*, 16(2), 99–108. <https://doi.org/10.33761/jsm.v16i2.474>
- Putra, Y., & Ariesmayana, A. (2020). Efektifitas Penguraian Sampah Organik Menggunakan Maggot (BSF) di Pasar Rau Trade Center. *Jurnal Lingkungan Dan Sumberdaya Alam (JURNALIS)*, 3(1). <https://ejournal.lppm-unbaja.ac.id/index.php/jls/article/view/888>
- Rofi, D. Y., Auvaria, S. W., Nengse, S., Oktorina, S., & Yusrianti, Y. (2021). Modifikasi Pakan Larva Black Soldier Fly (*Hermetia illucens*) sebagai Upaya Percepatan Reduksi Sampah Buah dan Sayuran. *Jurnal Teknologi Lingkungan*, 22(1), 130–137. <https://doi.org/10.29122/jtl.v22i1.4297>
- Salam, M., Shahzadi, A., Zheng, H., Alam, F., Nabi, G., Dezhi, S., Ullah, W., Ammara, S., Ali, N., & Bilal, M. (2022). Effect of different environmental conditions on the growth and development of Black Soldier Fly Larvae and its utilization in solid waste management and pollution mitigation. *Environmental Technology & Innovation*, 28, 102649. <https://doi.org/10.1016/j.eti.2022.102649>
- Saputra, R. F., & Nuryanti. (2018). Studi Kelayakan Kadar Air, Abu, Protein dan Tembaga (Cu) Pada Sayuran di Pasar Sunter Jakarta Utara sebagai Bahan Suplemen Makanan. *Indonesia Natural Research Pharmaceutical Journal*, 3(2), 150–161.
- Saragi, E. S. (2015). *Penentuan Optimal Feeding Rate Larva Black Soldier Fly (Hermetia illucens) dalam Mereduksi Sampah Organik Pasar*. Institut Teknologi Sepuluh Nopember.
- Sarpong, D., Odoro-Kwarteng, S., Gyasi, S. F., Buamah, R., Donkor, E., Awuah, E., & Baah, M. K. (2019). Biodegradation by composting of municipal organic solid waste into organic fertilizer using the black soldier fly (*Hermetia illucens*) (Diptera: Stratiomyidae) larvae. *International Journal of Recycling of Organic Waste in Agriculture*, 8(S1), 45–54. <https://doi.org/10.1007/s40093-019-0268-4>
- Triwandani, A., Purwaningrum, P., & Ruhayat, R. (2023). Efektivitas Penguraian Sampah Organik Pasar Menggunakan Larva Black Soldier Fly. *Jurnal Serambi Engineering*, 8(2), 5474–5484. <https://doi.org/10.32672/jse.v8i2.6003>
- U.S. Department of Agriculture. (2024, August 16). *FoodData Central*. <https://fdc.nal.usda.gov/>
- Waldbauer, G. P. (1968). *The Consumption and Utilization of Food by Insects* (pp. 229–288). [https://doi.org/10.1016/S0065-2806\(08\)60230-1](https://doi.org/10.1016/S0065-2806(08)60230-1)
- Wardhana, A. H. (2017). Black Soldier Fly (*Hermetia illucens*) as an Alternative Protein Source for Animal Feed. *Indonesian Bulletin of Animal and Veterinary Sciences*, 26(2), 069. <https://doi.org/10.14334/wartazoa.v26i2.1327>

ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1

lib.unnes.ac.id

Internet Source

1%

2

eprints.unmas.ac.id

Internet Source

1%

3

Submitted to Universitas Amikom

Student Paper

1%

4

ses.library.usyd.edu.au

Internet Source

1%

5

link.springer.com

Internet Source

1%

6

www.mdpi.com

Internet Source

1%

7

www.entomoljournal.com

Internet Source

1%

8

Edo Danilyan, Januarti Jaya Ekaputri, Enny Zulaika, Puput Risdanareni, Didier Snoeck.

"Utilization of soil microbial community to enhance mechanical properties of geopolymer paste for sustainable construction materials", Case Studies in Construction Materials, 2025

Publication

<1%

9

F. Agustina, A. Y. Bagastyo, E. Nurhayati.

"Electro-oxidation of landfill leachate using boron-doped diamond: role of current density, pH and ions", Water Science and Technology, 2019

Publication

<1%

10	backend.orbit.dtu.dk Internet Source	<1 %
11	arccjournals.com Internet Source	<1 %
12	journals.must.ac.ke Internet Source	<1 %
13	Sara Ruschioni, Daniele Duca, Francesca Tulli, Matteo Zarantoniello et al. "Evaluation of Growth Performance and Environmental Impact of <i>Hermetia illucens</i> Larvae Reared on Coffee Silverskins Enriched with <i>Schizochytrium limacinum</i> or <i>Isochrysisgalbana</i> Microalgae", <i>Animals</i> , 2024 Publication	<1 %
14	journal.umpr.ac.id Internet Source	<1 %
15	www.researchgate.net Internet Source	<1 %
16	scholar.sun.ac.za Internet Source	<1 %
17	Gabriel Tirtawijaya, Jin-Hwa Lee, Khawaja Muhammad Imran Bashir, Hae-Jeung Lee, Jae-Suk Choi. "Evaluating the Efficiency of Black Soldier Fly (<i>Hermetia illucens</i>) Larvae in Converting Mackerel Head Waste into Valuable Resources", <i>Animals</i> , 2024 Publication	<1 %
18	R Ismail, A Trisna, V S Tati, M Amran. "Evaluation Nutrient of Larvae Black Soldier Fly (<i>Hermetia Illucens</i>) on Fermented Palm Oil Sludge Culture Media", <i>IOP Conference Series: Earth and Environmental Science</i> , 2023 Publication	<1 %

19	web.archive.org Internet Source	<1 %
20	Shahida Anusha Siddiqui, Özge Süfer, Gülşah Çalışkan Koç, Hanif Lutuf et al. "Enhancing the bioconversion rate and end products of black soldier fly (BSF) treatment – A comprehensive review", Environment, Development and Sustainability, 2024 Publication	<1 %
21	Submitted to University of Venda Student Paper	<1 %
22	Submitted to University of the Philippines - Cebu Student Paper	<1 %
23	Daniel Pleissner, Birgit A. Rumpold. "Utilization of organic residues using heterotrophic microalgae and insects", Waste Management, 2018 Publication	<1 %
24	Submitted to Universitas Jambi Student Paper	<1 %
25	Zeshan, , and Chettiappan Visvanathan. "Evaluation of anaerobic digestate for greenhouse gas emissions at various stages of its management", International Biodeterioration & Biodegradation, 2014. Publication	<1 %
26	eprints.qut.edu.au Internet Source	<1 %
27	repository.itny.ac.id Internet Source	<1 %
28	www.researchsquare.com Internet Source	<1 %

- | | | |
|---|--|----------------|
| <div style="background-color: #008000; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">29</div> | <p>Sila Kriscahyanti, Mika Vernicia Humairo, Agung Kurniawan. "Efektivitas Penambahan Substraksi EM4, MOL, dan Eco-enzyme pada Proses Dekomposisi oleh Larva Black Soldier Fly", <i>ASPIRATOR - Journal of Vector-Borne Diseases Studies</i>, 2025</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #8B4513; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">30</div> | <p>Yaru Zhang, Guoxin Huang, Sicong Chen, Tinghao Yu, Xinrui Ren, Jinhua Xiao, Dawei Huang. "Enhanced waste-to-biomass conversion and reduced nitrogen emissions for black soldier fly larvae (<i>Hermetia illucens</i>) through modifying protein to energy ratio", <i>Journal of Environmental Management</i>, 2025</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #8B4513; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">31</div> | <p>brill.com</p> <p>Internet Source</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #0056b3; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">32</div> | <p>www.lrrd.org</p> <p>Internet Source</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #800080; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">33</div> | <p>www.researcherslinks.com</p> <p>Internet Source</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #8B4513; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">34</div> | <p>Shahida Anusha Siddiqui, Iskandar Azmy Harahap, Jonathan Osei-Owusu, Trideep Saikia et al. "Bioconversion of organic waste by insects – A comprehensive review", <i>Process Safety and Environmental Protection</i>, 2024</p> <p>Publication</p> | <p><1 %</p> |
| <hr/> | | |
| <div style="background-color: #00008B; color: white; display: inline-block; width: 30px; height: 30px; text-align: center; line-height: 30px;">35</div> | <p>Shouyu Zhang, Pu Xiong, Yongsheng Ma, Ning Jin, Shibo Sun, Xiaoying Dong, Xiaodong Li, Jianqiang Xu, Hao Zhou, Weiping Xu. "Transformation of food waste to source of antimicrobial proteins by black soldier fly larvae for defense against marine <i>Vibrio</i></p> | <p><1 %</p> |

36

Slameto Slameto, Danil Eka Fahrudin, Dimvy Rusefani Asetya, Muhamad Wahyu Saputra et al. "The influence of light-emitting diodes and sulfur-silica on the growth, yield, and biochemical content in lettuce Influence of LEDs and sulfur-silica on lettuce", Journal of Applied Biology & Biotechnology, 2025

Publication

<1 %

37

Weiping Tian, Zhouqiang Wang, Yu Xing, Yandong Xie, Yuling Jian, Haizhao Sun. "Black soldier fly larvae (Diptera: Stratiomyidae) enhance stabilization and modulate substrate microbial communities during municipal sewage sludge biotransformation", Journal of Environmental Chemical Engineering, 2025

Publication

<1 %

38

A. Isibika, P. Simha, B. Vinnerås, C. Zurbrügg, O. Kibazohi, C. Lalander. "Food industry waste - An opportunity for black soldier fly larvae protein production in Tanzania", Science of The Total Environment, 2022

Publication

<1 %

39

Aulia Ulfah Farahdiba, I.D.A.A. Warmadewanthi, Yunus Fransiscus, Elsa Rosyidah, Joni Hermana, Adhi Yuniarto. "The present and proposed sustainable food waste treatment technology in Indonesia: A review", Environmental Technology & Innovation, 2023

Publication

<1 %

40

E. Ermolaev, C. Lalander, B. Vinnerås. "Greenhouse gas emissions from small-scale fly larvae composting with *Hermetia illucens*", Waste Management, 2019

<1 %

41 Hanwen Guo, Chengliang Jiang, Zhijian Zhang, Wenjing Lu, Hongtao Wang. "Material flow analysis and life cycle assessment of food waste bioconversion by black soldier fly larvae (*Hermetia illucens* L.)", *Science of The Total Environment*, 2020

Publication

42 Iskandar Zulkarnaen, Eskar Tri Denatara, Anik Budiati, Julia Nurani et al. "Peningkatan Ketahanan Pangan Berbasis Security menggunakan Metode Eco Enzyme pada Perkebunan Sayuran", *Jurnal Kajian Ilmiah*, 2023

Publication

43 Tao Li, Samiullah Khan, Mao Wei, Haiyin Li, Tingchi Wen, Jianjun Guo, Daochao Jin. "Utilizing Black Soldier Fly Larvae to Improve Bioconversion and Reduce Pollution: A Sustainable Method for Efficient Treatment of Mixed Wastes of Wet Distiller Grains and Livestock Manure", *Molecules*, 2023

Publication

44 Wancheng Pang, Dejia Hou, Elhossen E. Nowar, Huanchun Chen, Jibin Zhang, Guoping Zhang, Qing Li, Shuca Wang. "The influence on carbon, nitrogen recycling, and greenhouse gas emissions under different C/N ratios by black soldier fly", *Environmental Science and Pollution Research*, 2020

Publication

45 corpus.ulaval.ca

Internet Source

46 edepot.wur.nl

Internet Source

47	eprints.whiterose.ac.uk Internet Source	<1 %
48	iopscience.iop.org Internet Source	<1 %
49	namibian-studies.com Internet Source	<1 %
50	psasir.upm.edu.my Internet Source	<1 %
51	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
52	repositori.usu.ac.id:8080 Internet Source	<1 %
53	res.mdpi.com Internet Source	<1 %
54	rosdok.uni-rostock.de Internet Source	<1 %
55	www.bdspublishing.com Internet Source	<1 %
56	www.diva-portal.se Internet Source	<1 %
57	www.research-collection.ethz.ch Internet Source	<1 %
58	www.usda.gov Internet Source	<1 %
59	E.M. Nyakeri, M.A. Ayieko, F.A. Amimo, H. Salum, H.J.O. Ogola. "An optimal feeding strategy for black soldier fly larvae biomass production and faecal sludge reduction", Journal of Insects as Food and Feed, 2019 Publication	<1 %
60	Eleonora De Santis, Alberto de Iudicibus, Francesca Lecce, Massimiliano De Mei et al. "A	<1 %

Multidisciplinary Approach for the Development of a Supply Chain in Biomass Conversion of Agrifood Waste Mediated by Larvae of *Hermetia illucens* L.: From Rearing to By-Product Exploitation", Agriculture, 2024

Publication

-
- 61 "Solid Waste Engineering and Management", Springer Science and Business Media LLC, 2022

Publication

-
- 62 A. Fuhrmann, M. Gold, L.H. Lau Heckmann, P. Pedersen, K. Shakhnovich, C.X. Chu, I. Haberkorn, N. Puniamoorthy, A. Mathys. "Comprehensive industry-relevant black soldier fly bioconversion characterisation by a novel chamber system", Waste Management, 2025

Publication

-
- 63 D. Purkayastha, S. Sarkar. "Sustainable waste management using black soldier fly larva: a review", International Journal of Environmental Science and Technology, 2021

Publication

-
- 64 Kashif ur Rehman, Cornelia Schwennen, Christian Visscher, Madeleine Plötz et al. "Closing the loop with pretreatment and black soldier fly technology for recycling lignocellulose-rich organic by-products: A progressive review", Carbohydrate Polymer Technologies and Applications, 2025

Publication

-
- 65 Riang Ursada, Arseto Yekti Bagastyo, Alfian Purnomo. "Effect of feeding rate and composition for bioconversion of wastewater treatment plant sludge from the milk and creamer processing industries using black

soldier fly larvae", BIO Web of Conferences, 2025

Publication

66

ejournal.undip.ac.id
Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On