

Successful of High Risk Pulseless-Balloon Aortic Valvuloplasty Procedure in Uncorrected Pulmonary Atresia with Severe Congenital Aortic Stenosis and Low Left Ventricular Function

Sefri Noventi Sofia^{✉*}, Bahrudin^{**}, Ignatius Riwanto^{***}, Hardhono^{*}, Anindita Soetadji^{*}, Adhi Gunawan Baskoro^{****}, Ananta Sidhi Prawara^{**}, Muhamamad Thifan Satyagraha^{**}, Yovie Kurniawati^{*****}, Radityo Prakoso^{*****}

*Doctoral Study Program of Medical and Health Science, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

**Department of Cardiology, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

***Department of Surgery, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

****Department of Anesthesiology and Intensive Care, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

*****Harapan Kita National Heart Center, Jakarta, Indonesia

✉Correspondence: sefrinoventisofia@mail.com

ABSTRACT

Background: Pulmonary atresia with ventricular septal defect (PA-VSD) and severe bicuspid aortic stenosis (AS) is an uncommon condition that presents significant surgical challenges. The dual obstruction leads to chronic pressure overload, resulting in ventricular hypertrophy and decreased systolic function. A low left ventricular ejection fraction (LVEF) increases the risk of mortality during surgical interventions. Percutaneous balloon aortic valvuloplasty (BAV) poses additional risks due to the intentional induction of pulselessness during balloon inflation. This case report aims to detail the perioperative management strategies employed during BAV in a patient with PA-VSD, severe AS, and low LVEF.

Case: We present a 19-year-old female patient weighing 45 kg with shortness of breath and fatigue. Her oxygen saturation was measured at 90% across all extremities. Electrocardiogram findings indicated sinus rhythm with biventricular hypertrophy and incomplete left bundle branch block. Chest X-ray revealed cardiomegaly and pulmonary artery dilation, while echocardiography confirmed PA-VSD, severe AS with a mean pressure gradient (mPG) of 55 mmHg, and an LVEF of 41.3%. A percutaneous pulseless-BAV was performed using a Tyshak balloon via an antegrade transvenous femoral approach under fluoroscopy and transesophageal echocardiography guidance.

Discussion: Anesthesia management focused on gradual medication titration, minimizing patient movement during the procedure, and ensuring comfort due to Transesophageal Echocardiogram (TEE) probe insertion. A temporary pacemaker was placed in the right ventricle apex for pacing at 220 beats per minute until cardiac arrest occurred post-balloon inflation; defibrillation successfully restored spontaneous circulation.

Conclusion: This case illustrates that percutaneous BAV in uncorrected PA-VSD with severe AS and low LVEF is feasible despite its inherent risks when conducted by a skilled team utilizing careful judgment throughout the procedure.

Keywords: Aortic stenosis, balloon aortic valvuloplasty, cardiovascular anesthesia, congenital heart disease, pulmonary atresia

INTRODUCTION

Pulmonary atresia with ventricular septal defect (PA-VSD) is a rare congenital heart defect, occurring in approximately 7 out of every 100,000 live births.¹ This condition represents a severe form of Tetralogy of Fallot (ToF), necessitating distinct management strategies, particularly in late-presenting patients, where clinical manifestations can vary significantly.² Accurate screening and diagnosis are essential and can be effectively achieved through segmental sequential analysis echocardiography.³

The presence of bicuspid aortic valves in patients with PA-VSD is uncommon.⁴ However, the coexistence of severe bicuspid aortic stenosis alongside PA-VSD has not been previously reported in the literature. This highlights the uniqueness and complexity of our case. The combination of these conditions poses significant challenges due to their impact on left ventricular function, which may be compromised as evidenced by decreased left ventricular ejection fraction (LVEF).

This case report aims to elucidate the management approach for a patient diagnosed with pulmonary atresia, VSD, and severe bicuspid aortic stenosis accompanied by diminished LV function. Specifically, we will detail the high-risk procedure involved in performing pulseless balloon aortic valvuloplasty (BAV). By sharing this experience, we hope to contribute valuable insights into managing similar cases that present unique challenges within pediatric cardiology.

CASE

A 19-year-old woman presented to the outpatient clinic with complaints of easy fatigability for the past year. Her symptoms were relieved by rest and worsened with physical activity. Her parents reported that she often turned blue while crying since the age of 3. There was no history of recurrent respiratory tract infections. She had previously been diagnosed with Tetralogy of Fallot (TOF) but was lost to follow-up.

On physical examination, her vital signs were within normal limits except for an oxygen saturation of 90% in all four extremities. Her height was 151 cm, and her body weight was 45 kg. Cardiac examination revealed a high-pitched ejection systolic murmur at the upper left sternal border. An ECG showed sinus rhythm at 75 beats per minute, left axis deviation, biventricular hypertrophy, and an incomplete left bundle branch block (LBBB) (Figure 1). Chest X-ray revealed cardiomegaly with dilation at the base of the heart, concavity at the region of the pulmonary arteries, a normal broncho-vascular pattern, and a mild left-sided pleural effusion (Figure 2). Laboratory investigations were unremarkable.

Transthoracic echocardiography (TTE) demonstrated situs solitus, concordant atrioventricular (AV) connections, pulmonary atresia, and an overriding aorta of approximately 50%. All pulmonary veins drained into the left atrium. There was no ASD, but a bidirectional shunt was present through a subaortic VSD. The LVEF was decreased at 41.3%, while right ventricular contractility was preserved,

with a tricuspid annular plane systolic excursion (TAPSE) of 25 mm. Severe valvular aortic stenosis with mild to moderate aortic regurgitation (AR) was also noted, caused by a calcified bicuspid aortic valve with post-stenotic dilation (Figure 3). The aortic annulus measured 30 mm, and the descending thoracic aorta measured 15.1 mm in diameter. Mild pericardial effusion was also observed.

Pre-procedural TTE and TEE revealed concentric LVH, decreased LV function, valvar AS with rudimentary noncoronary cusp / NCC (bicuspid physiology), mild AR, overriding 50% of the aorta, with aortic root size was 30 mm, Pulmonary Atresia, VSD subaortic bidirectional shunt. (Figure 4). Given the low LVEF, a decision was made to perform percutaneous balloon aortic valvuloplasty (BAV) using a retrograde transvenous approach. The patient underwent general anesthesia with premedication consisting of fentanyl 100 mcg, propofol 120 mg, midazolam 1 mg, and rocuronium 30 mg. She was intubated using an endotracheal tube (ETT) size 7 and ventilated with FiO₂ 30%, resulting in a post-intubation oxygen saturation of 88%.

The right internal jugular vein was punctured, followed by the insertion of a 5F sheath. A 5F bipolar temporary pacemaker (TPM) lead was introduced and advanced to the right ventricular apex. Pacing threshold was 1 A with an R-wave of 6 mV. The TPM was set at a sensitivity of 3 mV, output of 3 A, and rate of 50 bpm. Defibrillator pads were applied and connected to a defibrillation machine.

A 6F sheath was inserted into the right femoral artery (RFA), and a retrograde transvenous approach for BAV was

initiated. A pigtail catheter was advanced into the aorta via the RFA. A 0.035-inch guidewire was positioned across the aortic valve into the left ventricle. Initially, attempts to cross the severely stenotic aortic valve using a JR guiding catheter were unsuccessful due to the tight valve and tortuous anatomy. The guiding catheter was exchanged for a JR diagnostic catheter, and with the help of a 0.035" exchange-length Terumo wire, the catheter was successfully advanced across the VSD into the aorta. An extra-stiff Lunderquist wire was then placed into the distal descending aorta.

A balloon catheter (Tyshak II, 30 mm x 3 cm x 100 cm) was advanced over the stiff wire and positioned across the aortic valve (Figure 5). Rapid right ventricular pacing was performed using the indwelling TPM to induce pulseless ventricular tachycardia and stabilize balloon position during balloon inflation. The first inflation was suboptimal due to misalignment and incomplete balloon expansion. The TPM rate was reduced, and the patient returned to sinus rhythm.

A second balloon inflation was performed with careful repositioning, and this time the balloon fully expanded at 1.5 atm for 10 seconds. Following balloon deflation, the patient developed pulseless ventricular tachycardia (pVT) (Figure 6). A 200 J DC shock was delivered, followed by one minute of CPR, which successfully restored sinus rhythm with a palpable pulse.

A post-procedural pressure pullback from the ascending aorta to the right ventricle showed no significant residual pressure gradient (Figure 7). The criteria for a successful BAV—reduction of more than 50% in the mean transaortic pressure gradient—were achieved, with the gradient reduced from 55 mmHg to

21 mmHg. The patient was transferred to the intensive care unit (ICU) for monitoring and was sedated overnight.

The following day, the patient was fully awake without any neurological deficits. She showed significant clinical improvement. A follow-up TTE revealed improved LV function with LVEF increased to 66.1%, a reduced mean aortic gradient to 21 mmHg, no worsening of AR, and no increase in pericardial effusion (Figure 8). The patient was extubated and subsequently discharged in stable condition.

DISCUSSION

The present case report details a successful percutaneous balloon aortic valvuloplasty (PBAV) performed on a woman with pulmonary atresia, VSD, and severe bicuspid aortic stenosis, accompanied by diminished LV function.⁵ As noted by Olasińska-Wisniewska et al.⁶, while the significance of the disease is similar in children and adults, the etiology, course, and forms of treatment, including interventional cardiology methods, differ with the patient's age. This highlights the distinct considerations for managing AS in the pediatric population compared to adults.

Low LVEF in late-presenting ToF patients is usually caused by hypoxia; however, in this patient, the oxygen saturation was measured at 90% in all four extremities.⁷ Thus, hypoxia is not suspected to be the cause of the low LVEF. The presence of severe bicuspid aortic stenosis results in high afterload that must be overcome by the ventricles. Signs of this overload can be observed through increased thickness of the left ventricular muscle. When the ventricles fail to compensate for this pressure overload, LVEF decreases.⁸

The procedural success in this specific case, demonstrated by the reduction in the aortic gradient (Figure 7) and the subsequent improvement in left ventricular function with a reduced mean pressure gradient on follow-up TTE (Figure 8), is consistent with the findings of other studies on PBAV for congenital AS. For instance, a retrospective study by Pan et al.⁵ reported that the peak transaortic gradient significantly decreased immediately post-PBAV in patients with congenital AS under echocardiographic guidance. Similarly, Charafeddine et al.⁹ described their experience at a tertiary center, demonstrating the feasibility of balloon valvuloplasty for congenital aortic stenosis with high immediate success rates.

The BAV procedure can be performed using two approaches: the antegrade transvenous approach and the retrograde transarterial approach. Both methods have their own advantages and disadvantages. The antegrade transvenous approach typically requires a transseptal puncture; therefore, the retrograde transarterial approach is generally preferred. However, crossing the severely narrowed aortic cusps can sometimes be challenging. In this case, the antegrade transvenous approach was chosen due to the presence of a subaortic VSD with an overriding aorta, which allowed easier access to the left ventricular outflow tract. The size of the balloon should be carefully selected to maximize the benefits of BAV while minimizing potential risks. A balloon – to – annulus ratio of 0.9 : 1 to 1 : 1 is commonly used to guide balloon sizing. In this case, the aortic annulus measured 30 mm, which is considered quite large. Accordingly, a semi-compliant cylindrical Tyshak II balloon, size 30 mm, was selected to achieve a balloon-

to-annulus ratio of 1:1. While hourglass-shaped balloons, such as the Nucleus balloon, are now preferred for improved positioning and stability during inflation, their availability and size options remain limited.¹⁰

A reduction in the mean transaortic valvular gradient of 50 mmHg or 40–50% from the baseline is considered a criterion for successful BAV. In this patient, the criteria were met, with the initial mean gradient decreasing from 55 mmHg to 21 mmHg, representing a reduction of more than 50%. The patient's LVEF improved, from 41.3% to 66.1%.

Ołasińska-Wisniewska et al.⁶ emphasize that treatment strategies differ with age, suggesting that managing AS in conjunction with other critical congenital heart defects requires a specialized and multidisciplinary approach. The ACC/AHA guidelines on perioperative cardiovascular evaluation and management also underscore the importance of considering underlying cardiac conditions in the context of procedures.¹¹

The utilization of echocardiographic guidance was crucial in this case, as evidenced by the pre-procedural assessments using TTE and TEE (Figures 3 and 4) and intra-procedural monitoring (Figure 6). This is in line with the increasing adoption of echocardiography as the sole or primary guidance modality for PBAV in congenital AS, as demonstrated by Pan et al.⁵, who reported a high success and low complication rate using Doppler echocardiography alone.

While BAV in adults with severe AS is often considered a palliative measure or a bridge to definitive surgical, its role in

congenital AS, particularly in complex cases like ours, is often focused on palliating the stenosis, improving hemodynamics, and potentially delaying or avoiding early surgical intervention.¹² The long-term outcomes of BAV for congenital AS often involve the need for reintervention, and the progression of aortic regurgitation is a known concern.

Considering the broader context of aortic stenosis management, the ESC/EACTS and ACC/AHA guidelines primarily focus on the management of AS in adults.¹³ While these guidelines emphasize the importance of a Heart Team approach and risk assessment^{14,15}, the specific nuances of managing complex congenital heart disease in patients undergoing BAV are not their primary focus. However, the underlying principle of a multidisciplinary team evaluating the risks and benefits of intervention certainly applies to our case.

This case underscores the feasibility and immediate benefits of PBAV in a patient with severe congenital aortic stenosis complicated by pulmonary atresia and a VSD. The successful reduction of the aortic gradient and improvement in left ventricular function highlight the potential of BAV to provide effective palliation in this challenging clinical scenario. Last but not least, it is very important to have great teamwork. Delegate properly to every team member with an exact and clear job description before starting the procedure. Every wrong decision could lead to a worse and fatal outcome.

CONCLUSION

Percutaneous BAV in a patient with uncorrected PA-VSD, severe aortic stenosis, and low LVEF is a challenging and high-risk procedure. However, with careful consideration and a collaborative team approach, the procedure can be

performed successfully, resulting in significant hemodynamic and clinical improvement as a bridge to definitive therapy.

REFERENCES

1. Liang L, Wang Y, Zhang Y. Prenatal Diagnosis of Pulmonary Atresia With Ventricular Septal Defect and an Aberrant Ductus Arteriosus in a Dextrocardia by Two- and Three-Dimensional Echocardiography: A Case Report. *Front Med* [Internet]. 2022;9:904662. Available from: 10.3389/fmed.2022.904662
2. Moradian M. Pulmonary Atresia with VSD (PA-VSD) or Tetralogy of Fallot and Pulmonary Valve Atresia (TF-PA) BT - Atlas of Echocardiography in Pediatrics and Congenital Heart Diseases. In: Moradian M, Alizadehasl A, editors. Berlin, Heidelberg: Springer Berlin Heidelberg; 2021. p. 85–7. Available from: https://doi.org/10.1007/978-3-662-62341-1_27
3. Shakya S, Rajashekar P, Gupta SK. Sequential Segmental Approach to Congenital Heart Disease. *J Indian Acad Echocardiogr Cardiovasc Imaging* [Internet]. 2020;4(3). Available from: https://journals.lww.com/jiae/fulltext/2020/04030/sequential_segmental_approach_to_congenital_heart.5.aspx
4. Zhang ZJ, Zhang MK, Xue H, Fan LX. Ascending aorta dilatation for pulmonary atresia with ventricular septal defect: a report of three adult cases. *J Cardiothorac Surg* [Internet]. 2025 Jan;20(1):64. Available from: 10.1186/s13019-024-03293-7
5. Xie Y, Wang S, Zhao G, Li M, Zhang F, Ouyang W, et al. Percutaneous aortic balloon valvuloplasty under echocardiographic guidance solely. *J Thorac Dis* [Internet]. 2020 Mar;12(3):477–83. Available from: 10.21037/jtd.2020.01.16
6. Olasińska-Wiśniewska A, Trojnarowska O, Grygier M, Lesiak M, Grajek S. Percutaneous balloon aortic valvuloplasty in different age groups. *Postępy w Kardiologii interwencyjnej = Adv Interv Cardiol* [Internet]. 2013;9(1):61–7. Available from: 10.5114/pwki.2013.34029
7. Awasthy N, Radhakrishnan S, Iyer KS, Sharma R. Reversible ventricular dysfunction in cyanotic heart disease. *Indian Heart J* [Internet]. 2014;66(6):704–6. Available from: <https://www.sciencedirect.com/science/article/pii/S0019483214002570>
8. Ito S, Miranda WR, Nkomo VT, Connolly HM, Pislaru S V, Greason KL, et al. Reduced Left Ventricular Ejection Fraction in Patients With Aortic Stenosis. *J Am Coll Cardiol* [Internet]. 2018;71(12):1313–21. Available from: <https://www.sciencedirect.com/science/article/pii/S073510971830322X>
9. Charafeddine FA, Houssein HB, Kibbi NB, El-Rassi IM, Tabbakh AM, Abutaqa MS, et al. Balloon Valvuloplasty for Congenital Aortic Stenosis: Experience at a Tertiary Center in a Developing Country. *J Interv Cardiol* [Internet]. 2021;2021:6681693. Available from: 10.1155/2021/6681693
10. Keeble TR, Khokhar A, Akhtar MM, Mathur A, Weerackody R, Kennon S. Percutaneous balloon aortic valvuloplasty in the era of transcatheter aortic valve implantation: a narrative review.

- Open Hear. 2016;3(2):e000421.
11. Fleisher LA, Fleischmann KE, Auerbach AD, Barnason SA, Beckman JA, Bozkurt B, et al. 2014 ACC/AHA guideline on perioperative cardiovascular evaluation and management of patients undergoing noncardiac surgery: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* [Internet]. 2014 Dec;130(24):2215–45. Available from: [10.1161/CIR.0000000000000105](https://doi.org/10.1161/CIR.0000000000000105)
 12. González LF, Mata RB, Meabe JA, García ML, Miranda JMG. Percutaneous balloon valvuloplasty validity for congenital aortic stenosis. Experience since the beginning of the technique. *An Pediatría (English Ed)* [Internet]. 2022;96(3):265–7. Available from: <https://www.sciencedirect.com/science/article/pii/S2341287922000266>
 13. Lee G, Chikwe J, Milojevic M, Wijeyesundera HC, Biondi-Zoccai G, Flather M, et al. ESC/EACTS vs. ACC/AHA guidelines for the management of severe aortic stenosis. *Eur Heart J* [Internet]. 2023 Mar 7;44(10):796–812. Available from: <https://doi.org/10.1093/eurheartj/ehac803>
 14. Thourani VH, O'Brien SM, Kelly JJ, Cohen DJ, Peterson ED, Mack MJ, et al. Development and Application of a Risk Prediction Model for In-Hospital Stroke After Transcatheter Aortic Valve Replacement: A Report From The Society of Thoracic Surgeons/American College of Cardiology Transcatheter Valve Therapy Registry. *Ann Thorac Surg*. 2019 Apr;107(4):1097–103.
 15. Kiani S, Kamioka N, Black GB, Lu MLR, Lisko JC, Rao B, et al. Development of a Risk Score to Predict New Pacemaker Implantation After Transcatheter Aortic Valve Replacement. *JACC Cardiovasc Interv*. 2019 Nov;12(21):2133–42.






Figure 1 Electrocardiogram (ECG) showing sinus rhythm with biventricular hypertrophy and incomplete left bundle branch block (LBBB).




Figure 2 Chest X-ray demonstrating cardiomegaly with dilatation of the base of the heart and a concave region of the pulmonary arteries



3.1




3.2




3.3

Figure 3 TTE showed severe bicuspid aortic stenosis with mild to moderate AR and a low LVEF

- 3.1 The image displays the velocity-time integral (VTI) and pressure gradients, indicating significant aortic stenosis with a maximum velocity of 453 cm/s and mean pressure gradient of 55 mmHg.
- 3.2 Two-dimensional echocardiographic image showing the left ventricular outflow tract (LVOT) and aortic valve
- 3.3 Key parameters include left ventricular end-diastolic volume (EDV) of 92.04 ml, end-systolic volume (ESV) of 54.03 ml, and an ejection fraction (EF) of 41.3%.



4.1



4.2

Figure 4 Pre-procedure TTE and TEE showed severely doming of the aortic valve

- 4.1 The left panel displays the anatomical structure, while the right panel illustrates the flow dynamics with a distance measurement of 3.00 cm.
- 4.2 The left panel displays the anatomical structure, while the right panel illustrates significant flow dynamics with a peak velocity of 59.2 cm/s.





Figure 5 Tyshak II balloon 30 mm x 3 cm x 100 cm inflated




Figure 6 TEE showed pulseless-heart with VT rhythm during balloon inflation




Figure 7 Ao-RV pullback shows there was nearly no pressure gradient between Ao and RV




Figure 8 TTE evaluation showed mild AS with mild to moderate AR and an improved LVEF

8.1 The parameters include a maximum velocity (V_{max}) of 280 cm/s, mean pressure gradient (Mean PG) of 21 mmHg, and a velocity-time integral (VTI) of 62.9 cm.

8.2 Key parameters include left ventricular end-diastolic volume (EDV) of 100.27 ml, end-systolic volume (ESV) of 33.95 ml, and an ejection fraction (EF) of 66.1%

Review Form JAI

Please rate the paper by filling all fields below

Originality

- ☒ Excellent
- ☐ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Clarity in term of abstract and keyword

- ☐ Excellent
- ☒ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Syntetic method (objectives according to conclusions, methods of research and statistical analysis)

- ☐ Excellent
- ☒ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Systematic preparation of articles (clarity and refer to evidence based)

- ☒ Excellent
- ☐ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Suitability of citation method and writing bibliography with vancouver system

- ☐ Excellent
- ☒ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Impact of articles on science and technology

- ☐ Excellent
- ☒ Good
- ☐ Fair
- ☐ Poor
- ☐ Very Poor

Priority for publication

- ☐ Excellent
- ☒ Good
- ☐ Fair
- ☐ Poor

- Very Poor

Recommendations for editors

- Received without repair
- Received after minor repair
- Need a lot of repair
- Should be repaired and rewritten
- Not suitable for publication

Manuscript Accepted JAI