

## Characteristic of different level of fermented concentrate in the rumen metabolism based on *in vitro*

N.F. Sari\*, R. Ridwan, Rohmatussolihat, R. Fidriyanto, W.D. Astuti and  
Y. Widyastuti

Research Center for Biotechnology, Indonesian Institute of Sciences  
Jl. Raya Bogor km 46, Cibinong, Bogor, West Java, Indonesia 16911

\*Corresponding E-mail : nurulfitrisari@gmail.com

Received January 09, 2018; Accepted March 27, 2018

### ABSTRAK

Kecernaan merupakan gambaran dari kemampuan bahan pakan yang dapat digunakan oleh ternak. Semakin tinggi suatu kecernaan, semakin banyak bahan pakan yang dapat dicerna di dalam saluran pencernaan. Penelitian ini bertujuan untuk mendapatkan proporsi penggunaan konsentrat terfermentasi pada fermentasi rumen secara *in vitro*. Pakan yang digunakan terdiri dari tujuh perlakuan sebagai berikut : P1 (20% *Pennisetum purpureum* : 80% konsentrat), P2 (30% *P.purpureum* : 70% konsentrat), P3 (40% *P.purpureum* : 60% konsentrat), P4 (50% *P.purpureum* : 50% konsentrat), P5 (20% konsentrat: 80% *P.purpureum*), P6 (30% konsentrat: 70% *P.purpureum*) dan P7 (40% konsentrat : 60% *P.purpureum*). Kecernaan bahan kering (KcBK) dan kecernaan bahan organik (KcBO) dilakukan menggunakan metode Theodorou. Produksi gas diukur pada saat sampel diinkubasi pada suhu 39°C di dalam penangas air pada jam ke- 2, 4, 6, 8,10, 12, 24, dan 48. Sebanyak 0,75 g formulasi pakan dicampurkan dengan 75 mL cairan buffer rumen. Cairan rumen diambil dari 3 sapi fistula sebelum pemberian pakan di pagi hari. Rancangan acak lengkap dengan tiga kali ulangan digunakan. Produksi dan kinetika gas diestimasi menggunakan model Orskov. Hasil penelitian menunjukkan bahwa pada perlakuan P2 dan P3 menunjukkan kecernaan tertinggi dibanding perlakuan lainnya dengan KcBK masing-masing sebesar 71,63% ; 71,06% dan KcBO masing-masing sebesar 76,42% ; 71.65%. Disimpulkan bahwa perlakuan P2 dan P3 dapat digunakan sebagai pakan untuk sapi potong.

*Kata kunci* : sapi; fermentasi rumen, konsentrat terfermentasi, kecernaan, *in vitro*.

### ABSTRACT

Digestibility is a description of the ability of feed material that can be utilized by livestock. Higher digestibility of feed material means the total amount of feed content that can be digested in the digestive tract. This study aimed to obtain the proportion of the use of fermentation concentrate in rumen fermentation based on *in vitro*. The feed used consisted of 7 treatments as follows P1 (20% *Pennisetum purpureum*: 80% concentrate), P2 (30% *P.purpureum* : 70% concentrate), P3 (40% *P.purpureum* : 60% concentrate), P4 (50% *P.purpureum* : 50% concentrate), P5 (20% concentrate: 80% *P.purpureum*), P6 (30% concentrate: 70% *P.purpureum*) and P7 (40% concentrate: 60% *P.purpureum*). Dry matter digestibility (DMD) and organic matter digestibility (OMD) were done using Theodorou method. Gas production was measured when the samples were incubated at 39°C in the waterbath incubator at hours 2, 4, 6, 8, 10, 12, 24, and 48. Approximately 0.75 g of feed formulation was mixed with 75 mL of rumen fluid buffer. The rumen fluid is taken from 3 fistulated cattles prior to morning feeding. A complete randomized design with three replicates was used. Asymptotic gas production and

kinetics were estimated by the Orskov's model. The results of this study showed that P2 and P3 treatment showed the best digestibility among others treatment with 71.63% ; 71.06% of DMD and 76.42% ; 71.65% of OMD. In conclusion, P2 and P3 treatment can be used as feed for beef cattle.

*Keywords: cattle, rumen fermentation, fermented concentrate, digestibility, in vitro*

## INTRODUCTION

The improper storage technique of concentrate for a long time period and in humid conditions can trigger contamination of concentrate by microorganisms causing rancid odor (Canibe and Jensen, 2012). This may affect the nutritional quality of the concentrate as well as the palatability of feed. Therefore, to reduce the contamination of the concentrate by microorganisms, it can be processed through fermentation. The fermentation technique can inhibit the growth of microbial pathogens and also can improve the quality of nutrient concentrates (Lee and Mau., 2008; Cao *et al.*, 2012). In addition, the process of fermentation increases the vitamins, enzymes, and growth factors of fermented products (Ng *et al.*, 2011; Dei *et al.*, 2008).

The use of microbial inoculants to ferment concentrate have been commonly studied in livestock, especially in pigs and poultry. Nevertheless, there have been limited studies of the use of microbial inoculant to ferment concentrate in ruminant. In addition, there have been few studies about the use of lactic acid bacteria in fermented concentrate could increase the availability of phosphorus in cereal based-diets, able to reduce IP6 (Reale *et al.*, 2004 ). The effects of fermentation on feed depend on the activity and type of microbial used (Niba *et al.*, 2009).

Many researchers reported the beneficial of the use of microbial inoculant in fermented concentrate. As reported by Niba *et al* (2008) that the use of lactic acid bacteria can increase the concentration of lactic acid. Ahmed *et al* (2016) also reported that fermented feed contains large number of Lactobacilli with high concentrations of lactic acid and other volatile fatty acids and has a low pH. Moreover, Kobashi *et al* (2008) reported that the proportion of chlortetracycline-resistant *Escherichia coli* strains was significantly reduced in the gut of weaned piglets compared with dry feed. Fermented feed may provide more rich nutrition, better palatability, and increased digestibility so the needs of cattle protein could be met and could improve the performance of beef

cattle (Niba *et al.*, 2009; Humer *et al.*, 2013). However, further research is needed to determine optimal proportion of fermented concentrate. The objective of this study was to characterize the effects of different level of fermented concentrate and elephant grass on the rumen fermentation in vitro.

## MATERIALS AND METHODS

### Sample preparation

Concentrate was made from the following formulation : 22% of rice bran, 5% of maize flour, 5% of coffee pulp, 10% of corn gluten feed, 6% of cassava flour waste, 9% of coconut meal, 18% of palm kernel cake, 18% of pollard brand, 6% of soy bean meal, and 1% of mineral mix. The formulation concentrate was fermented by adding 2% of *L. plantarum* ( $2.5 \times 10^6$  cfu/mL) and kept for 24 hours incubation. *Lactobacillus plantarum* TSD-10 was from the laboratory of Applied Microbiology, Research Center for Biotechnology. *Pennisetum purpureum* were freshly collected from Agrostology Laboratory Field, Research Center for Biotechnology, LIPI. These materials were oven-dried at 60°C for 24 h, milled using a grinder to pass a 1 mm sieve.

### Chemical and pH analysis

Samples of *P.purpureum* and fermented concentrate were analyzed for dry matter (DM) and ash according to AOAC (2005). Crude protein (CP), crude fiber (CF), ether extract (EE) were determined followed the manufacture procedure of FOSS. The NH<sub>3</sub>-N concentration was determined by a Conway micro-diffusion method. Volatile Fatty Acids (VFA) were determined in centrifuged samples (1mL) by Gas Chromatography (GC) as described by Goering and Van Soest (1970). pH measurement of mix buffer rumen fluids were measured using pH meter (Jenway Model 3505, UK). Prior to the measurements, pH meter was calibrated by using pH 7 buffer solution and it was prepared carefully as the manufactures instructions. All chemical composition analyses were conducted in triplicate.

Table 1. Chemical Composition (g/kg DM)

Item <sup>a</sup>	(%) DM	(%) CP	(%) CF	(%) EE	pH
<i>Pennisetum purpureum</i>	97.40	9.04	29.72	1.53	
Concentrate	89.37	13.70	21.14	1.85	6.24
Fermented Concentrate	89.96	14.82	18.13	1.70	4.47

DM= dry matter; CP= crude protein; CF= crude fiber; EE= ether extract ; <sup>a</sup>Data are the means of three replicate analysis.




Figure 1. pH value of fermented concentrate. P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate ; P3 = 40% *P.purpureum* : 60% concentrate ; P4 = 50% *P.purpureum*: 50% concentrate ; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum* ; P7 = 40% concentrate: 60% *P.purpureum*

### ***In vitro* Incubation and Gas Production**

Samples of *P.purpureum* and fermented concentrate were arranged and mixed according to the following experimental treatments (dry matter basis). These treatments were subjected to *in vitro* analysis according to Theodorou *et al.* (1994):

- P1 : 20% *P. purpureum* : 80% concentrate feeds
- P2 : 30% *P. purpureum* : 70% concentrate feeds
- P3 : 40% *P. purpureum* : 60% concentrate feeds
- P4 : 50% *P. purpureum* : 50% concentrate feeds
- P5 : 80% *P. purpureum* : 20% concentrate feeds
- P6 : 70% *P. purpureum* : 30% concentrate feeds
- P7 : 60% *P. purpureum* : 40% concentrate feeds

Rumen contents were obtained from three fistulated cattles, then was mixed and strained through four layers of cheesecloth into an erlenmeyer flask with headspace. The use of the cattle in this experiment was approved No. 9879/WK/HK/XI/2015 by the Ethic Clearance Committee of Indonesian Institute of Sciences. Approximately 0.75 g of substrates were put inside the serum bottle glass and filled with 73

mL mixture of 24.33 mL rumen fluid and 48.67 mL of Mc'Dougall buffer (NaHCO<sub>3</sub> 0.98 g, Na<sub>2</sub>HPO<sub>4</sub>.7H<sub>2</sub>O 0.7 g, KCl 0.057 g, NaCl 0.0472 g, MgSO<sub>4</sub>.7H<sub>2</sub>O 0.012 g, CaCl 0.004 g, and H<sub>2</sub>O up to 100 mL). Anaerobic condition in the bottles was achieved by filling CO<sub>2</sub> gas until saturated. Serum bottles were sealed with butyl rubber stoppers and aluminum crimp seals shortly before starting the incubation. The bottles were placed in a water bath at 39°C for 48 h. After 2, 4, 6, 8, 10, 12, 24 and 48 h of incubation, gas production from each serum bottle was vented and recorded by using disposable syringe.

*In vitro* gas production was analysed in seven different treatments of P1, P2, P3, P4, P5, P6, and P7 in three replicates. In addition, three blank (no sample; rumen fluid + buffer mixture) were used to calculate the total gas production. Gas production kinetics were estimated by following Ørskov method (Ørskov and McDonald, 1979).

Table 2. Gas Production Kinetics during 48 Hours of Incubation

Time (Hr)	Treatment (mL)							
	Blank	P1	P2	P3	P4	P5	P6	P7
2	0 ± 0	8.5±1.5 <sup>ab</sup>	5.2±2.3 <sup>a</sup>	6.2±3.6 <sup>a</sup>	8.8±2.85 <sup>ab</sup>	20.0 ± 6.6 <sup>c</sup>	20.50±0.5 <sup>c</sup>	13.8 ± 5.4 <sup>bc</sup>
4	0 ± 0	18.7±2.8 <sup>ab</sup>	12.3±3.55 <sup>a</sup>	14.0±7.2 <sup>a</sup>	19.5±5.0 <sup>ab</sup>	37.3 ± 10.3 <sup>c</sup>	38.2± 0.8 <sup>c</sup>	28.3 ± 9.8 <sup>bc</sup>
6	1.2±1.0	35.5±4.3 <sup>ab</sup>	22.3±3.9 <sup>a</sup>	25.3±12.0 <sup>a</sup>	34.6 ± 7.3 <sup>ab</sup>	60.0 ± 13.3 <sup>c</sup>	62.0±03.9 <sup>c</sup>	45.8 ± 14.6 <sup>bc</sup>
8	2.7±4.5	53.5±3.8 <sup>ab</sup>	34.0±3.9 <sup>a</sup>	37.7±14.7 <sup>a</sup>	50.7±7.9 <sup>ab</sup>	78.5±14.5 <sup>c</sup>	79.8±4.0 <sup>c</sup>	61.7 ± 16.4 <sup>bc</sup>
10	7.3±6.8	79.3±3.8 <sup>c</sup>	51.7±3.6 <sup>a</sup>	55.3±17.5 <sup>ab</sup>	74.2±865 <sup>bc</sup>	101.3±13.8 <sup>d</sup>	101.7±3.7 <sup>d</sup>	8.02 ± 17.8 <sup>cd</sup>
12	14.3±7.4	95.3±2.8 <sup>b</sup>	65.7±3.3 <sup>a</sup>	70.0±17.7 <sup>a</sup>	90.8±8.8 <sup>b</sup>	117.7±14.4 <sup>c</sup>	118 ± 3.9 <sup>c</sup>	97.8 ± 18.8 <sup>bc</sup>
24	27.5±6.1	137.8±3.3 <sup>b</sup>	97.5±3.6 <sup>a</sup>	110.5 ± 23.4 <sup>a</sup>	143.2±15.5 <sup>b</sup>	173.2±13.3 <sup>c</sup>	175.0±5.7 <sup>c</sup>	154.8±19.6 <sup>bc</sup>
48	37.5±6.1	165.0 ± 3.8 <sup>bc</sup>	119.5±3.6 <sup>a</sup>	141.5±23.4 <sup>ab</sup>	175.7±15.9 <sup>bc</sup>	212.8±17.9 <sup>e</sup>	216.3±7.4 <sup>e</sup>	194.5±21.3 <sup>de</sup>

P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate; P3 = 40% *P.purpureum* : 60% concentrate; P4 = 50% *P.purpureum*: 50% concentrate; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum*; P7 = 40% concentrate: 60% *P.purpureum*; a-e Different superscripts within the same column are significantly different at P<0.05.




Figure 2. Total Gas Production during 48 Hours of Incubation. P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate; P3 = 40% *P.purpureum* : 60% concentrate; P4 = 50% *P.purpureum*: 50% concentrate; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum*; P7 = 40% concentrate: 60% *P.purpureum*

### Microbiology analysis

Total lactic acid bacteria and rumen bacteria were counted as colony forming units (CFU) per mL. The lactic acid bacteria were grown on MRS agar medium and counted by using total plate count (Cappucino, 2001), while the rumen

bacteria on 98-5 agar medium were grown anaerobically by following the method of roll tube (Hungate, 1969). Incubation was done at 38<sup>0</sup>C for 3-4 days. Aliquots of fresh filtrate (1 mL) were added to 1 mL of protozoal staining solution (MFS solution, Ogimoto and Imai, 1987). Ciliate




Figure 3. The Comparison among Total Bacteria, Total Lactic Acid Bacteria and Protozoa in Each Treatment . P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate; P3 = 40% *P.purpureum* : 60% concentrate; P4 = 50% *P.purpureum*: 50% concentrate; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum*; P7 = 40% concentrate: 60% *P.purpureum*

protozoa were counted as described by Veira *et al.* (1983) using a Fuchs-Rosenthal counting chamber. Duplicate preparations of each sample were counted, and if either value differed from the average by more than 5%, the counts were repeated. After being incubated for 24 hours, protozoa can be observed by using microscope at 200x magnification

#### Statistical analysis

The experiment design was a block randomized. The statistical analysis of data was performed using SPSS 16 for Windows. The difference among groups was determined by oneway ANOVA analysis. The Duncan's Multiple Range Test, one of the multiple comparison tests, was used when the difference among groups was found to be significant. Significance was defined at  $P < 0.05$ . Microbial diversity data were analyzed descriptively.

## RESULTS AND DISCUSSIONS

#### Chemical and pH analysis

Chemical analysis of feed material is very important to evaluate the nutrient content of feed. Proximate analysis is one of parameter to

determine the quality of feed material. The nutrition composition of feed material that used in this study is shown in Table 1.

The pH rumen indicate rumen conditions that are appropriate or inappropriate for the growth of rumen microbes. This study showed that the use of different levels of fermented concentrate did not affect the pH of rumen (Figure 1). The same pH values as control showed that the use of different levels of fermented concentrate does not interfere rumen fermentation activity. According to Dehority (2004), the rumen's normal pH is about 5.5-7.0. The pH value actually represents the amount of acid produced by microbial activity in the rumen (Elghandour *et al.*, 2015; Hutton *et al.*, 2012). The lowest pH value is generally achieved about 2-6 hours after meals, in accordance with the maximum acid production. Changes in pH are affected by time after meal, lactic acid supply, feed properties, and feeding frequency of animal. (Dehority, 2004; Seo *et al.* 2010).

#### *In vitro* Incubation and Gas Production

Total gas production for 48 hours incubation is presented in Table 2. The gas production comes from direct substrate fermentation ( $\text{CO}_2$  and  $\text{CH}_4$ )

Table 3. Production of VFA and NH<sub>3</sub> during 48 h Incubation

Variable	Treatment						
	P1	P2	P3	P4	P5	P6	P7
C <sub>2</sub>	44.51 <sup>a</sup>	41.38 <sup>a</sup>	42.94 <sup>a</sup>	43.86 <sup>a</sup>	39.48 <sup>a</sup>	43.65 <sup>a</sup>	47.43 <sup>b</sup>
C <sub>3</sub>	30.17 <sup>b</sup>	31.40 <sup>b</sup>	31.89 <sup>b</sup>	27.84	21.57	23.99	24.46
C <sub>4</sub>	13.45 <sup>a</sup>	14.71 <sup>a</sup>	13.73 <sup>a</sup>	16.28 <sup>ab</sup>	19.62 <sup>ab</sup>	16.64 <sup>ab</sup>	12.73 <sup>a</sup>
Iso-C <sub>4</sub>	5.26 <sup>a</sup>	7.08 <sup>a</sup>	5.56 <sup>a</sup>	5.87 <sup>a</sup>	13.11 <sup>ab</sup>	9.33 <sup>a</sup>	7.95 <sup>a</sup>
C <sub>5</sub>	2.5 <sup>a</sup>	1.27 <sup>a</sup>	1.59 <sup>a</sup>	1.82 <sup>a</sup>	1.53 <sup>a</sup>	2.16 <sup>a</sup>	3.26 <sup>ab</sup>
Iso-C <sub>5</sub>	4.11 <sup>a</sup>	4.16 <sup>a</sup>	4.29 <sup>a</sup>	4.33 <sup>a</sup>	4.69 <sup>a</sup>	4.23 <sup>a</sup>	4.17 <sup>a</sup>
C <sub>2</sub> /C <sub>3</sub>	1.47 <sup>a</sup>	1.31 <sup>a</sup>	1.35 <sup>a</sup>	1.58 <sup>a</sup>	1.83 <sup>a</sup>	1.82 <sup>a</sup>	1.94 <sup>a</sup>
NH <sub>3</sub>	6.09 <sup>bc</sup>	6.59 <sup>c</sup>	6.39 <sup>c</sup>	6.94 <sup>c</sup>	4.98 <sup>b</sup>	5.00 <sup>b</sup>	4.80 <sup>b</sup>

P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate; P3 = 40% *P.purpureum* : 60% concentrate; P4 = 50% *P.purpureum*: 50% concentrate; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum*; P7 = 40% concentrate: 60% *P.purpureum*; a-eDifferent superscripts within the same column are significantly different at P<0.05.

and indirect gas production (CO<sub>2</sub>) reactions from organic acids with bicarbonate buffers (Goel *et al.*, 2008; McDonald *et al.* 2010; Hook *et al.*, 2010). In addition, according to McDonald *et al.* (2010) gas composition in the rumen is consist of 40% carbon dioxide (CO<sub>2</sub>), 30-40% methane (CH<sub>4</sub>), 5% hydrogen, and a small portion of oxygen and nitrogen. Based on 24 hours incubation presented in Table 2, it was showed that the average of gas production was over 75% of the total gas during 48 hours incubation, this result is coherent with Jayanegara *et al.* (2009), which stated that more than 75% of the gas is generated after 24 hours of incubation period. Differences in gas production between treatments can be caused by rumen fluids, incubation conditions, and also substrates. Table 2 showed that the rate of gas production of each treatment decreased every hour in accordance with the reduced amount of substrate that can be fermented during the incubation period. Therefore, the rate of gas production will demonstrate the time required for rumen microorganisms to digest the entire substrate.

Figure 2 showed that the gas was still being produced until 48 hours of incubation time. This means that the tested substrate remained until the 48 hours of incubation. Consequently, the zero point of gas production could not be known. It takes up to 96 hours to obtain an accurate kinetic

coefficient (Jayanegara *et al.*, 2009) because incubation for 96 hours the substrate will be exhausted. If the substrate has been fermented out, then the zero point of the gas production rate will be known and the shape of the graph will be constant by forming horizontally as in the blank (Figure 2).

#### Microbiology Analysis

In this study, it was showed in the Figure 3 that the treatment of P1, P2, P3, and P4 had higher amounts of lactic acid bacteria than other treatments. In contrast, the total number of bacteria on treatment P1, P2, P3, and P4 was decreased. We assumed that *L.plantarum* TSD 10 has an antimicrobial ability that can inhibit the presence of pathogenic bacteria in the rumen thus affecting the reduction of total bacteria in the rumen. Fermentation with with *L. plantarum* might prevent harmful microbe in the digestive tract and increase the immune enhancing activity of fermented products (Rizzello *et al.*, 2013; Ashraf and Shah., 2014; Fu *et al.*, 2014; Li *et al.*, 2012).

In addition, concentrate which is fermented by lactic acid bacteria not only as a preservation, but also as probiotic , therefore animals can obtain both nutrients and the probiotics from fermented feed (Sánchez *et al.*, 2015; Hu *et al.*, 2015). Ellis *et al.* (2016) found that the viable cells of lactic

Table 4. Dry Matter Digestibility (DMD) and Organic Matter Digestibility (OMD) *In vitro*

Treatment	DMD (%)	OMD (%)
P1	65.80 ± 1.32 <sup>cd</sup>	66.48 ± 1.53 <sup>b</sup>
P2	71.63 ± 2.73 <sup>d</sup>	76.42 ± 2.39 <sup>c</sup>
P3	71.06 ± 3.69 <sup>d</sup>	71.65 ± 3.93 <sup>c</sup>
P4	60.36 ± 2.94 <sup>bc</sup>	60.76 ± 4.52 <sup>b</sup>
P5	50.89 ± 3.49 <sup>a</sup>	59.28 ± 1.75 <sup>b</sup>
P6	55.07 ± 5.81 <sup>ab</sup>	45.49 ± 5.75 <sup>a</sup>
P7	50.03 ± 1.59 <sup>a</sup>	46.93 ± 4.54 <sup>a</sup>

P1 = 20% *Pennisetum purpureum*: 80% concentrate; P2 = 30% *P.purpureum*: 70% concentrate; P3 = 40% *P.purpureum* : 60% concentrate; P4 = 50% *P.purpureum*: 50% concentrate; P5 = 20% concentrate: 80% *P.purpureum*; P6 = 30% concentrate: 70% *P.purpureum*; P7 = 40% concentrate: 60% *P.purpureum*; a-e Different superscripts within the same column are significantly different at P<0.05.

acid bacteria in silage can play a role as probiotic in the rumen during fermentation process. Sari *et al.* (2017) also reported that lactic acid bacteria has an important role for producing lactic acid, which results in quicker drop of pH. This finding suggested that *L. plantarum* population in fermented concentrate was sufficiently established during fermentation process and able to utilize the high concentration of NH<sub>3</sub>-N. The NH<sub>3</sub>-N in the fermented concentrate was used as a source of N for microbial protein synthesis and for microbial growth (Calsamiglia *et al.*, 2010; Belanche *et al.*, 2012b).

The population of ciliated protozoa is more dominant in the rumen, while the population of flagellated protozoa is minor (Dehority, 2004). In this study the protozoa population presented in Figure 3 in the treatment of P4, P5, P6, and P7 were higher among other treatments. This is because the presence of protozoa naturally develops in the rumen to assist digestion of nutrients from forage crops containing high crude fiber (Noziere *et al.*, 2011). However, the impact of protozoa on the rumen digestion depends on their concentration and the generic composition of their population (Belanche *et al.*, 2012a).

### NH<sub>3</sub> and VFA Production

Ammonia is an important component for the synthesis of amino acids and microbial cell proteins. Protein from feed will be degraded by proteolytic bacteria with the help of protease enzyme into peptide then hydrolyzed to amino acid, and amino acid will go through deamination process to become ammonia. McDonald *et al.* (2010) states that ammonia is produced in conjunction with several small peptides and free amino acids, then it is utilized by rumen organisms for microbial protein synthesis. In addition, some of the amino acids will be further digested into organic acids, ammonia (NH<sub>3</sub>), and carbon dioxide (CO<sub>2</sub>).

The results of the average concentration of NH<sub>3</sub> are presented in Table 3. This study showed that proportion of fermented concentrate can increase the production of NH<sub>3</sub> in the treatment of P2, P3 and P4, while in the treatment of P5, P6, and P7 the amount of NH<sub>3</sub> decreases. This is due to the tendency of increasing bacterial population in the treatment of P5, P6, and P7 (Figure 3) will reduce the value of NH<sub>3</sub> concentration due to NH<sub>3</sub> is an important source of N for rumen microbes for the microbial synthesis process.

Production of VFA after incubation for 48 hours is presented in Table 3. The use of feed which is dominated by fermented concentrate such as P1, P2, and P3 had significant effect (P<0.05) on propionic acid formation. In contrast, the use of feed which is dominated by forage on treatment P7 had significant effect (P <0.05) on acetate production.

C<sub>2</sub>/C<sub>3</sub> ratio shows a pattern of microbial fermentation in the rumen that has the potential to form acetic acid or propionate. The lowest C<sub>2</sub>/C<sub>3</sub> ratio is produced by P2 as shown in Table 3, an be explained that in this treatment has a higher fermentable organic content than others with 76.42% of IVOMD (Table. 4). The low of C<sub>2</sub>/C<sub>3</sub> ratio is due to the high amount of fermentable organic matter enabling the formation of higher propionic acid than acetic acid. As reported by Moran (2005) that the fermentative digestion of carbohydrates especially starch will produce lactic acid, this lactic acid will be converted to propionic acid by microbial lactate utilization, such as *Propionibacteria sp*, *Veillonella alkalescens* and *Peptostreptococcus elsdeini*. The low of C<sub>2</sub>/C<sub>3</sub> ratio will stimulate fattening and tend to body fat formation. In the treatment of P5, P6, and P7 showed higher C<sub>2</sub>/C<sub>3</sub> ratio values than other treatments, this indicates that the microbial

fermentation pattern leads to the formation of acetic acid because the carbohydrates contained in the feed are still dominated by fibrous carbohydrates and microbes that grow well are cellulolytic and hemicellulolytic microbes.

### **Dry Matter Digestibility (DMD) and Organic Matter Digestibility (OMD)**

The results of DMD and OMD during 48 hours incubation for all treatments are presented in Table 4. Dry matter digestibility value is one of indication for feed quality, the higher digestibility value, the more nutrition will be absorbed by the body.

Based on DMD value it was found that P2 and P3 resulted the highest percentage of DMD with 71.63 and 71.06 respectively. It also has a similarity in OMD pattern, that P2 and P3 resulted the highest percentage of OMD with 76.42 dan 71.65 respectively. This study found that digestibility is affected by the presence of protozoa. The lower protozoa population, the higher digestibility value will be generated. This is because protozoa will consume fibrolytic bacteria so that the population will be less and will affect the process of feed degradation in the rumen. This is evidenced by the number of protozoa in the treatment of P2 and P3 is less than other treatments (Figure 3). In Table 4. It is showed that the increase of DMD value will increase the OMD value. This is in accordance with as reported by Muhtarudin and Liman (2006), that the dry matter digestibility will be in line with the organic matter digestibility value. Digestibility can also be affected by components in the feed. As reported by De Boever *et al.* (2005) that the structural components of plant feed material such as cellulose, lignin, cell wall, NDF and ADF will reduce the nutrient digestibility of feed, whereas soluble carbohydrates (starch) and crude protein can increase the nutrient digestibility.

### **CONCLUSION**

Based on this study it could be concluded that fermented concentrate with the range proportion from 60 -70% (P2 and P3) can be used for feed, particularly in beef cattle. In addition, characteristic of fermented concentrate can improve the digestibility of feed in the rumen metabolism.

### **ACKNOWLEDGMENTS**

This research was funded by Unggulan-LIPI DIPA Grant for Food Security and Drug's Discovery of the Research Center for Biotechnology, Indonesian Institute of Sciences (2015-2017). In addition, the authors would like to thank the Center of Excellence of Biotechnology Animal of Beef Cattle and Dairy Cattle (PUI Kemenristekdikti 2018) for supporting this publication.

### **REFERENCES**

- Ahmed, S.T., Mun, H.S., Islam, M., M., Ko, S.Y., Yang, C.J. 2016. Effects of dietary natural and fermented herb combination on growth performance, carcass traits and meat quality in grower-finisher pigs. *Meat Sci*, 122 :7-15.
- AOAC. 2005. Official Methods of Analysis. 18th edition. AOAC International, Arlington, VA, USA
- Ashraf, R. and N. P. Shah. 2014. Immune system stimulation by probiotic microorganisms. *Critical reviews in Food Science and Nutrition*. 54(7):938-956.
- Belanche, A., G. delaFuente, J.M. Moorby and C.J. Newbold. 2012a. Bacterial protein degradation by different rumen protozoal groups. *J.Anim.Sci*. 90:4495-4504
- Belanche, A., M. Doreau, J.E. Edwards, J.M. Moorby, E. Pinloche and C.J. Newbold. 2012b. Shifts in the rumen microbiota due to the type of carbohydrate and level of protein ingested by dairy cattle areas associated with changes in rumen fermentation. *J.Nutr*. 142:1684-1692.
- Calsamiglia, S., A. Ferret, C. Reynolds, N. Kristensen and A. Van Vuuren. 2010. Strategies for optimizing nitrogen use by ruminants. *Animal*. 4(7) : 1184-1196.
- Canibe, N and B.B. Jensen. 2012. Fermented liquid feed—Microbial and nutritional aspects and impact on enteric diseases in pigs. *Anim. Feed Sci. Technol*. 173:17-40.
- Cao, F. L., X.H. Zhang, W.W. Yu, L.G. Zhao and T. Wang. 2012. Effect of feeding fermented Ginkgo biloba leaves on growth performance, meat quality, and lipid metabolism in broilers. *Poult. Sci*. 91: 1210-1221.
- Cappuccino, J.G and N. Sherman. 2001. *Microbiology: A Laboratory Manual*. 6th



- Edition, Benjamin Cummings, San Francisco.
- De Boever, J.L., J.M. Aerts, J.M. Vanacker and D.L. De Brabander. 2005. Evaluation of the nutritive value of maize silages using a gas production technique. *Anim. Feed Sci. Technol.* 123-124: 255-265.
- Dehority, B. A and P. A. Tirabasso. 2004. Effect of feeding frequency on bacterial and fungal concentration, pH, and other parameters in the rumen. *J. Anim. Sci.* 79 : 2908 - 2912.
- Dei, H. K., S.P. Rose, A.M. Mackenzie and R. Amarowicz. 2008. Growth performance of broiler chickens fed diets containing shea nut (*Vitellaria paradoxa*, Gaertn.) meal fermented with *Aspergillus niger*. *Poult. Sci.* 87:1773-1778.
- Elghandour M M, A.Z. Salem, J.S.M. Castañeda, L.M. Camacho, A.E. Kholif and J.C.V. Chagoyán. 2015. Direct-fed microbes: A tool for improving the utilization of low quality roughages in ruminants. *Journal of Integrative Agriculture.* 14:526-533.
- Ellis, J. L., A. Bannink, I. K. Hindrichsen, R. D. Kinley and W. F. Pellikaan. 2016. The effect of lactic acid bacteria included as a probiotic or silage inoculant on in vitro rumen digestibility, total gas, and methane production. *Anim. Feed Sci. Technol.* 211: 61-74.
- Fu, R., R. Chen., Y. Huang, H. Zhao and H. Li H., 2014. Effects of fermented protein feed on the antioxidant ability and immune ability of growing-finishing pigs. *Modern. J. Anim. Husbandry Vet. Med.* 2:21-24.
- Goel, G., H.P.S. Makkar and K. Becker. 2008. Effects of *Sesbania Sesban* and *Carduus Pycnocephalus* leaves and fenugreek (*Trigonella Foenum-Graecum* L.) seeds and their extracts on partitioning of nutrients from roughage- and concentrate based feeds to methane. *Anim. Feed Sci. Tech.* 147:72-89.
- Goering, H. K and P. J. Van Soest. 1970. Forage Fiber Analyses (Apparatus, Reagents, Procedures, and Some Applications). Agriculture handbook no. 379, Agriculture Research Service USDA. Washington (DC), USA.
- Hook, S.E., Wright, A.D.G., McBride, B.W. 2010. Methanogens: methane producers of the rumen and mitigation strategies. *Archaea* : 945785.
- Humer, E., W. Wetscherek, C. Schwarz, and K. Schedle. 2013. Effect of maize conservation technique and phytase supplementation on total tract apparent digestibility of phosphorus, calcium, ash, dry matter, organic matter and crude protein in growing pigs. *Anim. Feed Sci. Technol.* 185:70-77.
- Hungate, R. E. 1969. A Roll Tube Method for Cultivation of Strict Anaerobes. Academic Press. London.
- Hutton, P G, Z. Durmic, E.L. Ghisalberti, G.R. Flematti, R.M. Duncan, C.F. Carson and T.V. Riley. 2012. Inhibition of ruminal bacteria involved in lactic acid metabolism by extracts from Australian plants. *Anim. Feed Sci. Technol.* 176:170-177.
- Jayanegara, A., A. Sofyan, H.P.S. Makkar and K. Becker .2009. Kinetika produksi gas, kecernaan bahan organik dan produksi gas metana in vitro pada hay dan jerami yang disuplementasi hijauan mengandung tanin. *Med Pet.* 32(2):120-129.
- Kobashi Y., H. Ohmori, K. Tajima, T. Kawashima and H. Uchiyama. 2008. Reduction of chlortetracycline-resistant *Escherichia coli* in weaned piglets fed fermented liquid feed. *Anaerobe* 14: 201-204.
- Lee, Y. L., J.H. Yang and J.L. Mau. 2008. Antioxidant properties of water extracts from *Monascus* fermented soybeans. *Food Chem.* 106:1128-1137.
- Li, G., P.J. Brown, J.X. Tang, J. Xu, E.M. Quardokus, C. Fuqua and Y.V. Brun. 2012. Surface contact stimulates the just-in-time deployment of bacterial adhesins. *Mol. Microbiol.* 83:41-51.
- McDonald P., R. Edwards and J. Greenhalgh. 2010. *Animal Nutrition.* 7th Edition. Ashford Colour Press. New York.
- Moran, J. 2005. *Tropical Dairy Farming.* Landlinks Press. Australia.
- Muhtarudin and Liman. 2006. Penentuan tingkat penggunaan mineral organik untuk memperbaiki bioproses rumen pada kambing secara in vitro. *JIPI.* 8(2):132-140.
- Ng, C. C., C.Y. Wang, Y.P. Wang, W.S. Tzeng and Y.T. Shyu. 2011. Lactic acid bacterial fermentation on the production of functional antioxidant herbal *Anoetochilus formosanus* Hayata. *J. Biosci. Bioengineer.* 111:289-293.
- Niba, A. T., J. D. Beal, A. C. Kudi and P. H. Brooks. 2009. Potential of bacterial fermentation as a biosafe method of

- improving feeds for pigs and poultry. *Afr. J. Biotechnol.* 8(9):1758-176.
- Niba, A.T., H. Kouchika, A.C. Kudi, J.D. Beal, P.H. Brooks. 2008. Effect of particle size and micro-organism on lactic acid fermentation of sorghum, In: BSAS (Ed.), *Proceedings of the British Society of Animal Science*, Cambridge University Press, Scarborough, U. K., p. 43.
- Noziere, P., F. Glasser and D. Sauvant. 2011. In vivo production and molar percentages of volatile fatty acids in the rumen : a quantitative review by an empirical approach. *Animal.* 5(3):403-414.
- Ogimoto, K and S. Imal. 1987 . *Atlas of Rumen Microbiology*. Japan Scientific Societies Press, Tokyo.
- Ørskov, E. R. and I. McDonald. 1979. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J. Agric. Sci.* 92: 499-503.
- Reale, A., L. Mannina, P. Tremonte, A.P. Sobolev, M. Succi, E. Sorrentino and R. Coppola. 2004. Phytate degradation by lactic acid bacteria and yeasts during the wholemeal dough fermentation: a <sup>31</sup>P NMR study. *J. Agric. Food Chem* 52:6300-6305.
- Rizzello, C. G., R. Coda, D.S. Macías, D. Pinto, B. Marzani, P. Filannino and M. Gobbetti. 2013. Lactic acid fermentation as a tool to enhance the functional features of *Echinacea* spp. *Microbial Cell Factories.* 12: 44
- Sánchez, B., M. Gueimonde, A.S. Peña and D. Bernardo. 2015. Intestinal microbiota as modulators of the immune system. *J. Immunol. Res.* Article ID. 159094.
- Sari, N.F., R. Ridwan and Y. Widyastuti. 2017. The quality of corn silage product from technopark of Banyumulek Lombok, West Nusa Tenggara. *Buletin Peternakan.* 41(2):156-162.
- Seo, J. K, S.W. Kim, M.H. Kim, S.D. Upadhaya, D.K. Kam and J.K. Ha. 2010. Direct-fed microbials for ruminant animals. *Asian-Australas. J. Anim Sci.* 23:1657-1667.
- Theodorou, M. K., B. A. Williams, M. S. Dhanoa, A. B. McAllan and J. France. 1994. A simple gas production method using a pressure transducer to determine the fermentation kinetics of ruminant feeds. *Anim. Feed Sci. Technol.* 48:185-197.
- Veira, D.M., M. Ivan and P.J. Jui. 1983. Rumen ciliate protozoa: effects on digestion in the stomach of sheep. *J. Dairy Sci.* 66:1015-1022.
- Xin-Xu, H. U. 2015. Effects of complex probiotics fermented feed without antibiotic on growth performance, plasma biochemical parameters, immune function and meat quality in growing-finishing pig. *J. Huazhong Agric. Univer.* 1:72-77.