

EFFECT OF WATER EXTRACT OF PLANTS CONTAINING TANNIN ON IN VITRO METHAGONESIS AND FERMENTATION CHARACTERISTICS OF THE GRASS *Pennisetum purpureophoides*

B. Santoso, E. W. Saragih and B. Tj. Hariadi

Department of Animal Nutrition, Faculty of Animal Science, Fishery and Marine Science,
State University of Papua, Manokwari 98314, Papua Barat - Indonesia
Corresponding E-mail: santosob@lycos.com

Received January 08, 2013; Accepted February 20, 2013

ABSTRAK

Penelitian ini dilaksanakan untuk mengevaluasi pengaruh perbedaan ekstrak tanaman yang mengandung tanin terhadap produksi CH₄, karakteristik fermentasi dan degradasi nutrisi secara *in vitro*. Enam jenis daun tanaman yaitu *Gliricidia sepium*, *Acacia mangium*, *Leucaena leucocephala*, *Desmodium intortum*, *Camellia sinensis*, *Calliandra calothyrsus* dan biji *Areca catechu* diekstraksi menggunakan pelarut air. Perlakuan percobaan terdiri atas *P. purpureophoides* (300±5 mg) diinkubasi tunggal atau ditambah 1,2 mL ekstrak tanaman. Degradasi NDF diukur menggunakan prosedur Tilley and Terry tahap pertama. Hasil penelitian menunjukkan bahwa konsentrasi tanin total pada ekstrak tanaman bervariasi antara 34-95 g/kg BK, terendah pada *D. intortum* dan tertinggi pada *A. mangium*. Produksi CH₄ pada inkubasi 48 jam signifikan lebih rendah ($P < 0,001$) dengan penambahan ekstrak *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* dan *C. calothyrsus* dibandingkan kontrol. Tanin total mempunyai hubungan yang erat dengan produksi CH₄ ($r = -0,79$). Terdapat korelasi yang kuat antara produksi CH₄ dan konsentrasi NDF ($r = 0,61$). Disimpulkan bahwa ekstrak tanaman *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* dan *C. calothyrsus* menggunakan air berpotensi sebagai manipulator untuk menurunkan produksi CH₄ pada ternak ruminansia.

Kata kunci: in vitro, metana, ruminansia, tanin, ekstrak

ABSTRACT

This experiment was conducted to evaluate the effect of extract of plants containing tannin on *in vitro* CH₄ production, fermentation characteristics and nutrient degradability. Six of plant leaves *i.e.* *Gliricidia sepium*, *Acacia mangium*, *Leucaena leucocephala*, *Desmodium intortum*, *Camellia sinensis*, *Calliandra calothyrsus* and seed of *Areca catechu* were extracted by using water. Experimental treatments consisted of *P. purpureophoides* (300±5 mg) incubated alone or added with 1.2 mL of plant extracts. The *in vitro* neutral detergent fibre (NDF) degradability was determined using the first stage technique of Tilley and Terry. The results showed that total tannin concentration of plant extract ranged from 34 to 95 g/kg DM, and was lowest in *D. intortum* and highest in *A. mangium*. Methane production was significantly ($P < 0.001$) lower with addition of *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* and *C. calothyrsus* extracts compared to control. Total tannin had a close relationship with CH₄ production ($r = -0.79$). There was strong correlation between CH₄ production and NDF degradability ($r = 0.61$). It was concluded that water extracts of *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* and *C. calothyrsus* have potential to be used as rumen manipulator in order to reduce CH₄ production in ruminants.

Keywords: in vitro, methane, ruminant, tannin, extract

INTRODUCTION

Methane (CH₄) is produced as a result of anaerobic fermentation of the soluble and structural carbohydrates by methanogens in the rumen of ruminant animals, which is released into

the environment by eructation. The CH₄ emissions from ruminant animals range from 2 to 12% of the gross energy intake (Johnson and Johnson, 1995). Besides, it has been estimated that the world's population of ruminants produce about 15% of total atmospheric CH₄ emissions

(Moss, 1993). This means that CH₄ production from ruminant is not only represents a substantial loss in efficiency of animal production, but also contributes significantly to global warming as the greenhouse gas.

Recently, there is an increasing interest in research activities to evaluate the potential of secondary plant compound as feed additives instead of chemical compounds *i.e.* ionophores and antibiotics as manipulators of rumen fermentation including decrease CH₄ production. As previously stated by Russell and Rychlik (2001), there has been an increased perception that antibiotics and chemical compounds should not be routinely used as feed additives.

Tannin is a phenolic plant secondary compound and is widely distributed through plant kingdom especially legume and browse. Makkar *et al.* (1995) noted that secondary compound tannin is prevalent in many tropical fodder plants. Effect of tannin from some plants such *Acacia mangium*, *Biophytum petersianum* and *Psidium guajava* has been demonstrated as supplement to the tropical grass *Pennisetum purpureum* (Hariadi and Santoso, 2010). Plant extract containing tannin have been shown to decrease CH₄ production (Śliwiński *et al.*, 2002; Sirohi *et al.*, 2009) and ruminal NDF digestibility (McSweeney *et al.*, 2001; Oliveira *et al.*, 2007). Previous study with tropical plants, Jayanegara *et al.* (2011) concluded that total tannin had close relationship with CH₄/digestible OM ($r=-0.74$). However, use of water as a solvent is more applicable and safe for animals than chemical solvents *i.e.* methanol, ethanol or acetone. The objective of this study was to evaluate the effect of water extracts from plant containing tannin on *in vitro* CH₄ production, fermentation characteristics, and nutrient degradability.

MATERIALS AND METHODS

Samples and Extract Preparations

King grass (*P. purpureophoides*) was planted in a 6 m² plot without fertilizer at the Animal Research Station of State University of Papua in Manokwari. Grass was harvested after 50 days, chopped to 5 cm in length and oven-dried at 60°C for 48 h.

The leaves of *G. sepium*, *A. mangium*, *L. leucocephala*, *C. sinensis*, *D. intortum* and *C. calothyrsus* and seed of *A. catechu* were collected from the Manokwari Regency. The collected

samples were then pooled and oven-dried at 60°C for 48 h. A commercial leaf of *C. sinensis* was used in this experiment. Samples of grass, leaves and seed of plants were ground to pass a 1 mm sieve in a Wiley mill.

Plants extract were prepared in water following a modified method of Patra *et al.* (2006). About 5 g of finely ground plants material were weighed into 100 ml beaker glass and added 50 ml of water. Plant materials were boiled for 10 min on a hotplate. The beakers were stoppered and incubated at 39°C on a shaker for 24 h and filtered through 2 layers of cheesecloth. The filtrates were stored at 4°C for further use.

Experimental Design and Treatments

The experiment was arranged in a completely randomized design with eight treatments and three replications. Experimental treatments consisted of *P. purpureophoides* incubated alone as control or added with plant extract at level of 1.2 mL/300 mg of substrate.

In vitro CH₄ Production and Nutrient Degradability Measurements

The *in vitro* gas production method was essentially according to Menke and Steingass (1988) as previously demonstrated by Hariadi and Santoso (2010). Oven-dried samples of about 300±5 mg were weighed into 100 mL glass syringes with pistons lubricated with Vaseline. Rumen liquor was obtained from two ruminally fistulated Holstein Frisian cross-bred cows fed elephant grass twice a day at maintenance level. About 30 mL of rumen liquor-buffer mixtures (1 : 2, v/v) was added into each syringe and then incubated in a water bath at 39°C for 48 h. The volume of gas released from each syringe was recorded before incubation (0 h) and 2, 4, 6, 12, 24 and 48 h of incubation.

Ten milliliter of gas was collected at 24 and 48 h of incubation from the silicon tube at the syringe tip using Terumo syringe and pooled to vacutainer tube for CH₄ analysis. Methane was determined by injection 100 mL of gas into a chromatograph gas. The volume of CH₄ production was calculated by using formula: CH₄ production (ml) = total gas produced (ml) × % CH₄ in the sample.

At the end of the incubation period, about 10 ml of syringe contents were sampled. The pH of medium incubation was recorded using a digital pH meter. Subsequently, 0.2 mL of sub-

samples were pipetted into 1.5 mL micro centrifuge tube containing 1 ml of 25 g/100 mL (w/v) metaphosphoric acid and centrifuged at 9000 x g for 10 min for volatile fatty acids (VFA) determination. Further on 2 mL of sub-samples were added to 2 mL of 20 g/l (w/v) NaCl for NH₃-N (Chaney and Marbach, 1962).

The *in vitro* organic matter (OM) and NDF degradability was determined using the first-stage technique as proposed by Tilley and Terry (1963) as previously demonstrated by Hariadi and Santoso (2010).

Laboratory Analyses

Dried samples were used to determine DM, ash and crude protein (CP) according to procedure of AOAC (1990). Total tannin and condensed tannin contents were assayed using Folin-Ciocalteu and butanol-HCl methods respectively (Makkar, 2003). Concentrations of NDF and acid detergent fibre (ADF) were determined following Van Soest *et al.* (1991).

Statistical Analysis

Data was analyzed using the procedure GLM of SAS (version 6.12, SAS institute, Cary, NC, USA). Duncan's multiple range test was used to separate treatment means. Correlation analysis was done to establish the relationship between variables.

RESULTS AND DISCUSSION

Results

Chemical Composition of Plants

The chemical composition of experimental plants is presented in Table 1. The NDF and ADF content in plants containing tannin ranging from 269 to 596 and 115 to 381 g/kg DM, respectively. The ranking order of plant samples on the basis their total tannin content was *A. mangium* > *C. calothyrsus* < *L. leucocephala* < *A. catechu* > *C. sinensis* > *G. sepium* > *D. intortum*. However, on the basis of their condensed tannin concentration the plant could be ranked as *A. mangium* > *A. catechu* > *L. leucocephala* > *G. sepium* > *C. calothyrsus* > *C. sinensis* > *D. intortum*.

In vitro Gas and CH₄ Productions

Effect of plant extract containing tannin on gas and CH₄ production are given in Table 2. Gas production was significantly (P<0.001) different among treatments at 6, 24 and 48 h of incubation. Addition of plant extracts in grass substrate decreased (P<0.001) gas production at 48 h of incubation compared to control. Methane production in substrate with addition of *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* and *C. calothyrsus* extracts was significantly (P<0.001) lower as compared to control. The CH₄ produced at 48 h of incubation averaged 12.3% of total gas at 48 h of incubation.

Table 1. Chemical Composition (g/kg DM) of *P. purpureophoides* Incubated as Substrate, and Leaves or Seed Plant Containing Tannin

	OM	CP	NDF	ADF	Hemi-celullose	TT	CT
<i>P. purpureophoides</i>	953	148	751	404	347	5	N.D.
<i>G. sepium</i>	916	219	466	315	261	45	25
<i>A. mangium</i>	934	241	477	216	261	95	36
<i>L. leucocephala</i>	920	344	269	115	154	81	28
<i>A. catechu</i>	975	102	496	135	361	77	35
<i>D. intortum</i>	881	207	403	209	194	34	5
<i>C. sinensis</i>	936	197	424	269	155	69	20
<i>C. calothyrsus</i>	944	188	596	381	215	82	21

DM = dry matter; OM = organic matter; CP = crude protein; NDF = neutral detergent fiber; ADF = acid detergent fiber; TT = total tannin; CT = condensed tannin; ND = not detected

Table 2. Gas and CH₄ Productions of *in vitro* Incubation of *P. purpureophoides* Alone or with Addition of Plant Extracts

	Total gas (ml/300 mg of DM)			CH ₄ (ml/300 mg of DM)	CH ₄ /total gas 48 h (ml/l)
	6 h	24 h	48 h		
<i>P. purpureophoides</i>	12.6 ^a	54.1 ^a	71.2 ^a	11.0 ^a	154 ^A
<i>G. sepium</i>	9.3 ^c	43.6 ^{cde}	64.0 ^{cd}	8.4 ^{ab}	131 ^{AB}
<i>A. mangium</i>	7.3 ^{cd}	37.6 ^{ef}	61.3 ^e	6.2 ^b	101 ^B
<i>L. leucocephala</i>	10.3 ^b	50.9 ^{ab}	66.3 ^{de}	7.6 ^b	114 ^B
<i>A. catecú</i>	6.9 ^d	33.3 ^f	69.5 ^e	7.6 ^b	129 ^{AB}
<i>D. intortum</i>	12.1 ^a	49.6 ^{abc}	68.3 ^b	8.3 ^{ab}	121 ^B
<i>C. sinensis</i>	6.8 ^d	40.3 ^{de}	60.5 ^e	7.0 ^b	115 ^B
<i>C. calothyrsus</i>	11.1 ^{ab}	45.3 ^{bcd}	64.8 ^c	7.5 ^b	115 ^B
SEM	0.52	1.54	0.67	0.56	9.15
P	<0.001	<0.001	<0.001	<0.001	0.03

Means in the same column followed by different letters are different (^{A-B}P<0.05; ^{a-f}P<0.01)

Fermentation Characteristics

The pH value, concentrations of NH₃-N and VFA in the medium incubation are present in Table 3. The pH value in medium incubation was not significantly (P>0.05) different among treatments. Concentration of NH₃-N significantly (P<0.05) increased in grass substrate with addition of extract of *L. leucocephala* as compared to control. Addition of extract of *G. sepium*, *A. mangium*, *A. catechu*, *C. sinensis* and *C. calothyrsus* in grass substrate decreased (P<0.001) total VFA concentration. Extracts of *G. sepium*, *A. catechu* and *C. sinensis* increased (P<0.001) proportion of acetate compared to control, whereas *A. mangium* and *C. sinensis* extracts reduced (P<0.001) proportion of propionate. Acetate:propionate ratio was highest in *A. mangium* and lowest in *A. catechu* extracts.

In vitro Nutrient Degradability

In vitro OM and NDF degradability of grass substrate with addition of tannin extracts is presented Table 4. Addition of *L. leucocephala*, *C. sinensis* and *C. calothyrsus* extracts to grass substrate decreased (P<0.001) IVOMD, whereas IVNDFD was decreased (P<0.001) by addition of *A. mangium*, *L. leucocephala*, *A. catechu*, *D.*

intortum, *C. sinensis* and *C. calothyrsus* extracts as compared to control.

Relationships between Tannin Content and Fermentation Variables

The correlations between total tannin, condensed tannin and fermentation variables are in Table 5. There was a negative correlation between total tannin and the gas production at 6, 24 and 48 h of incubation and CH₄ production, indicating that as total tannin concentration increased, the gas and CH₄ production decreased. Total tannin concentration had closer correlation with CH₄ and CH₄/total gas production than condensed tannin concentration. There was negative correlation between total tannin content and both IVOMD and IVNDFD.

Discussion

The total tannin content in *A. mangium* was comparable to that value reported by Jayanegara *et al.* (2011), but condensed tannin content was lower than value of 42 g/kg DM as reported by Jayanegara *et al.* (2011) Concentration of total tannin in *D. intortum* was lower than in *Desmodium intortum* (Getachew *et al.*, 2000). Concentration of condensed tannin in *L.*

Table 3. The pH, Concentrations of NH₃-N and VFA in the Medium Incubation After 48 h of *in vitro* Incubation of *P. purpureophoides* Alone or with Addition of Plant Extracts

	pH	NH ₃ -N (mg/100 ml)	Total VFA (mmol/l)	Acetate (C2) (mol/ 100 mol)	Propionate (C3) (mol/ 100 mol)	Butyrate (mol/100 mol)	C2/C3
<i>P. purpureophoides</i>	6.53	19.5 ^{BC}	139.2 ^a	71.0 ^d	15.9 ^{ab}	13.1 ^a	4.5 ^{cd}
<i>G. sepium</i>	6.53	22.2 ^C	111.5 ^b	75.8 ^{abc}	14.2 ^{bc}	10.0 ^{abc}	5.4 ^{ab}
<i>A. mangium</i>	6.54	21.2 ^{ABC}	88.5 ^d	76.7 ^{ab}	13.2 ^c	10.1 ^{abc}	5.8 ^a
<i>L. leucocephala</i>	6.51	25.4 ^A	137.9 ^a	75.0 ^{abc}	15.6 ^{ab}	9.3 ^{bc}	4.8 ^{bcd}
<i>A. catecú</i>	6.50	17.2 ^{BC}	87.9 ^d	73.8 ^c	17.0 ^a	9.2 ^{bc}	4.3 ^d
<i>D. intortum</i>	6.48	22.6 ^{AB}	128.9 ^a	76.0 ^{abc}	17.0 ^a	7.1 ^c	4.5 ^{cd}
<i>C. sinensis</i>	6.50	19.4 ^{ABC}	97.7 ^{cd}	74.3 ^c	14.4 ^c	11.3 ^{ab}	5.2 ^{abc}
<i>C. calothyrsus</i>	6.53	19.3 ^{ABC}	103.3 ^{bc}	77.2 ^a	15.7 ^{ab}	7.1 ^c	4.9 ^{bcd}
SEM	0.01	1.54	2.69	0.63	0.53	0.73	0.17
P	0.11	0.04	<0.001	<0.001	<0.001	<0.001	<0.001

Means in the same column followed by different letters are different (^{A-C}P<0.05; ^{a-d}P<0.01)

Table 4. The IVOMD and IVNDFD after 48 h of Incubation of *P. purpureophoides* Alone or with Addition of Plant Extracts

	IVOMD (g/kg)	IVNDFD (g/kg)
<i>P. purpureophoides</i>	606 ^a	336 ^a
<i>G. sepium</i>	588 ^{abc}	309 ^a
<i>A. mangium</i>	594 ^{ab}	222 ^{bc}
<i>L. leucocephala</i>	562 ^c	260 ^b
<i>A. catechu</i>	589 ^{abc}	189 ^c
<i>D. intortum</i>	575 ^{abc}	207 ^c
<i>C. sinensis</i>	560 ^c	187 ^c
<i>C. calothyrsus</i>	484 ^d	180 ^c
SEM	0.74	10.1
P	<0.001	<0.001

Means in the same column followed by different letters are different (^{A-B}P<0.05; ^{a-d}P<0.01)

leucocephala and *C. calothyrsus* was lower than values of 76 and 240 g/kg DM, respectively (Tiemann *et al.*, 2008). Different of plant nutrients content in this study as compared to previous study could be due to difference in location of sample source and plants maturity.

Higher gas production with addition of plant extract could be due to the presence of higher of soluble sugar from these extract. In previous studies of Śliwiński *et al.* (2002); Patra *et al.* (2010), increasing gas production by plant extracts that contain phenol or saponin caused by increasing soluble sugar from these plant extract.

No differences in pH values obtained in the present study, consistent with previous studies of Śliwiński *et al.* (2002); Oliviera *et al.* (2007); Animut *et al.* (2008) who found that H was not changed by addition of tannin in both sheep and cattle rumens.

Tannin has beneficial effect on protection on dietary protein in the rumen and subsequently enhancement of amino acid absorption and utilization by the ruminant animal (Waghorn *et al.*, 1994). McSweeney *et al.* (2001) revealed that tannin has ability to bind protein by forming hydrogen bonds between the phenolic sub-units of

Table 5. Coefficient of Correlation (*r*) Between Total Tannin, Condensed Tannin Contents and Gas Production, Fermentation Characteristics, *in vitro* Nutrients Degradability

Variables	Total Tannin	Condensed Tannin
Gas 6 h	-0.63 ^{***}	0.70 ^{***}
Gas 24 h	-0.61 ^{**}	0.60 ^{**}
Gas 48 h	-0.72 ^{**}	0.67 ^{**}
CH ₄	-0.79 ^{***}	-0.37 ^{ns}
CH ₄ /total gas	-0.66 ^{***}	-0.12 ^{ns}
Total VFA	-0.65 ^{***}	-0.53 [*]
pH	0.10 ^{ns}	0.41 ^{ns}
C2	0.56 ^{**}	-0.15 ^{ns}
C3	-0.30 ^{ns}	-0.35 ^{ns}
C4	-0.32 ^{ns}	0.43 [*]
C2/C3	0.41 [*]	0.33 ^{**}
NH ₃ -N	-0.04 ^{ns}	-0.17 ^{ns}
IVOMD	-0.40 [*]	0.21 ^{ns}
IVNDFD	-0.21 ^{ns}	0.17 ^{ns}

ns: not significant ($P > 0.05$); * ($P < 0.05$);

** ($P < 0.01$); *** ($P < 0.001$)

the polymer and the carbonyl groups of peptides of the protein. *In vivo* study, Min *et al.* (2005) suggested that the action of condensed tannin in forages markedly reduced both growth and population of proteolytic bacterial. However, the extent of positive or negative effects of tannins may vary depending on the type and level of tannins in plants and their biological activity (Getachew *et al.*, 2000). Relatively higher concentration ammonia-N in the medium incubation with addition of extract of *G. sepium*, *A. mangium*, *L. leucocephala*, *A. catechu* and *D. intortum* could be due to higher crude protein content in those plants or tannin concentration is lower than minimum concentration needed to produce maximum inhibition of proteolysis.

The inhibition of methanogenesis has long been considered from nutritional aspects, and more recently from the perspectives on greenhouse gas emissions. In the previous studies by Śliwiński *et al.* (2002); Patra *et al.* (2006);

Patra *et al.* (2010); Rodríguez *et al.* (2011) who reported that CH₄ production reduced by addition of plant extract containing tannin. In contrast, Beauchemin *et al.* (2007) described that supplementing a forage-based diet with quebracho tannin extract failed to reduce CH₄ production from growing cattle. The same result has been reported by Oliviera *et al.* (2007) that feeding sorghum silage containing high or low concentrations of tannin did not affect nutrient digestibility and CH₄ production in cattle. However, effect of tannin on methanogenesis depends on the source, type and dose of tannins (Patra *et al.*, 2006). In this study, relative to control, addition 1.2 ml extract of *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* or *C. calothyrsus* decreased CH₄ production by 43.6, 30.9, 30.9, 36.3 and 31.8%, respectively. Decreased CH₄ production in this study, however might be related to decreasing of fibre degradation which is shown by high correlation coefficient value between CH₄ production and NDF degradability ($r = 0.61$; $P < 0.01$) (result not shown). This result was supported by previous study of Estermann *et al.* (2002) that there was a strong relationship between CH₄ production and digestible NDF for cows and calves. Bhatta *et al.* (2009) concluded that tannins suppress methanogenesis by reducing methanogenic populations in the rumen either directly or by reducing the protozoal population, thereby reducing methanogens symbiotically associated with the protozoal population. Tavendale *et al.* (2005) revealed that condensed tannin reduce CH₄ production could be due to indirect effect by reduced hydrogen production as result of reduced feed degradability, and by direct inhibitory effect on methanogens. A high correlation ($r = -0.79$, $P < 0.001$) between total tannin concentration and CH₄ production in this study (Table 5) was agreed with previous studies of Jayanegara *et al.* (2009, 2011); Hariadi and Santoso (2010) that found *r* values -0.60, -0.66 and -0.76, respectively. A similar observation was found by Bhatta *et al.* (2009) that total tannin had close correlation with CH₄ output ($r = -0.97$). In our study, IVNDFD in substrate with addition of *L. leucocephala*, *A. catechu*, *D. intortum*, *C. sinensis* or *C. calothyrsus* extracts was significantly lower than control. This finding agrees with previous study of Bhatta *et al.* (2009) who found that tannin significantly suppressed bacteria population, through a direct effect (Koike and Kobayashi, 2009) and by reducing nutrient availability

(Sallam *et al.*, 2010). In addition, tannin could reduce fibre digestion by complexing with lignocellulose and preventing microbial digestion or by directly inhibiting cellulolytic microorganism or both (McSweeney *et al.*, 2001). In our results, IVOMD decreased ranged from 7.3 to 20.1% when addition of *L. leucocephala*, *C. sinensis* or *C. calothyrsus* extract to *P. purpureophoides* substrate. Similar result has been reported by Patra *et al.* (2006) that addition of extract of plants containing tannin reduced IVDMD and IVOMD by about 7% in comparison to control. Hess *et al.* (2003) also reported that tannins from *C. calothyrsus* may cause significant shifts in rumen microbial populations, especially a reduced total number of cellulolytic bacteria, which could have contributed to lower OM degradation.

CONCLUSION

Plant extracts from *A. mangium*, *L. leucocephala*, *A. catechu*, *C. sinensis* and *C. calothyrsus* have potential to be used as rumen manipulator in order to reduce CH₄ production in ruminants. Among plant extracts, *A. mangium* had the strongest effect on *in vitro* CH₄ production. The mode of action tannin on methanogenesis might be partially attributed to reduce fibre degradability.

ACKNOWLEDGMENT

This study was funded by the Directorate General of Higher Education, Ministry of National Education of Indonesia (Contract No. 262/SP2H/PP/DP2M/V/2009).

REFERENCES

Animut, G., R. Puchala, A.L. Goetsch, A.K. Patra, T. Sahlu, V.H. Varel and J. Wells. 2008. Methane emissions by goats consuming different sources of condensed tannins. *Anim. Feed Sci. Technol.* 144:228-241.

AOAC. 1990. Official Methods of Analysis. 15th ed. Association of Official Analytical Chemist. Arlington. VA. USA.

Beauchemin, K.A., S.M. McGinn, T.F. Martinez and T.A. McAllister. 2007. Use of condensed tannin extract from quebracho trees to reduce methane emissions from cattle. *J. Anim. Sci.* 85:1990-1996.

Bhatta, R., Y. Uyeno, K. Tajima, A. Takenaka, Y.

Yabumoto, I. Nonaka, O. Enishi and M. Kurihara. 2009. Difference in the nature of tannins on *in vitro* ruminal methane and volatile fatty acid production and on methanogenic archaea and protozoal populations. *J. Dairy Sci.* 92:5512-5522.

Chaney, A.L. and E.P. Marbach. 1962. Modified reagents for determination of urea and ammonia. *Clin. Chem.* 8:130-132.

Estermann, B.L., F. Sutter, P.O. Schlegel, D. Erdin, H.R. Wettstein and M. Kreuzer. 2002. Effect of calf age and dam breed on intake, energy expenditure, and excretion of nitrogen, phosphorus, and methane of beef cows with calves. *J. Anim. Sci.* 80:1124-1134.

Getachew, G., H.P.S. Makkar and K. Becker. 2000. Effect of polyethylene glycol on *in vitro* degradability of nitrogen and microbial protein synthesis from tannin-rich browse and herbaceous legumes. *Br. J. Nutr.* 84:73-83.

Hariadi, B.T. and B. Santoso. 2010. Evaluation of tropical plants containing tannin on *in vitro* methanogenesis and fermentation parameters using rumen fluid. *J. Sci. Food Agric.* 90:456-461.

Hess, H.D., L.M. Monsalve, C.E. Lascano, C.E. Carulla, T.E. D'iaz and M. Kreuzer. 2003. Supplementation of a tropical grass diet with forage legumes and *Sapindus saponaria* fruits: effects on *in vitro* ruminal nitrogen turnover and methanogenesis. *Aust. J. Agric. Res.* 54:703-713.

Jayanegara, A., N. Togtokhbayar, H.P.S. Makkar and K. Becker. 2009. Tannins determined by various methods as predictors of methane production reduction potential of plants by an *in vitro* fermentation system. *Anim. Feed Sci. Technol.* 150:230-237.

Jayanegara, A., E. Wina, C.R. Soliva, S. Marquardt, M. Kreuzer and F. Leiber. 2011. Dependence of forage quality and methanogenic potential of tropical plants on their phenolic fractions as determined by principal component analysis. *Anim. Feed Sci. Technol.* 163:231-243.

Johnson, K.A. and D.E. Johnson. 1995. Methane emissions from cattle. *J. Anim. Sci.* 73: 2483-2492.

Koike, S., and Y. Kobayashi. 2009. Fibrolytic rumen bacteria: their ecology and functions. *Asian-Austral J. Anim. Sci.* 22:131-138.

Makkar, H.P.S. 2003. Quantification of Tannins in

- Tree and Shrub Foliage – A Laboratory Manual. Joint FAO/IAEA, Division of Nuclear Techniques in Food and Agriculture. Kluwer Academic Publisher, Dordrecht, The Netherlands.
- Makkar, H.P.S., M. Blümmel and K. Becker. 1995. *In vitro* effects of and interactions between tannins and saponins and fate of tannins in the rumen. *J. Sci. Food Agric.* 69:481-493.
- McSweeney, C.S., B. Palmer, D.M. McNeill and D.O. Krause. 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim. Feed Sci. Technol* 91:83-93.
- Menke, K.H. and H. Steingass. 1998. Estimation of the energetic feed value obtained from chemical analysis and *in vitro* gas production using rumen fluid. *Anim. Res. Develop.* 28:7-55.
- Min, B.R., G.T. Atwood, W.C. McNabb, A.L. Molan and T.N. Barry. 2005. The effect of condensed tannin from *Lotus corniculatus* on the proteolytic activities and growth of rumen bacteria. *Anim. Feed Sci. Technol.* 121:45-48.
- Moss, A.R. 1993. Methane Global Warming and Production by Animals. Chalcombe Publications, Canterbury, p.105.
- Oliveira, S.G., T.T. Berchielli, M.S. Pedreira, O. Primavesi, R. Frighetto and M.A. Lima. 2007. Effect of tannin levels in sorghum silage and concentrate supplementation on apparent digestibility and methane emission in beef cattle. *Anim. Feed Sci. Technol.* 135:236-248.
- Patra, A.K., D.N. Kamra and N. Agarwal. 2010. Effects of extracts of spices on rumen methanogenesis, enzyme activities and fermentation of feeds *in vitro*. *J. Sci. Food Agric.* 90:511-520.
- Patra, A.K., D.N. Kamra and N. Agarwal. 2006. Effect of plant extracts on *in vitro* methanogenesis, enzyme activities and fermentation of feed in rumen liquor of buffalo. *Anim. Feed Sci. Technol.* 128:276-291.
- Rodríguez, R, Britos, A., Rodríguez-Romero, N, Fondevila, M. 2011. Effect of plant extracts from several tanniferous browse legumes on *in vitro* microbial fermentation of the tropical grass *Pennisetum purpureum*. *Anim. Feed Sci. Technol.* 168:188-195.
- Russell, J.B. and J.L. Rychlik. 2001. Factors that alter rumen microbial ecology. *Sci.* 292:1119-1122.
- Sallam, S.M.A., I.C.S. Bueno, P.B. Godoy, E.F. Nozella, D.M.S.S. Vitti and A.L. Abdalla. 2010. Ruminant fermentation of tannins bioactivity of some browses using a semi-automated gas production technique. *Trop. Subtrop. Agroecosyst.* 12:1-10.
- Sirohi, N.K., N. Pandey, N. Goel, B. Singh, M. Mohini, P. Pandey and P.P. Chaudhry. 2009. Microbial activity and methanogenesis as affected by plant secondary metabolites in different plant extracts. *Int. J. Environ. Sci. Eng.* 1:52-58.
- Śliwiński, B.J., M. Kreuzer, H.R. Wettstein and A. Machmüller. 2002. Rumen fermentation and nitrogen balance of lambs fed diets containing plant extracts rich in tannins and saponins, and associated emissions of nitrogen and methane. *Arch. Anim. Nutr.* 56:379-392.
- Tavendale, M.H., M.P. Meagher, D. Pacheco, N. Walker, G.T. Atwood and S. Sivakumaran. 2005. Methane production from *in vitro* incubations with *Lotus pedunculatus* and *Medicago sativa*, and effects of extractable condensed tannin fractions on methanogenesis. *Anim. Feed Sci. Technol.* 123-124:403-419.
- Tiemann, T.T., P. Avila, G. Ramirez, C.E. Lascano, M. Kreuzer and H.D. Hess. 2008. *In vitro* ruminal fermentation of tanniniferous tropical plants: Plant-specific tannin effects and counteracting efficiency of PEG. *Anim. Feed Sci. Technol.* 146:222-241.
- Tilley, J.M. and R.A. Terry. 1963. A two stage technique for the *in vitro* digestion of forage crops. *J. Brit. Grassl. Soc.* 18:104-111.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74:3583-3597.
- Waghorn, G.C., I.D. Shelton, W.C. McNabb and S.N. McCutcheon. 1994. Effect of condensed tannins in *Lotus pedunculatus* on its nutritive value for sheep. 2. Nitrogenous aspects. *J. Agric. Sci.* 123:09-119.