Antioxidant Activity of *Daemonorops draco* Resin

Sri Purwanti^a, Wulan Tri Wahyuni^{a,b}, Irmanida Batubara^{a,b,*}

^a Department of Chemistry, Faculty of Mathematics and Natural Sciences, IPB University, Kampus IPB Darmaga, Bogor, Indonesia

^b Tropical Biopharmaca Research Center, IPB University, Kampus IPB Taman Kencana. Jalan Taman Kencana No 3, Bogor, Indonesia.

* Corresponding author: ime@apps.ipb.ac.id

<https://doi.org/10.14710/jksa.22.5.179-183>

Article Info

Article history:

Received: 6 May 2019

Revised: 5 August 2019

Accepted: 13 September 2019

Online: 30 September 2019

Keywords:

Antioxidant;
chromatography;
Daemonorops draco;
fractionation; resin

Abstract

Jernang resin is secretion of jernang rattan (*Daemonorops draco*, *Arecaceae* family) fruits which is endemic in Southeast Asia. This resin has various biological activities and empirically used as wound healing, headache medicines, and fever remedies by Anak Dalam ethnic group from Jambi. This study was performed to evaluate the antioxidant activity of nonpolar fraction of *D. draco* resin which collected from Jambi Province, Sumatera, Indonesia. Resin was extracted with n-hexane, ethyl acetate, and methanol respectively. The antioxidant properties of the extracts were then evaluated using 1,1-diphenyl-2-picryl-hydrazyl radical scavenging assay. The most active extract was further fractionated using n-hexane and methanol and separated using column chromatography and preparative thin layer chromatography. Separation of the extract was conducted through antioxidant assay-guided fractionation. Characterization of the active fraction was carried out by infrared spectroscopy. The result shows that ethyl acetate extract provides higher antioxidant activity ($IC_{50} = 27.61 \mu\text{g/mL}$) compare to methanol and n-hexane extracts. N-hexane fraction of ethyl acetate extract used for further separation using column and preparative thin layer chromatography due to its antioxidant activity. Separation using column chromatography resulting in 9 fractions (F.1–9). Fraction F.5 provide high antioxidant activity ($IC_{50} = 17.27 \mu\text{g/mL}$) and further separated using preparative thin layer chromatography resulting two fractions with lower antioxidant activity F.5.1 ($IC_{50} = 85.18 \mu\text{g/mL}$) and F.5.2 ($IC_{50} = 34.94 \mu\text{g/mL}$). Characterization of fraction F.5.2 using infrared spectroscopy showed that component in fraction F.5.2 contains NH-substituted benzene.

1. Introduction

Most of world's health problems are caused by the presence of free radicals. Uncontrolled free radicals development causes damages in tissues and biomolecules extensively, which lead to pathological disorders, such as aging [1], cancer [2], cardiovascular [3] and alzheimer disease [4]. Therefore, natural antioxidant is required as a free radicals scavenger and able to prevent the human body from any kind of oxidative damages. Natural resins from natural sources, which can be described as a sticky yet water-insoluble materials produced from damaged or infected plants, are acknowledged have a potential antimicrobial, antibacterial antibiofilm [5] and antioxidant activities [6, 7].

One of natural resin was produced in *Daemonorops draco* plant from Indonesia and well known as “Dragon’s blood” resin. It was a bright red resin, which has been continuously used as dyes, varnish, incense and medicine [8]. Dragon’s blood resin generally obtained from two main genus: *Daemonorops* (Palmae) found in Indonesia, India and Malaysia, and *Dracaena* (Liliaceae) grown in Hainan province of China, Vietnam, and Cambodia. Indonesian *D. draco* is considered as the most common species available for traditional medication, for instance, it was used for healing a wound, headache, and even fever by Anak Dalam ethnic group from Jambi, Indonesia [9]. Furthermore, *D. draco* is commercially produced by Meer corporation has been reported for having antibacterial

activity [10]. *D. draco* from Jambi Province, Indonesia is also reported has antibacterial activity [11]. Moreover, it has been reported that *D. draco* resin also has antiviral [8], anticancer [12], and anti-inflammatory activity [13]. Activity of the resin as antioxidant need to be explored. This study was conducted to evaluate antioxidant activity of *D. draco* resin collected from Jambi Province, Sumatera, Indonesia. Since some polar constituent from *D. draco* had been isolated [10, 14–16], this study also needs to evaluate nonpolar constituent of *D. draco* resin.

2. Methods

2.1. Extraction and Fractionation

D. draco resin was collected from Sarolangun Jambi, Sumatera, Indonesia. Approximately 150 g dried resin was extracted by 300 mL of solvents with a gradual increase of polarity in each solvent. Firstly, *n*-hexane was used, then ethyl acetate, and finally methanol as polar solvent. Ethyl acetate extract (1 g) was then fractionated by solvent-solvent extraction (methanol:*n*-hexane = 30 mL:30 mL) for 4 times, then the *n*-hexane fraction was collected. A total of 1.45 gram of *n*-hexane fraction was then separated by column chromatography with silica gel as the stationary phase and chloroform:*n*-hexane (9:1) as the mobile phase. From this separation step, 9 fraction was collected (number 1–9). Fraction 5 was again separated by preparative thin layer chromatography using GF₂₅₄ silica glass plate and dichloromethane:*n*-hexane:methanol (9:1:0.1) solvent system, which resulting fraction 5.1 and fraction 5.2.

2.2. Antioxidant Assay

Antioxidant activity was carried out by 1,1-diphenyl-2-picryl hydrazyl (DPPH) radical scavenging assay. Each 100 μ L of sample and DPPH solution (11.8 mg in 100 mL ethanol) were added into 96-well of microplate. The absorbance was then measured at 514 nm after 30 minutes of incubation. Finally, the inhibition activity of sample was calculated and determined [17]. In this assay, ascorbic acid was used as a positive control and ethanol was used as blank.

2.3. TLC Bioautography

Briefly, 10 μ L of each extract (2 g in ethanol) was applied to the TLC Silica GF₂₅₄. Chromatography method was conducted using chloroform: methanol: water =9:1:0.1 as mobile phase. The TLC plate was then sprayed by DPPH solution following chromatographic elution. After 30 min, the yellow spots from reduced DPPH were clearly observed against the purple background.

2.4. Characteritation of Fraction

Characterization was conducted by infrared spectroscopy. 2 mg of sample was measured using attenuated total reflectance (ATR) FTIR spectrophotometer.

2.5. Statistical Analysis

Statistical analysis was performed as means $\pm$ SD from three independent replicates. One ways analysis of variance (ANOVA) was applied for comparison of the mean values with 95% confidence levels.

3. Results and Discussion

3.1. Extract and Fraction of *Daemonorops draco*

The extracts yields of *D. draco* resin using *n*-hexane, ethyl acetate, and methanol as extraction solvent were 0.22%, 73% and 6.38%, respectively (Table 1). The previous study reported by Hao *et.al.* [14] showed that the extraction of *D. draco* using chloroform give 87.33% yield. The result indicated that *D. draco* resin contains mainly semipolar constituents followed by the polar and then the nonpolar constituent. Ethyl acetate is a semipolar solvent with the index polarity of 4.4, likewise the chloroform. Those semipolar solvent successfully extracted constituents from *D. draco* resin.

Some of the semipolar and polar constituent from *D. draco* resin had been isolated and some of them are known to be the flavonoid compounds. Dracorhodin and dracorubin were isolated from chloroform-methanol extract [10], dracoflavan B1, B2, C1, C2, D1 and D2 were isolated from ethyl acetate extract [15], daemonorol group (A–F) were isolated from acetone extract [16] and dimethoxyflavan group were isolated from chloroform extract [14]. Since the compounds were mainly isolated from semipolar and polar extracts, it was predicted that the constituents in ethyl acetate and methanol extract should provide antioxidant activity.

Flavonoids play an important role in plant growth and development, and as the defense system of plants against the microorganisms and pests. The best-described property of almost every group of flavonoids is their ability to act as antioxidants. The flavonoids seem to be the most powerful compound to protect the body against reactive oxygen species [18].

The ethyl acetate extract provided the highest maceration yield and was fractionated by solvent-solvent extraction with methanol and *n*-hexane. This separation yielded 24.46% of *n*-hexane fraction. The nonpolar fraction was further fractionated by column chromatography with the eluent system of chloroform:*n*-hexane (9:1) and obtained 9 fractions (1–9), where the fraction number 6 was provided the greatest yield. Furthermore, fraction number 4 showed better antioxidant activity followed by fraction number 5, but the difference was not significant ($p > 0.05$). TLC profile (Figure 1) showed that fraction 4 has more components than those in fraction number 5. In order to obtain greater weight and higher purity of active component, fraction number 5 was then separated with the preparative TLC using dichloromethane: *n*-hexane: methanol (9:1:0.1) eluent system.

Figure 1. TLC chromatogram of fraction 4 and 5 at UV 254 nm

Table 1. Weight, % yield, and antioxidant activity of extracts and fraction of *D. draco*

Sample	Weight (g)	% Yield	IC ₅₀ (μg/mL)
<i>n</i> -hexane extract	0.2950	0.22	63.06±2.28
Ethyl acetate extract	100.2595	73.31	27.61±0.40
Methanol extract	8.7293	6.38	63.86±5.40
Non polar fraction	1.4510	24.46	-
F.1*	0.0105	0.72	5697.49±500.94
F.2*	0.0033	0.23	651.86±33.21
F.3*	0.0905	6.24	56.93±9.55
F.4*	0.1124	7.75	14.19±2.04
F.5*	0.1122	7.73	17.27±4.31
F.6*	0.1302	8.97	78.34±7.91
F.7*	0.0289	1.99	94.20±18.81
F.8*	0.0265	1.83	21.20±1.15
F.9*	0.0179	1.23	81.41±8.08
F.5.1**	0.0036	9.00	85.18±0.52
F.5.2**	0.0057	14.25	34.94±2.88
Ascorbic acid			5.12±0.15

* Separated by column chromatography

** Separated by preparative TLC

(-) not tested

3.2. Antioxidant Activity of Extracts and Fractions of *Daemonorops draco*

The antioxidant activities of *D. draco* extracts were evaluated by their ability as the DPPH radicals scavenging through the hydrogen donating mechanism, which could reduce the stable violet of DPPH radical to the yellow DPPH-H. The result in Table 1 showed that the ethyl acetate extract of *D. draco* exhibited higher antioxidant activity (IC₅₀ 27.61 μg/mL) than the activity of *n*-hexane and methanol extract.

The antioxidant activities of *D. draco* showed at IC₅₀ values range of 14.19–569.49 μg/mL (Table 1). Other

species showed IC₅₀ such *Dracaena cinnabari* Balf f. resin (IC₅₀ 94.2–135.7 μg/mL) [7], *Pinus oocarpa* (154.50 μg/mL), *Pinus insularis* (99.328 μg/mL), *Pinus merkusii* (60.20 μg/mL) and *Agathis loranthifolia* (245.99–438.55 μg/mL) [6], it showed that *D. draco* extracts has a higher activity. However, it still has a lower activity than ascorbic acid (IC₅₀ 5.12±0.15 μg/mL).

A qualitative test of the extracts was performed to identify the compounds which contribute in the antioxidant activity by using thin layer chromatography method followed by bioautography method using DPPH reagent, where active compound was identified as a pale-yellow spot on a violet background. Component with the highest DPPH reducing activity shown in Figure 2C, was a component with the retardation factor (R_f) of 0.71 (indicated with red arrow) in *n*-hexane, ethyl acetate, and methanol extract, which was then become the targetted component in this research. Fractionation was then conducted by solvent-solvent extraction method (methanol:*n*-hexane) in order to increase the component yield. The *n*-hexane fraction was then further fractionated by column chromatography.

Figure 2. TLC chromatogram of *D. draco* resin extract monitored at 254 nm (A), 366 nm (B), visible light after sprayed by DPPH, antioxidant bioautography (C), *n*-hexane extract (H), ethyl acetate extract (E), methanol extract (M), the mobile phase was chloroform: methanol: water = 9:1:0.1, while the stationary phase was silica GF₂₅₄

TLC-bioautography showed that the component with R_f of 0.71 possesses antioxidant activity. Solvent-solvent extraction was conducted to separate the nonpolar from the polar component. The nonpolar component was then fractionated by column chromatography. According to Table 1, Fraction 4 from column chromatography showed the highest antioxidant activity but not significantly different with Fraction 5 (*p*>0.05). From the TLC profile, it showed that there are more components in Fraction 4 than the components in the Fraction 5, therefore in order to obtain the active component with a greater yield, the separation of Fraction 5 with less components was carried out by preparative TLC. Fraction 5 (IC₅₀: 17.27 ± 4.31 μg/mL) has a better antioxidant activity than F.5.1 and F.5.2. It is estimated that there is a synergistic effect between the

compounds in fraction number 5 in their mechanism as an antioxidant.

Fraction 5.2, which has an active component as antioxidant, has R_f 0.30 in TLC with F_{254} silica gel as the stationary phase and dichloromethane:*n*-hexane:methanol (9:1:0.1) as the mobile phase. Fraction 5.2 is a brownish-white amorphous solid which has a melting point at 160–162 °C. The results of the analysis of F.5.2 with FTIR spectrophotometer (Figure 3) shows the absorption at wave number of 3445 cm^{-1} as one single peak which suggested that there is N-H band of the secondary amine. The absorption at 1600 cm^{-1} indicates the presence of aromatic groups. The absorption that appears at 2843 cm^{-1} shows the vibration of the C-H strain. Vibration of C=C ring indicated at 1500 cm^{-1} and 1458 cm^{-1} . This result indicates that F.5.2 contains amine ($-\text{NH}_2$) substituted aromatic group.

Figure 3. IR spectrum of F.5.2

The UV spectrum of F.5.2 showed the peak in the wavenumber of 230 nm indicating a transition of $\pi \rightarrow \pi^*$ and the peak at 280 nm indicating a forbidden transition of $\pi \rightarrow \pi^*$. Therefore, it roughly indicates that F.5.2 contains NH-substituted benzene. A further isolation and characterization could be suggested for future work since this compound was potential as antioxidant agent.

4. Conclusions

Ethyl acetate extract of *D. draco* resin provide higher antioxidant activity compare to methanol and *n*-hexane extracts. Fractionation of *n*-hexane fraction of ethyl acetate extract using column chromatography resulting fraction F.4 and F.5 which provide high antioxidant activity with IC_{50} value 14.19 $\mu\text{g/mL}$ and 17.27 $\mu\text{g/mL}$, respectively. Separation of F.5 using preparative thin layer chromatography resulting two fractions (F.5.1 and F.5.2) with lower antioxidant activity compare to F.5. Characterization of fraction F.5.2 using infrared spectroscopy showed that component in fraction F.5.2 contains NH-substituted benzene.

Acknowledgment

This research was financially supported by Ministry of Research, Technology and Higher Education in accordance with the Letter of Appointment

Implementation of Research Program Number: 079/SP2H/LT/DRPM/ II/2016 17 February 2016

References

- [1] Dong-Hoon Hyun, Joe O. Hernandez, Mark P. Mattson and Rafael de Cabo, The plasma membrane redox system in aging, *Ageing Research Reviews*, 5, 2, (2006) 209–220
<https://doi.org/10.1016/j.arr.2006.03.005>
- [2] Vuokko L. Kinnula and James D. Crapo, Superoxide dismutases in malignant cells and human tumors, *Free Radical Biology and Medicine*, 36, 6, (2004) 718–744
<https://doi.org/10.1016/j.freeradbiomed.2003.12.010>
- [3] Uma Singh and Ishwarlal Jialal, Oxidative stress and atherosclerosis, *Pathophysiology*, 13, 3, (2006) 129–142 <https://doi.org/10.1016/j.pathophys.2006.05.002>
- [4] Mark A. Smith, Catherine A. Rottkamp, Akihiko Nunomura, Arun K. Raina and George Perry, Oxidative stress in Alzheimer's disease, *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*, 1502, 1, (2000) 139–144
[https://doi.org/10.1016/S0925-4439\(00\)00040-5](https://doi.org/10.1016/S0925-4439(00)00040-5)
- [5] Termentzi Aikaterini, Fokialakis Nikolas and Skaltsounis Alexios Leandros, Natural Resins and Bioactive Natural Products thereof as Potential Antimicrobial Agents, *Current Pharmaceutical Design*, 17, 13, (2011) 1267–1290
<http://dx.doi.org/10.2174/138161211795703807>
- [6] Mardho Tillah, Irmanida Batubara and Rita Kartika Sari, Antimicrobial and Antioxidant Activities of Resins and Essential Oil From Pine (*Pinus merkusii*, *Pinus ocarpa*, *Pinus insularis*) and *Agathis* (*Agathis loranthifolia*), *Biosaintifika: Journal of Biology & Biology Education*, 9, 1, (2017) 134–139
<https://doi.org/10.15294/biosaintifika.v9i1.8371>
- [7] Deepika Gupta and Rajinder K. Gupta, Bioprotective properties of Dragon's blood resin: In vitro evaluation of antioxidant activity and antimicrobial activity, *BMC Complementary and Alternative Medicine*, 11, 1, (2011) 1–9
<https://doi.org/10.1186/1472-6882-11-13>
- [8] Deepika Gupta, Bruce Bleakley and Rajinder K. Gupta, Dragon's blood: Botany, chemistry and therapeutic uses, *Journal of Ethnopharmacology*, 115, 3, (2008) 361–380 <https://doi.org/10.1016/j.jep.2007.10.018>
- [9] Rana Rio Andhika, Bambang Hariyadi and Fachruddin Saudagar, Etnobotani Penghasil Getah oleh Suku Anak Dalam di Taman Nasional Bukit Duabelas Kabupaten Sarolangun, Jambi, *Jurnal Ilmu Pertanian Indonesia*, 20, 1, (2015) 33–38
- [10] G. S. R. Rao, M. A. Gerhart, R. T. Lee, L. A. Mitscher and S. Drake, Antimicrobial Agents From Higher Plants. Dragon's Blood Resin, *Journal of Natural Products*, 45, 5, (1982) 646–648
<http://doi.org/10.1021/np50023a024>
- [11] Wulan Tri Wahyuni, Sri Purwanti and Irmanida Batubara, Antibacterial and Antibiofilm Activity of *Daemonorops draco* Resin, *Biosaintifika: Journal of Biology & Biology Education*, 10, 1, (2018) 138–144
<https://doi.org/10.15294/biosaintifika.v10i1.13554>

- [12] Jing-hua Yu, Gui-bin Zheng, Chun-yu Liu, Li-ying Zhang, Hong-mei Gao, Ya-hong Zhang, Chun-yan Dai, Lin Huang, Xian-ying Meng, Wen-yan Zhang and Xiao-fang Yu, Dracorhodin Perchlorate Induced Human Breast Cancer MCF-7 Apoptosis through Mitochondrial Pathways, *International Journal of Medical Science*, 10, 9, (2013) 1149–1156
<http://doi.org/10.7150/ijms.6275>
- [13] Ping-Chung Kuo, Hsin-Yi Hung, Tsong-Long Hwang, Wen-Ke Du, Hsiang-Chih Ku, E. Jian Lee, Shih-Huang Tai, Fu-An Chen and Tian-Shung Wu, Anti-inflammatory Flavan-3-ol-dihydroretro-chalcones from *Daemonorops draco*, *Journal of Natural Products*, 80, 4, (2017) 783–789
<http://doi.org/10.1021/acs.jnatprod.7b00039>
- [14] Qian Hao, Yoshinori Saito, Yosuke Matsuo, Hai-Zhou Li and Tanaka Takashi, Three new flavans in dragon's blood from *Daemonorops draco*, *Natural Product Research*, 29, 15, (2015) 1419–1425
<http://doi.org/10.1080/14786419.2014.1003137>
- [15] Alberto Arnone, Gianluca Nasini, Orso Vajna de Pava and Lucio Merlini, Constituents of Dragon's Blood. 5. Dracoflavans B1, B2, C1, C2, D1, and D2, New A-Type Deoxyproanthocyanidins, *Journal of Natural Products*, 60, 10, (1997) 971–975
<http://doi.org/10.1021/np9702188>
- [16] Ken-ichi Nakashima, Naohito Abe, Fumiko Kamiya, Tetsuro Ito, Masayoshi Oyama and Munekazu Iinuma, Novel Flavonoids in Dragon's Blood of *Daemonorops draco*, *Helvetica Chimica Acta*, 92, 10, (2009) 1999–2008
<http://doi.org/10.1002/hlca.200900086>
- [17] Irmanida Batubara, Tohru Mitsunaga and Hideo Ohashi, Screening antiacne potency of Indonesian medicinal plants: antibacterial, lipase inhibition, and antioxidant activities, *Journal of Wood Science*, 55, 3, (2009) 230–235
<http://doi.org/10.1007/s10086-008-1021-1>
- [18] Bishnu Prasad Pandey, Rupak Thapa and Anil Upreti, Chemical composition, antioxidant and antibacterial activities of essential oil and methanol extract of *Artemisia vulgaris* and *Gaultheria fragrantissima* collected from Nepal, *Asian Pacific Journal of Tropical Medicine*, 10, 10, (2017) 952–959
<https://doi.org/10.1016/j.apjtm.2017.09.005>