Microwave-assisted synthesis of 1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one and its activities as an antioxidant, sunscreen, and antibacterial

Ihsan Ikhtiarudin^{a,*}, Nesa Agistia^a, Neni Frimayanti^a, Tria Harlianti^a, Jasril^b

^a Pharmacy Study Program, Sekolah Tinggi Ilmu Farmasi Riau, Jl. Kamboja, Kel. Simpang Baru, Kec. Tampan, Panam, Pekanbaru, 28293, Indonesia

^b Chemistry Department, FMIPA Universitas Riau, Jl. H.R. Subrantas Km. 12,5 Panam, Pekanbaru, 28293, Indonesia

* Corresponding author: ihsanikhtiarudin@stifar-riau.ac.id

<https://doi.org/10.14710/jksa.23.2.51-60>

Article Info

Article history:

Received: 9th December 2019

Revised: 10th February 2020

Accepted: 24th February 2020

Online: 29th February 2020

Keywords:

chalcone analog;
antioxidant; DPPH,
sunscreen; disk diffusion

Abstract

Chalcone analogs have been reported to have a variety of exciting biological activities, such as anticancer, anti-inflammatory, antioxidant, photoprotective, antibacterial, and antidiabetic activities. Therefore, analogs of these compounds have been widely synthesized as intermediate compounds or target molecules in the discovery of bioactive compounds to be applied in the pharmaceutical field. The purpose of this study is to synthesize chalcone analog, (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one under microwave irradiation and explore some of the biological activities of this compound, including testing the antioxidant activity by 1,1-diphenyl-2-picrylhydrazyl (DPPH) method, *in vitro* sunscreen activity by microplate method, and antibacterial activity by disk diffusion method. DPPH test results indicate that the compound has weak antioxidant activity, with an IC₅₀ value of 300.43 µg/mL. However, the compound showed excellent potential as a UV B and UV A filter at a concentration of 150 µg/mL with a %Te value of 0.73±0.10% (sunblock), %Tp value of 0.07±0.00% (sunblock), SPF value of 21.10±1.46 (ultra-protection) and potentially better than benzophenone-3 as a standard sunscreen. Then, disk diffusion testing showed that the compound had weak antibacterial activity against *Staphylococcus aureus* and did not show antibacterial activity against *Escherichia coli* at test concentrations of 30, 60, and 120 µg/disk.

1. Introduction

Chalcone (1,3-diphenylprop-2-en-1-one) is a precursor of flavonoid compounds that can be found in several types of plants. This compound is known to have a variety of exciting biological activities. For example, Broussonchalcone A isolated from the plant *Broussonetia papyrifera* Vent is reported to have free radical scavenging activity stronger than α-tocopherol as a comparative antioxidant [1]. Some other natural chalcone analogs are also reported to have antibacterial activity; for example, the lichochalcone E. This compound is reported can inhibit the growth of several strains of *Staphylococcus*

aureus with a minimum inhibitory concentration of 1-4 µg/mL [2]. Some hydroxy-substituted synthetic chalcone analogs are also reported to have inhibitory properties against *Escherichia coli* and *Staphylococcus aureus* bacteria [3]. Then, some nitro-substituted [4] and hydroxy-substituted [5] synthetic chalcone analogs have also been patented regarding their potential use as active ingredients in some cosmetic products, because they possessed good sunscreen activity. Other studies have also shown that hydroxy and methoxy substituted chalcone analogs, such as (E)-1-(2-hydroxyphenyl)-3-(4-methoxy phenyl)prop-2-en-1-one shows the

maxima absorption close to the dibenzoyl methane as one of the active ingredients of sunscreen [6]. Then, the latest research shows that the analog chalcone (*E*)-1-(3-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one has the potential to be developed as an active ingredient in sunblock preparations at a concentration of 200 µg/mL with an SPF value of 30.61 ± 0.38 [7].

Figure 1. Structure of chalcone

Besides showing a variety of potential biological activities, chalcone analogs have also been used extensively as intermediates in the synthesis of their derivatives [8]. Various chalcone-derived compounds, such as flavonols [9] and pyrazolines [10], have been reported to have a variety of biological activity potentials. The diverse biological activities of chalcone analog compounds and their derivatives have made many researchers interested in synthesizing and exploring the potential use of these compounds in the context of discovering bioactive compounds as drug candidates.

Chalcone analog compounds can be synthesized by several conventional methods, such as the grinding method [11], stirring with magnetic stirrer [8], and reflux [12]. In some cases, this conventional method has several disadvantages, such as low reaction selectivity [13] and requires a relatively long reaction time [9]. On the other hand, synthesis by the microwave irradiation method has been widely reported can increase reaction selectivity and also shorten the reaction time [14].

Synthesis using conventional methods and anticancer potential of chalcone analog (*E*)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one has been reported by several previous researchers [15, 16, 17, 18]. However, the application of microwave irradiation to synthesize this compound has never been reported. Therefore, in this study, we are interested in applying the microwave irradiation method to synthesize this chalcone analog, then evaluate some other biological activities, including antioxidants, sunscreen, and antibacterial activities *in vitro* to explore the potential use of this compound widely. Because, in many cases, the fact is found that a chalcone analog possessed various biological activities. For example, xanthohumol, besides showing antidiabetic activity, this compound also has anti-cancer, antioxidant, and anti-inflammatory activity. In addition, isobavachalcone, besides showing anticancer activity, it also shows antibacterial and antifungal activity [19].

2. Methodology

2.1. Equipment and Materials

The equipment used for synthesis is the Samsung microwave (ME109F). The progress of the synthesis reaction was observed by TLC analysis using GF₂₅₄ silica gel plates (Merck). The spots on the TLC plates were observed under UV lamps 254 and 366 nm (CamagTM). HPLC analysis was carried out with UFLC Prominence-Shimadzu LC Solution, using a Shim-pack VP-ODS column (250 x 4.6 mm), a mixture of methanol and water used as mobile phase in a gradient elution system, with a flow rate of 0.75 mL/min, detected by a UV detector SPD 20AD and analysis time for 25 minutes. The melting point of the synthesized compound was measured by Stuart Melting Point (SMP-11). The UV spectrum was measured by UV-Vis Spectrophotometer (Genesys 10S UV-VIS v4.002 2L9N175013). The FT-IR spectrum was measured by FT-IR Spectrophotometer (Shimadzu, IR Prestige-21). The ¹H NMR spectrum was measured by the NMR spectrometer (Agilent, 500 MHz). The high-resolution mass spectrum is measured by premier XE mode positive LCT water. Antioxidant activity was evaluated with a 96-wells microplate reader (Berthold), and the sunscreen activity was evaluated with a 96-wells microplate reader (Epoch Biotech).

The materials used are 4'-hydroxy acetophenone (Merck), 4-methoxy benzaldehyde (Merck), potassium hydroxide (Merck), hydrochloric acid (Merck), distilled water (Bratachem), universal indicators (Merck), filter papers (Whatman 42), 1,1-diphenyl-2-picrylhydrazyl (Sigma-Aldrich), ascorbic acid (Merck), benzophenone-3 (Cosmetic Grade), Nutrient Agar (Merck), physiological NaCl (Otsuka), chloramphenicol 30 µg/disk antibiotic (Oxoid), and some organic solvents such as *n*-hexane (Merck), ethyl acetate (Merck), chloroform (Merck), dichloromethane (Merck), methanol (Merck), absolute ethanol and dimethylsulfoxide (Merck).

2.2. Synthesis of Chalcone Analog

Synthesis of chalcone analog (*E*)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one as can be seen in Figure 2, was carried out by modifying the previous synthesis method [15, 16, 17, 18]. As much as 20 mmol 4-hydroxy acetophenone was dissolved in 20 mL absolute ethanol in an erlenmeyer, then added with 50 mL of 6N KOH solution and homogenized. 20 mmol of 4-methoxy benzaldehyde was added to the solution, and then the reactant mixture was irradiated in the microwave using 180 W power. The progress of the reaction was observed every 2 minutes by TLC analysis. After the reaction was complete, the reaction mixture was neutralized by adding 3N hydrochloric acid solution to form a precipitate. The precipitate was filtered with a Buchner funnel, and the solid was dried in a desiccator. The crude product solid obtained was tested by TLC to determine its purity and purified by recrystallization technique in hot ethanol.

Then, the purity and physical properties of the recrystallized product are determined by the TLC test, melting point measurement, and HPLC analysis. The structure of the pure compound was determined through UV, FT-IR, ¹H NMR, and HRMS spectroscopic analysis.

Figure 2. The synthesis route of chalcone analog

(*E*)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one. Yellow solids. Yield 89%. Melting point 184–185 °C. TLC chromatogram: *R_f* 0.45 (*n*-hexane:ethyl acetate = 6:4). HPLC Chromatogram: *t_R* = 11.382 minutes. UV Spectrum (MeOH) (λ_{max} , nm): 203, 235 and 345. FT-IR spectrum (KBr) (ν , cm^{-1}): 3161, 3089, 2975, 2941, 2908, 1643, 1601, 1508, 1466, 1466, 1466 and 1037. The ¹H NMR spectrum (acetone-*d*₆, 500 MHz) (δ , ppm): 9.23 (*br-s*, ¹H, Ar-4-OH); 8.09 (*d*, 2H, Ar-2', 6'-H, *J* = 8.5 Hz); 7.79 (*d*, 2H, Ar-2,6-H, *J* = 9.0 Hz); 7.74 (*s*, 2H, H_a, H_b); 7.02 (*d*, 2H, Ar-3,5-H, *J* = 9.0 Hz); 6.98 (*d*, 2H, Ar-3', 5'-H, *J* = 8.5 Hz); 3.87 (*s*, 3H, Ar-4-OCH₃). ¹H NMR spectrum based on literature [20] (polysol, 300 MHz) (δ , ppm): 10.10 (*s*, ¹H, Ar-4-OH), 8.00 (*d*, 2H), 7.69 (*d*, ¹H, H_a), 7.66 (*d*, 2H), 7.58 (*d*, ¹H, H_a), 6.93 (*d*, 4H), 3.80 (*s*, 3H). ¹H NMR spectrum based on literature [21] (CDCl₃, 200 MHz) (δ , ppm): 7.99 (*d*, 2H, *J* = 8.6 Hz), 7.78 (*d*, ¹H, H_a, *J* = 15.6 Hz), 7.60 (*d*, 2H, *J* = 8.6 Hz), 7.41 (*d*, ¹H, H_a, *J* = 15.6 Hz), 6.93 (*d*, 4H, *J* = 7.2 Hz), 5.85 (*s*, ¹H, Ar-4-OH), 3.86 (*s*, 3H, Ar-4-OCH₃). Mass spectra (HRMS, ES⁺): mass was calculated as C₁₆H₁₅O₃ [*M* + *H*]⁺: 255.1021 *m/z* and found at *m/z* = 255.1010 *m/z*.

2.3. Measurement of antioxidant activity by DPPH method

The *in vitro* antioxidant activity assay was conducted by modifying the previous procedure [22]. A sample solution (1000 µg/mL) and DPPH solution (80 µg/mL) were prepared in a methanol solvent. A total of 100 µL of 1000 µg/mL sample solution was pipetted to row A wells on 96-wells microplate. Then, the sample solution in line A was diluted by the two-fold serial dilution method by filling wells in line B–G with 50 µL of methanol and followed by transferring 50 µL of the solution from line A to line B. Then as much as 50 µL of the solution from line B was pipetted to line C, and so on up to line F. Furthermore, a total of 50 µL of the solution from row F was pipetted and discarded, so that several sample solutions were obtained with a concentrations of 1000 µg/mL (line A), 500 µg/mL (line B), 250 µg/mL (line C), 125 µg/mL (line D), 62.5 µg/mL (line E), and 31.25 µg/mL (line F). Ascorbic acid solutions as standard antioxidant were prepared in the same way, but with a variety of

concentrations of 100; 50; 25; 12.5; 6.25; and 3.125 µg/mL. Then, as much as 80 µL of DPPH solution was pipetted into lines A to line G and as much as 130 µL of methanol as a blank was pipetted to line H. The mixture on 96-wells microplate was then incubated for 30 minutes in a dark room [23]. Then, the absorbance of the solutions was measured with a 96-wells microplate reader using a 520 nm filter [24], and the percentage of free radical scavenging activity (percentage of inhibition) was calculated based on equation 1 below.

$$\text{Inhibition(\%)} = \frac{(A_{\text{control}} - A_{\text{sample}})}{A_{\text{control}}} \times 100\% \quad (1)$$

Inhibition Concentration 50 (IC₅₀) value is calculated based on a linear regression equation (*y* = *ax* + *b*) obtained from the curve by plotting the value of Ln concentration on the x-axis and the percentage inhibition value on the y-axis. Antioxidant activity is categorized as very strong if the IC₅₀ value < 50 µg/mL, strong if the IC₅₀ value is between 50–100 µg/mL, while if the IC₅₀ value is between 101–250 µg/mL, it is weak if the IC₅₀ value is between 251–500 µg/mL, and not active as an antioxidant if the IC₅₀ value > 500 µg/mL [25].

2.4. *In vitro* sunscreen activity test

2.4.1. Determination of %Te and %Tp values

Determination of %Te and %Tp values were conducted by modifying the previous method [7]. A total of 10 mg of sample was dissolved in 10 mL absolute ethanol to obtain the mother liquor with a concentration of 1000 µg/mL. Then the mother liquor was diluted to obtain a 200 µg/mL test solution. As much as of 120, 90, 60, 30, and 15 µL of the 200 µg/mL test solution were pipetted successively into the wells in lines A, B, C, D, and E using multichannel micropipettes. Then as much as 30, 60, 90, and 105 µL absolute ethanol was pipetted into wells in rows B, C, D, and E using a multichannel micropipette. In order to obtain a sample concentration of 200 µg/mL (row A), 150 µg/mL (line B), 100 µg/mL (line C), 50 µg/mL (line D), 25 µg/mL (line F). The benzophenone-3 solutions as standard sunscreen were prepared in the same way as the sample solutions. The wells in line F were filled with 120 µL of absolute ethanol as blank. A 96-wells microplate reader measured the absorbance of the solution in each well at the wavelengths that cause erythema (293–318 nm) and pigmentation (323–373 nm) with 5 nm intervals. Then, to calculate the %Te and %Tp values, the absorbance value (*A*) was first converted to the transmittance percentage value (% *T*) using the following equation.

$$A = -\log T \quad (2)$$

Next, the % *T* value obtained is entered into equation 3 to calculate the %Te value and entered into equation 4 to calculate the %Tp value. In this case, *F_e* is the erythema flux value, and *F_p* is the pigmentation flux. *F_e* and *F_p* values are constants and have been explained in previous literature [26].

$$\%Te = \frac{\sum(T \times Fe)}{\sum Fe} \quad (3)$$

$$\%Tp = \frac{\sum(T \times Fp)}{\sum Fp} \quad (4)$$

Sunscreen is categorized as a sunblock if it has %Te <1% and %Tp of 3–40%, extra protection if it has %Te of 1–6% and %Tp of 42–86%, standard suntan if it has %Te of 6–12% and %Tp of 45–86%, fast tanning if it has %Te of 10–18% and %Tp of 45–86% [26].

2.4.2. Determination of SPF value

The determination of the SPF value was performed by modifying the previous method [7]. A total of 10 mg of sample was dissolved in 10 mL absolute ethanol to obtain the mother liquor with a concentration of 1000 µg/mL. Then the mother liquor was diluted to obtain a 200 µg/mL test solution. As much as of 120, 90, 60, 30, and 15 µL of the 200 µg/mL test solution were pipetted successively into the wells in lines A, B, C, D, and E using multichannel micropipettes. Then, as much as 30, 60, 90 and 105 µL absolute ethanol is pipetted into wells in rows B, C, D, and E using a multichannel micropipette, so that the concentration of the test solution obtained in row A (200 µg/mL), B (150 µg/mL), C (100 µg/mL), D (50 µg/mL), and E (25 µg/mL). The benzophenone-3 solutions as standard sunscreen were prepared in the same way as the sample solutions. The wells in line F were filled with 120 µL of absolute ethanol as blank. A 96-wells microplate reader measured the absorbance of the solution in each well at the wavelength range of 290–320 nm. The absorbance value of the sample at each test concentration was recorded, and then the SPF value was calculated using the Dutra equation [27].

$$SPF = CF \times \sum_{290}^{320} EE(\lambda) \times I(\lambda) \times Abs(\lambda) \quad (5)$$

In this case, CF is a correction factor of 10. EE is Erythral Effect, and I is the intensity of the sun. The value of EE x I is a constant, as explained elsewhere [27]. Then Abs is the absorbance of the sample after deducting the absorbance of the blank. The protective strength of a sunscreen is categorized as ultra-protection if the SPF value is ≥ 15, the maximum protection is if the SPF value is 8–≤ 15, extra protection is if the SPF value is 6–≤ 8, moderate protection is if the SPF value is 4–<6 and minimal protection is if the SPF value is 2–<4 [28].

2.4.3. Determination of antibacterial activity by the disk diffusion method

The antibacterial activity test was carried out by modifying the previous procedure [29, 30]. Before testing, the equipment has been sterilized, and the bacteria used were also rejuvenated. The bacterial suspension was made in a physiological NaCl solution with a %T of 25% at a wavelength of 580 nm. A total of 10 mg of chalcone analog was weighed and dissolved in 10 mL DMSO, then diluted to obtain test solutions with concentrations of 120, 60, and 30 µg/disk. A total of 300 µL of bacterial suspension was put into a petri dish and

added with 15 mL of NA media, then homogenized and then allowed to stand until it solidified. As much as 10 µL of each concentration of test solutions was pipetted and dropped on each disk paper. Then, the disk paper was placed on a compacted inoculum media and incubated for 24 hours at 37°C by turning the petri dish upside down. After 24 hours, the formed inhibition zone around the disk was measured. The 10 µL of DMSO was used as a negative control, and 30 µg chloramphenicol antibiotic disk was used as a positive control. The strength of the sample antibacterial activity was expressed as the percentage of the inhibitory power by comparing the sample inhibitory diameter with the inhibitory diameter of the positive control following equation 6 below.

$$\text{inhibitory power (\%)} = \frac{\text{Inhibitory diameter of the sample}}{\text{Inhibitory diameter of the positive control}} \times 100\% \quad (6)$$

Antibacterial activity is categorized as strong if the percentage of inhibition power ≥ 70%, as moderate if the percentage of inhibition ranges from 50–70% and as weak if the percentage of inhibition < 50% [31].

3. Results and Discussion

3.1. Chalcone analog synthesis

The chalcone analog compound, (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one has been successfully synthesized by the microwave irradiation method through the Claisen-Schmidt condensation reaction between 4'-hydroxy acetophenone and 4-methoxy benzaldehyde. The microwave irradiation method was chosen because it has various advantages, such as faster reaction time, more comfortable to control, more energy-efficient, and can increase product yield, but not under all reaction conditions [32]. In this study, the use of the microwave irradiation method was proven to accelerate reaction times and increase the yield of synthesized compounds when compared to the literature [15, 17]. However, the yield obtained in this study is slightly lower when compared to the literature [21], as can be seen in Table 1.

Table 1. Comparison of the synthesis results of compound (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one by conventional methods from several references and microwave irradiation methods

Synthesis Method	Reaction Conditions	Yield (%)	Melting Point (°C)	Ref.
Stirrer	Room Temperature	86	182	[17]
		86	180–182	[15]
		90	184–185	[21]
Microwave irradiation	180 W power	89	184–185	This work

Based on Table 1, it can be observed that the melting point of the synthesized compound is reached at a temperature of 184–185°C. The melting point range of ≤ 2°C indicates that the solid product obtained was pure.

This conclusion was supported by a TLC chromatogram that showed the single spot with R_f 0.45 and HPLC chromatogram, which also showed a single peak at a retention time of 11.38 minutes.

The structure of synthesized chalcone analog was confirmed based on the analysis of UV, FT-IR, ^1H NMR, and HRMS spectrum. The UV spectrum of chalcone analog shows absorption at wavelengths of 203, 235, and 345 nm. This absorption indicates the characteristics of conjugated double bonds found in the conjugation system of benzoyl and cinnamoyl of chalcone analog. The presence of a hydroxy group (OH) in chalcone analog is confirmed by the appearance of a broad absorption band in the FT-IR spectrum, at a wavenumber of 3162 cm^{-1} which is supported by the appearance of a broad singlet peak (δ 9.23 ppm) with the integration of one proton in the ^1H NMR spectrum. The presence of a methoxy group (OCH_3) is confirmed by the appearance of several absorption bands in the FT-IR spectrum at wavenumbers of $2975\text{--}2908\text{ cm}^{-1}$ (aliphatic C-H) and 1037 cm^{-1} (C-O), which is supported by the appearance of a singlet peak in the ^1H NMR spectrum with the integration of three protons at δ 3.87 ppm. In addition, the presence of a carbonyl group ($\text{C}=\text{O}$) of ketone that conjugated with an aromatic ring A and with α , β double bond (keto ethylene group) is confirmed by the appearance of the absorption band at the wavenumber of 1643 cm^{-1} . Then, the presence of phenyl groups is confirmed by the appearance of absorption bands at wavenumbers of 1601, 1508, and 1466 cm^{-1} (aromatic C=C) and 3089 cm^{-1} (aromatic C-H and alkene C-H) in the FT-IR spectrum, which is supported by the appearance of peaks in the ^1H NMR spectrum with a total integration of 10 protons in the chemical shift area of 8.01–6.97 ppm, as can be seen in Figure 3.

Figure 3. ^1H NMR spectrum (acetone- d_6 , 500 MHz) of chalcone analog (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one

Generally, the signals of α and β protons appear as two doublet signals (1H), with a coupling constant of 15–16 Hz. Nevertheless, in this case, the signals of protons α and β appear as a singlet (2H) signal. This can be caused by the effect of the solvent used in the measurement of the spectrum so that the chemical environment of α and β

protons in chalcone analog becomes equivalent. As a result, the spectrum does not show a spin splitting, even though the protons are not equivalent, structurally. When measured with polysol solvent [20], there is a slight spin splitting between α and β protons because the chemical environment becomes slightly different, but the 3'/5' and 3/5 proton signals were overlapped and appear as a doublet (4H) signal. When measured with the CDCl_3 solvent [22], the difference in the chemical environment of α and β protons becomes greater, as a result, the two doublet signals become further apart, but the 3'/5' and 3/5 proton signals still overlap, so that they appear as a doublet (4H) signals. Another difference can be observed from the hydroxy (OH) proton signal. In the acetone- d_6 solvent, the hydroxy proton signal appears at a chemical shift of 9.23 ppm as a broad singlet. However, when measured in polysol and CDCl_3 solvents, the hydroxy proton signal appears as a singlet at a chemical shift of 10.10 ppm and 5.85 ppm, respectively. Comparison of ^1H NMR spectrum data of synthesized chalcone analog as measured in an acetone- d_6 with ^1H NMR spectrum data of reference compounds measured in polysol and CDCl_3 has been shown in section 2.2. Then the comparison of the three ^1H NMR spectra of this compound in the aromatic chemical shift region can be seen in Supplementary Materials. Based on the comparison, it can be concluded that the type of solvent used in the ^1H NMR analysis can affect the chemical shift of protons of chalcone analog.

Then, a high-resolution mass spectrum measurement is performed to confirm the molecular mass of the synthesized chalcone analog. Based on the results of mass spectrum analysis, the molecular ion peak $[\text{M}+\text{H}^+]$ of chalcone analog in the MS spectrum was found at m/z 255.1010 with a mass difference of 0.0011 from the calculated mass. This small difference in mass shows that the compound obtained has excellent purity. Thus, the overall results of the spectroscopic analysis showed that the compound obtained had the structure as expected.

3.2. Biological Activity Assay

3.2.1. Measurement of Antioxidant Activity by DPPH Method

In this study, the measurement of antioxidant activity was carried out by the DPPH method. This method has several advantages, including being more accessible and cheaper compared to other antioxidant activity testing methods. Therefore, this method is widely used to predict the potential antioxidant activity of a pure compound or plant extract. The activity of capturing free radicals by chalcone analog and ascorbic acid as a standard antioxidant can be observed visually through the color changes that occurred in the test solution. DPPH solution has maximum absorbance in the visible light region, which is in the wavelength range of 515–520 nm. When the sample stabilizes a dark purple DPPH free radical through a proton or electron donor mechanism, the DPPH radical will be reduced to its stable form,

namely 1,1-diphenyl-2-picrylhydrazine (DPPH-H) which is yellow [22], as can be seen in Figure 4.

Figure 4. Color changes in antioxidant activity test with the DPPH method

The results of the antioxidant activity assay of chalcone analog by the DPPH method can be seen in Table 2. Based on these results, the higher the test concentration, the higher the DPPH free radical scavenging activity. A hydroxy-substituted chalcone analog can donate its hydroxyl proton to the DPPH free radical because the conjugation system in this compound will direct resonance into the aromatic ring. Then, the presence of electron-donating groups such as methoxy in this compound can also increase the stability of produced aryloxy radicals through electron delocalization [33]. The mechanism of proton donation of chalcone analog to DPPH free radical can be seen in Figure 5.

Based on the calculation of IC_{50} values, the antioxidant activity of this chalcone analog is included in the weak category [25], with an IC_{50} value of 300.43 $\mu\text{g/mL}$. In contrast, ascorbic acid as a standard antioxidant shows extreme free radical scavenging

activity, with an IC_{50} value of 9.45 $\mu\text{g/mL}$. This is assumed because the chalcone analog has only one proton that can be donated to DPPH free radical, the hydroxyl proton which is located in the *para* position of the ring A, while ascorbic acid can donate two protons rapidly to DPPH free radicals [34] as can be seen in Figure 6.

Table 2. DPPH test result of chalcone analog

Compounds	Concentration ($\mu\text{g/mL}$)	Inhibition (%)	IC_{50} ($\mu\text{g/mL}$)
Chalcone analog	1000	77.61 $\pm$ 1.84	300.43
	500	59.61 $\pm$ 1.71	
	250	46.29 $\pm$ 0.62	
	125	30.09 $\pm$ 0.45	
	62.5	15.62 $\pm$ 0.66	
	31.25	1.44 $\pm$ 0.87	
Ascorbic acid	100	98.63 $\pm$ 0.12	9.45
	50	85.03 $\pm$ 0.76	
	25	69.83 $\pm$ 0.76	
	12.5	57.24 $\pm$ 1.14	
	6.25	40.39 $\pm$ 1.69	
	3.125	27.07 $\pm$ 0.45	

Some literature also supports this explanation that the phenolic compound antioxidant activity was affected by the position of the hydroxyl group and also by the number of hydroxyl groups attached to the aromatic ring. The more hydroxyl groups that are bound to a phenolic compound will increase the ability of the compound to donate its protons to DPPH free radicals, thereby increasing the potential for antioxidant activity [35]. For example, compound (*E*)-1-(2,4-dihydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one containing two hydroxy substituents in the aromatic ring has a stronger antioxidant activity (IC_{50} = 47.45 μM) [36].

Figure 5. Proton donor mechanisms of chalcone analog to DPPH free radicals

Figure 6. Proton donor mechanisms of ascorbic acid to DPPH free radicals

3.2.2. *In vitro* Sunscreen Activity Test

Sunscreen activity testing was carried out *in vitro* using a 96-wells microplate reader. This test includes determining the %Te, %Tp, and SPF values. The %Te value represents the ability of a molecule to protect the skin from UV B rays (290–320 nm), which are transmitted and can cause redness of the skin (erythema). While the %Tp value illustrates the ability of a molecule to protect the skin from UV A rays (320–375 nm), which are passed and can cause darkening of the skin (pigmentation) [37]. The lower the %Te and %Tp values of a substance indicates that the smaller the intensity of UV rays transmitted to the skin. In other words, the better the potential of the substance acts as a UV B or UV A filter.

Table 3. Results of determining the %Te and %Tp values of chalcone analog in various concentrations

Compounds	Concentration (µg/mL)	%Te values	UV B Filter Categories	%Tp values	UV A Filter Categories
Chalcone analog	200	0.19±0.01	Sunblock	0.05±0.00	Sunblock
	150	0.73±0.10	Sunblock	0.07±0.00	Sunblock
	100	3.29±0.19	Extra protection	0.27±0.12	Sunblock
	50	21.74±1.40	-	4.03±1.68	Sunblock
	25	42.78±0.13	-	16.24±0.63	Sunblock
Benzophenone-3	200	0.30±0.01	Sunblock	9.72±0.58	Sunblock
	150	0.85±0.03	Sunblock	13.04±0.23	Sunblock
	100	5.73±0.66	Extra protection	22.82±0.23	Sunblock
	50	21.09±2.04	-	39.93±2.58	Sunblock
	25	44.49±1.99	-	60.97±3.05	Fast tanning

Based on sunscreen activity test, as can be seen in Table 3, the results show that at the lowest test concentration (25 µg/mL), chalcone analog, (*E*)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one has been able to provide proper protection from UV A rays, but it has not been able to protect from UV B rays. The chalcone analog can only provide proper protection from UV A and UV B rays at a test concentration of 150 µg/mL with %Te and %Tp values of 0.73±0.10 and 0.07±0.00 %, respectively. At these concentrations, the protective strength by chalcone analog can be categorized as sunblock [26]. When compared with a standard sunscreen compound (benzophenone-3 or oxybenzone), the chalcone analog proved to be slightly better in acting as a UV B filter, which is shown by the %Te value which is slightly lower than the %Te value of benzophenone-3 at a test concentration of 100–200 µg/mL. The chalcone analog also possessed better potential as UV A filter. Because, at the lowest test concentration (25 µg/mL), the chalcone analog can already act as UV A filter in the sunblock category, while benzophenone-3 at the same concentration still causes the fast tanning effect. This is caused by the presence of the conjugation system of *p*-methoxy cinnamoyl in chalcone analog. The presence of a

more extended conjugation system causes the chalcone analog to have a better ability to absorb UV light at higher wavelengths compared to benzophenone-3, so that, the intensity of UV A light transmitted to the skin (%Te) is lower compared to benzophenone-3. Therefore, the chalcone analog can act better as a UV A filter.

Then, the SPF values measurement is performed to determine the ability of a sunscreen to protect the skin from sun exposure. The SPF value states how many times the natural endurance of a person's skin is doubled so that it is safe in the sun without getting burnt [38]. Based on measurements of the SPF values, as can be seen in Table 4. The results show that at a concentration of 100 µg/mL, the chalcone analog was able to provide ultra-protection as indicated by an SPF value ≥ 15 [29], while the benzophenone-3 can provide protection ultra at concentrations above 100 µg/mL. Thus, it can be concluded that the chalcone analog has better potential in protecting the skin from UV B and UV A radiation compared to benzophenone-3.

Table 4. Results of determining SPF values of chalcone analog in various concentrations

Compounds	Concentrations (µg/mL)	SPF values	Protection categories
Chalcone analog	200	29.60±0.55	Ultra
	150	21.10±1.46	Ultra
	100	15.83±2.28	Ultra
	50	7.99±0.66	Extra
	25	3.94±0.15	Minimum
Benzophenone-3	200	26.71±0.31	Ultra
	150	19.98±0.52	Ultra
	100	12.10±1.44	Maximum
	50	8.49±1.95	Maximum
	25	2.85±0.50	Minimum

3.2.3. Antibacterial Activity Test with Disk Diffusion Method

The results of the antibacterial activity test of the chalcone analog, (*E*)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one with the disk diffusion method can be seen in Figure 7 and Table 5. Based on these results, it can be observed that at test concentrations of 30, 60, and 120 µg/disk, this compound only shows antibacterial activity against Gram-positive bacteria, *Staphylococcus aureus* but does not show inhibitory zones against Gram-negative bacteria, *Escherichia coli*.

As presented in Table 5, it can be observed that the higher the concentration of the test, the greater the diameter of the inhibition zone against Gram-positive bacteria, *Staphylococcus aureus*. However, based on literature [31], the antibacterial activity produced at all concentrations of the test is still relatively weak, with a percentage of inhibition <50% compared to the inhibition produced by chloramphenicol as a positive control.

Figure 7. Results of antibacterial activity test of chalcone against (a) *Staphylococcus aureus*, (b) *Escherichia coli*

Table 5. Results for antibacterial activity test of chalcone analog

Bacteria	Treatments	Inhibition zone diameters (mm)				Inhibition powers (%)
		1	2	3	Average	
<i>Staphylococcus aureus</i>	Positive control	20.2	20.9	19.7	20.3±0.6	100
	Negative control	0	0	0	0	0
	30 µg/disk	7	6.5	6.5	6.7±0.3	33
	60 µg/disk	7.3	7	7	7.1±0.2	35
	120 µg/disk	7.5	7.5	7.5	7.5±0.0	37
<i>Escherichia coli</i>	Positive control	20.3	20.1	19.6	20.0±0.4	100
	Negative control	0	0	0	0	0
	30 µg/disk	0	0	0	0	0
	60 µg/disk	0	0	0	0	0
	120 µg/disk	0	0	0	0	0

Based on previous research on the antibacterial mechanism of another chalcone analog with a similar structure (hydroxy substituent located in the *meta* position), the chalcone analog exhibits antibacterial mechanisms by damaging the cell wall of *Staphylococcus aureus* bacteria [39]. To damage the bacterial cell wall, a compound must be able to interact with bacterial cell wall components. Based on the literature, the cell wall structure of Gram-negative bacteria is more complex and dynamic than Gram-positive bacteria. The cell wall of Gram-negative bacteria has an outer membrane above a thin layer of peptidoglycan. This structure causes Gram-negative bacteria to have a strong, robust, and elastic cell wall [40]. Therefore, the molecule size, the capacity of the electron-withdrawing and donating groups, and the hydrophilicity of a molecule will determine its antibacterial activity. Besides, the antibacterial activity of chalcone analog is also affected by the hydrophobicity of the bacterial cell surface. Based on previous research, chalcone analog compound with a hydroxy substituent in the *meta* position provide better antibacterial activity against bacteria with more hydrophobic cell surfaces. According to the study, the surface of *Staphylococcus aureus* cells is more hydrophobic compared to *Escherichia coli* [39]. This might cause chalcone analog (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-

one only shows inhibition against *Staphylococcus aureus* but does not show inhibition against *Escherichia coli* at the test concentrations.

4. Conclusion

In this study, chalcone analog (E)-1-(4-hydroxyphenyl)-3-(4-methoxyphenyl)prop-2-en-1-one has been successfully synthesized using a microwave irradiation method with shorter time and more yield high compared to some previous synthesis methods. Based on bioactivity testing that has been done, the chalcone analog compound shows a better potential for sunscreen activity compared to the standard sunscreen, benzophenone-3. However, the compound has no potential as an antioxidant and antibacterial. Further studies are needed to determine the stability of this compound under UV radiation and also its toxicity to human skin.

Acknowledgment

The authors would like to thank DPRM-Ministry of Research Technology and Higher Education (KEMENRISTEK DIKTI) for funding provided for this research through the Hibah Penelitian Dosen Pemula (PDP) 2019 with the research contract number 06.15.LP2M.STIFAR.V.2019. The authors also thank Prof. Dr. Adel Zamri, MS, DEA for the synthesis facility support that has been provided at the Organic Synthesis Laboratory, FMIPA, University of Riau.

References

- [1] N. K. Sahu, S. S. Balbhadra, J. Choudhary and D. V. Kohli, Exploring Pharmacological Significance of Chalcone Scaffold: A Review, *Current Medicinal Chemistry*, 19, 2, (2012), 209–225
<http://dx.doi.org/10.2174/092986712803414132>
- [2] Zhou Tiezhong, Xu Ming Deng and Jia Zhang Qiu, Antimicrobial activity of licochalcone E against *Staphylococcus aureus* and its impact on the production of staphylococcal alpha-toxin, *Journal of microbiology and biotechnology*, 22, 6, (2012), 800–805 <https://doi.org/10.4014/jmb.1112.12020>
- [3] Ismiyarto Ismiyarto, Suyanti Suyanti, Ngadiwiyan Ngadiwiyan, Purbowatiningrum Ria Sarjono and Nor Basid Adiwibawa Prasetya, Synthesis of 4-Hydroxy-2-Methylchalcone from meta-Cresol Formulation Product and Its Activities as an Antibacteria, *Jurnal Kimia Sains dan Aplikasi*, 21, 4, (2018), 193–197
<https://doi.org/10.14710/jksa.21.4.193-197>
- [4] Florian George and Toufik Fellague, Compose Chimique De Derive Azote De Chalcone, Procede D'obtention D'un Tel Compose Chimique, Composition Cosmetique Et/Ou Ftlttrante Contenant Un Tel Compose Chimique, in: O.M.d.l.P. Intellectuelle (Ed.), France, 2003, pp. 38
- [5] Serge Forestier, Claudine Moire and Gerard Lang, Cosmetic composition containing hydroxylated chalcone derivatives and its use for protecting the skin and the hair against luminous radiations, new

- hydroxylated chalcone derivatives employed and process for their preparation, in: U.S.P. DOCUMENTS (Ed.), L'Oreal, Paris, France, United State, 1989,
- [6] Silvina Quintana Lazópulos, Federico Svarc, Gabriel Sagrera and Lelia Dixelio, Absorption and Photo-Stability of Substituted Dibenzoylmethanes and Chalcones as UVA Filters, *Cosmetics*, 5, 2, (2018), <https://doi.org/10.3390/cosmetics5020033>
 - [7] Ihsan Ikhtiarudin, Nesa Agistia, Tria Harlianti and Adel Zamri, Sintesis dan Potensi Aktivitas Tabir Surya Senyawa Analog Kalkon Turunan 3'-Hidroksiasetofenon dan 4-Metoksibenzaldehid, *Jurnal Photon*, 10, 1, (2019), 1-12
 - [8] Ihsan Ikhtiarudin, Lelani, Adel Zamri, Hilwan Yuda Teruna and Yuharmen, Sintesis dan Uji Toksisitas Senyawa Analog Kalkon Turunan 2'-Hidroksiasetofenon dan Halobenzaldehid, *Jurnal Photon*, 5, 1, (2014), 57-63
 - [9] Ihsan Ikhtiarudin, Neni Frimayanti, Hilwan Y Teruna and Adel Zamri, Microwave-assisted synthesis, molecular docking study and in vitro evaluation of halogen substituted flavonols against P388 Murine leukemia cells, *Applied Science and Technology*, 1, 1, (2017), 375-381
 - [10] Neni Frimayanti, Jasril Jasril, Ihsan Ikhtiarudin, Syilfia Hasti and Anisa Indah Reza, Microwave-Assisted Synthesis, in Silico Studies and in Vivo Evaluation for Antidiabetic Activity of New Brominated Pyrazoline Analogues, *Thai Journal of Pharmaceutical Sciences (TJPS)*, 43, 2, (2019), 83-89
 - [11] Nora M. Rateb and Hussein F. Zohdi, Atom-Efficient, Solvent-Free, Green Synthesis of Chalcones by Grinding, *Synthetic Communications*, 39, 15, (2009), 2789-2794
<https://doi.org/10.1080/00397910802664244>
 - [12] Jae-Chul Jung, Yongnam Lee, Dongguk Min, Mankil Jung and Seikwan Oh, Practical Synthesis of Chalcone Derivatives and Their Biological Activities, *Molecules*, 22, 11, (2017),
<https://doi.org/10.3390/molecules22111872>
 - [13] Edmont V. Stoyanov, Yves Champavier, Alain Simon and Jean-Philippe Basly, Efficient liquid-Phase synthesis of 2'-Hydroxychalcones, *Bioorganic & Medicinal Chemistry Letters*, 12, 19, (2002), 2685-2687
[https://doi.org/10.1016/S0960-894X\(02\)00553-X](https://doi.org/10.1016/S0960-894X(02)00553-X)
 - [14] Tahseen Razzaq and C. Oliver Kappe, On the Energy Efficiency of Microwave-Assisted Organic Reactions, *ChemSusChem*, 1, 1-2, (2008), 123-132
<https://doi.org/10.1002/cssc.200700036>
 - [15] Jonathan R. Dimmock, N. Murthi Kandepu, Mark Hetherington, J. Wilson Quail, Uma Pugazhenth, Athena M. Sudom, Mahmood Chamankhah, Patricia Rose, Eric Pass, Theresa M. Allen, Sarah Halleran, Jen Szydłowski, Bulent Mutus, Marie Tannous, Elias K. Manavathu, Timothy G. Myers, Erik De Clercq and Jan Balzarini, Cytotoxic Activities of Mannich Bases of Chalcones and Related Compounds, *Journal of Medicinal Chemistry*, 41, 7, (1998), 1014-1026
<https://doi.org/10.1021/jm970432t>
 - [16] S Sathiya Moorthi, K Chinnakali, S Nanjundan, P Selvam, H-K Fun and X-L Yu, 3-(3-Hydroxyphenyl)-1-(4-methoxyphenyl) prop-2-en-1-one, *Acta Crystallographica Section E: Crystallographic Communications*, 61, 3, (2005), o743-o745
<https://doi.org/10.1107/S1600536805004915>
 - [17] Halise I. Gul, Kadir O. Yerdelen, Mustafa Gul, Umashankar Das, Bulbul Pandit, Pui-Kai Li, Hasan Secen and Fikretin Sahin, Synthesis of 4'-Hydroxy-3'-piperidinomethylchalcone Derivatives and Their Cytotoxicity Against PC-3 Cell Lines, *Archiv der Pharmazie*, 340, 4, (2007), 195-201
<https://doi.org/10.1002/ardp.200600072>
 - [18] Halise Inci Gul, Murat Cizmecioglu, Sevil Zencir, Mustafa Gul, Pakize Canturk, Mustafa Atalay and Zeki Topcu, Cytotoxic activity of 4'-hydroxychalcone derivatives against Jurkat cells and their effects on mammalian DNA topoisomerase I, *Journal of Enzyme Inhibition and Medicinal Chemistry*, 24, 3, (2009), 804-807
<https://doi.org/10.1080/14756360802399126>
 - [19] Daniela I. Batovska and Iva Todorova Todorova, Trends in Utilization of the Pharmacological Potential of Chalcones, *Current Clinical Pharmacology*, 5, 1, (2010), 1-29
<https://doi.org/10.2174/157488410790410579>
 - [20] Bio-Rad Laboratories Inc., SpectraBase; SpectraBase Compound ID=1PXRd97MNom SpectraBase Spectrum ID=8HbcjN7xboX, in, 2020,
 - [21] Ram Pratap, Mavurapu Satyanarayana, Chandeshwar Nath, Ram Raghubir, Anju Puri, Ramesh Chander, Priti Tiwari, Brajendra Kumar Tripathi and Arvind Kumar Srivastava, Oxy Substituted Chalcones as Anthyperglycemic and Antidyslipidemic agents, in, United State, 2006, pp. 19
 - [22] Philip Molyneux, The use of the stable free radical diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity, *Songklanakarinn Journal of Science and Technology*, 26, 2, (2004), 211-219
 - [23] Yueting Lu, Rudi Hendra, Aaron J. Oakley and Paul A. Keller, Efficient synthesis and antioxidant activity of coelenterazine analogues, *Tetrahedron Letters*, 55, 45, (2014), 6212-6215
<https://doi.org/10.1016/j.tetlet.2014.09.065>
 - [24] L. R. Fukumoto and G. Mazza, Assessing Antioxidant and Prooxidant Activities of Phenolic Compounds, *Journal of Agricultural and Food Chemistry*, 48, 8, (2000), 3597-3604
<https://doi.org/10.1021/jf000220w>
 - [25] MR Marjoni and A Zulfisa, Antioxidant activity of methanol extract/fractions of senggani leaves (*Melastoma candidum* D. Don), *Pharmaceutica Analytica Acta*, 8, 8, (2017), 1-6
<https://doi.org/10.4172/2153-2435.1000557>
 - [26] M.S. Balsam and E. Sagarin, *Cosmetics, Science and Technology*, Krieger Publishing Company, 1992
 - [27] Elizângela Abreu Dutra, Daniella Almança Gonçalves da Costa e Oliveira, Erika Rosa Maria Kedor-Hackmann and Maria Inês Rocha Miritello Santoro, Determination of sun protection factor (SPF) of sunscreens by ultraviolet spectrophotometry, *Revista Brasileira de Ciências Farmacêuticas*, 40, (2004), 381-

- 385
<https://doi.org/10.1590/S1516-93322004000300014>
- [28]Fitriyanti Jumaetri Sami, Syamsu Nur and Megawati M Martani, Uji Aktivitas Tabir Surya pada Beberapa Spesies dari Family Zingiberaceae dengan Metode Spektrofotometri, *As-Syifaa Jurnal Farmasi*, 7, 2, (2015), 164-173
- [29]Vivi Anggia, Amri Bakhtiar and Dayar Arbain, The Isolation of xanthonenes from trunk latex of *Garcinia mangostana* Linn. and their antimicrobial activities, *Indonesian Journal of Chemistry*, 15, 2, (2015), 187-193
<https://doi.org/10.22146/ijc.21213>
- [30]Melzi Octaviani, Haiyul Fadhli and Erenda Yuneistya, Uji Aktivitas Antimikroba Ekstrak Etanol Kulit Bawang Merah (*Allium cepa* L.) dengan Metode Difusi Cakram, *Pharmaceutical Sciences & Research*, 6, 1, (2019), 62-68 <https://doi.org/10.7454/psr.v6i1.4333>
- [31]E. W. C. Chan, Y. Y. Lim and Mohammed Omar, Antioxidant and antibacterial activity of leaves of *Etlingera* species (Zingiberaceae) in Peninsular Malaysia, *Food Chemistry*, 104, 4, (2007), 1586-1593
<https://doi.org/10.1016/j.foodchem.2007.03.023>
- [32]Adel Zamri, Hilwan Yuda Teruna and Ihsan Ikhtiarudin, The influences of power variations on selectivity of synthesis reaction of 2'-hydroxychalcone analogue under microwave irradiation, *Molekul*, 11, 2, (2016), 299-307
<http://dx.doi.org/10.20884/1.jm.2016.11.2.220>
- [33]Sindhu Mathew, T. Emilia Abraham and Zainul Akmar Zakaria, Reactivity of phenolic compounds towards free radicals under in vitro conditions, *Journal of Food Science and Technology*, 52, 9, (2015), 5790-5798
<https://doi.org/10.1007/s13197-014-1704-0>
- [34]Satish Balasaheb Nimse and Dilipkumar Pal, Free radicals, natural antioxidants, and their reaction mechanisms, *RSC Advances*, 5, 35, (2015), 27986-28006 <https://doi.org/10.1039/C4RA13315C>
- [35]E. Bendary, R. R. Francis, H. M. G. Ali, M. I. Sarwat and S. El Hady, Antioxidant and structure-activity relationships (SARs) of some phenolic and anilines compounds, *Annals of Agricultural Sciences*, 58, 2, (2013), 173-181
<https://doi.org/10.1016/j.aos.2013.07.002>
- [36]Yogesh Murti, A Goswami and Pradeep Mishra, Synthesis and antioxidant activity of some chalcones and flavanoids, *International Journal of PharmTech Research*, 5, 2, (2013), 811-818
- [37]Siti Hasanah, Islamudin Ahmad and Laode Rijai, Profil Tabir Surya Ekstrak dan Fraksi Daun Pidada Merah (*Sonneratia caseolaris* L.), *Jurnal Sains dan Kesehatan*, 1, 4, (2015), 175-180
<https://doi.org/10.25026/jsk.v1i4.36>
- [38]Marvin E Wala, Suryantom Edy and Defny S Wewengkang, Aktivitas Antioksidan Dan Tabir Surya Fraksi Dari Ekstrak Lamun (*Syringodium isoetifolium*), *Pharmacon*, 4, 4, (2015), 282-289
- [39]Ponnurengam Malliappan Sivakumar, Sobana Priya and Mukesh Doble, Synthesis, Biological Evaluation, Mechanism of Action and Quantitative Structure-Activity Relationship Studies of Chalcones as Antibacterial Agents, *Chemical Biology & Drug Design*, 73, 4, (2009), 403-415 <https://doi.org/10.1111/j.1747-0285.2009.00793.x>
- [40]Terry J. Beveridge, Structures of Gram-Negative Cell Walls and Their Derived Membrane Vesicles, *Journal of Bacteriology*, 181, 16, (1999), 4725